

Validation of ‘sasLM’ Package

Kyun-Seop Bae MD PhD

2022-03-21 12:08:04

Contents

1	Tested Version and Books used for the Validation	5
1.1	Packages Used	5
1.2	Books and Articles used for the Test	5
2	ARS20-8	6
2.1	p8	6
2.2	p42	6
2.3	p101	8
3	Snee EMS ANOVA 1974	11
4	Goodnight	13
4.1	Type I SS	13
4.2	Type II SS	18
4.3	Type III SS	19
5	SAS for Linear Models 4e	22
5.1	Chapter 2	22
5.2	Chapter 3	26
5.3	Chapter 4	31
5.4	Chapter 5	35
5.5	Chapter 6	37
5.6	Chapter 7	41
5.7	Chapter 8	52
5.8	Chapter 11	53

6 Sahai - Unbalanced	69
6.1 Table 11.2	69
6.2 Table 12.6	70
6.3 Table 13.6	70
6.4 Table 14.2	71
6.5 Table 15.3	72
6.6 Table 16.3	74
7 Federer - Variations	76
7.1 Example 1.1	76
7.2 Example 1.2	77
7.3 Example 2.1	78
7.4 Example 2.2	79
7.5 Example 3.1	82
7.6 Example 4.1	90
7.7 Example 5.1	93
7.8 Example 7.1	100
7.9 Example 7.2	101
7.10 Example 7.3	102
7.11 Example 8.1	104
7.12 Example 9.1	105
7.13 Example 9.2	106
7.14 Example 10.1	108
7.15 Example 10.2	111
7.16 Example 11.1	112
7.17 Example 11.2	115
7.18 Example 11.3	120
8 Hinkelmann & Kempthorne - Volume 1	124
8.1 Chapter 6	124
8.2 Chapter 7	125
8.3 Chapter 8	127
8.4 Chapter 9	130
8.5 Chapter 10	135

8.6 Chapter 11	139
8.7 Chapter 12	145
8.8 Chapter 13	148
8.9 Chapter 14	150
9 Hinkelmann & Kempthorne - Volume 2	152
9.1 Chapter 1	152
9.2 Chapter 2	153
9.3 Chapter 6	155
9.4 Chapter 7	157
9.5 Chapter 8	160
9.6 Chapter 9	163
9.7 Chapter 10	167
9.8 Chapter 14	169
9.9 Chapter 16	173
9.10 Chapter 17	178
9.11 Chapter 19	180
10 Lawson - DAE with SAS	183
10.1 Chapter 2	183
10.2 Chapter 3	185
10.3 Chapter 4	192
10.4 Chapter 5	196
10.5 Chapter 7	198
10.6 Chapter 8	201
10.7 Chapter 9	206
10.8 Chapter 11	211
10.9 Chapter 12	214
11 Searle - Linear Models 2e	225
11.1 7.2 (p390, 59%)	225
11.2 7.2 (p393, 60%)	226
12 Web site examples	228
12.1 https://github.com/djnavarro/psyr	228

13 Test Summary **230**

14 Session Information **231**

1 Tested Version and Books used for the Validation

1.1 Packages Used

- ‘sasLM’ version: 0.8.0
- ‘SAS’ version: 9.4 Licensed and University Edition
- ‘car’ version: 3.0.12
- R version: R version 4.1.3 (2022-03-10)

The ‘car’ package is not necessary for ‘sasLM.’ It is used for the comparison of the results.

If you see any difference between ‘car’ and ‘sasLM’, ‘SAS’ results coincide with ‘sasLM’, not with ‘car’.

Before ‘sasLM’ is available on CRAN, you can download using the following command in R.

```
install.packages("sasLM", repos="http://r.acr.kr")
```

1.2 Books and Articles used for the Test

1. Harvey WR. Least-Squares Analysis of Data with Unequal Subclass Frequencies. USDA, Agriculture Research Service, ARS 20-8. 1960. reprinted with corrections as ARS H-4, 1975, also reprinted 1979.
2. Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.
3. Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User’s Group, SAS Institute, Raleigh, N.C. 1976.
4. Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.
5. Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.
6. Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.
7. Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.
8. Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 2 Advanced Experimental Design. John Wiley & Sons Inc. 2005.
9. Lawson J. Design and Analysis of Experiments with SAS. Taylor and Francis Group. 2010.
10. Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

2 ARS20-8

Reference

- Harvey WR. Least-Squares Analysis of Data with Unequal Subclass Frequencies. USDA, Agriculture Research Service, ARS 20-8. 1960. reprinted with corrections as ARS H-4, 1975, also reprinted 1979.

2.1 p8

(1) MODEL

```
p8 = read.csv("C:/G/Rt/ANOVA/ARS20-8p8.csv")
p8 = af(p8, c("PigNo", "Ration"))
GLM(Barrow ~ Ration, p8)
```

```
$ANOVA
Response : Barrow
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       2 11.111  5.5556  1.2626 0.3113
RESIDUALS    15 66.000   4.4000
CORRECTED TOTAL 17 77.111

$Fitness
  R-square Coef Var Root MSE Barrow Mean
0.1440922 40.16715 2.097618     5.222222

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration    2 11.111  5.5556  1.2626 0.3113

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration    2 11.111  5.5556  1.2626 0.3113

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration    2 11.111  5.5556  1.2626 0.3113
```

2.2 p42

(2) MODEL

```
p42 = read.csv("C:/G/Rt/ANOVA/ARS20-8p42.csv")
p42 = af(p42, c("Ration", "Pig", "Sire"))
GLM(Y ~ Sire + Ration, p42)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       3 20.819  6.9397  1.7259 0.2075
RESIDUALS    14 56.292  4.0209
CORRECTED TOTAL 17 77.111
```

```
$Fitness
R-square Coef Var Root MSE   Y Mean
0.2699867 38.39764 2.00521 5.222222
```

```
$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Sire      2 11.1111  5.5556  1.3817 0.2834
Ration    1  9.7079  9.7079  2.4144 0.1425
```

```
$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Sire      2 15.6829  7.8414  1.9502 0.1790
Ration    1  9.7079  9.7079  2.4144 0.1425
```

```
$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Sire      2 15.6829  7.8414  1.9502 0.1790
Ration    1  9.7079  9.7079  2.4144 0.1425
```

(3) MODEL

```
GLM(Y ~ Sire + Ration + Sire:Ration, p42)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       5 51.044 10.2089  4.6997 0.01311 *
RESIDUALS    12 26.067  2.1722
CORRECTED TOTAL 17 77.111
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
R-square Coef Var Root MSE   Y Mean
0.6619597 28.22258 1.473846 5.222222
```

```
$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Sire      2 11.1111  5.5556  2.5575 0.118799
Ration    1  9.7079  9.7079  4.4691 0.056129 .
```

```

Sire:Ration 2 30.2255 15.1127 6.9573 0.009859 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Sire       2 15.6829 7.8414 3.6099 0.059238 .  

Ration     1  9.7079  9.7079 4.4691 0.056129 .  

Sire:Ration 2 30.2255 15.1127 6.9573 0.009859 **  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Sire       2 21.0007 10.5004 4.8339 0.028853 *  

Ration     1  3.5919  3.5919 1.6535 0.222736  

Sire:Ration 2 30.2255 15.1127 6.9573 0.009859 **  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

2.3 p101

(4) MODEL

```

p101 = read.csv("C:/G/Rt/ANOVA/ARS20-8p101.csv")
p101 = af(p101, c("Line", "Sire", "Dam", "Steer"))
GLM(Gain ~ Line + Sire + Dam + Line:Dam + Age + Weight, p101)

```

```

$ANOVA
Response : Gain
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      16 2.4972 0.156073 3.0675 0.001364 **
RESIDUALS   48 2.4422 0.050879
CORRECTED TOTAL 64 4.9394
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Gain Mean
0.5055646 9.354136 0.2255642 2.411385

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Line      2 0.38009 0.190046 3.7352 0.03107 *  

Sire      6 0.92634 0.154391 3.0345 0.01347 *  

Dam      2 0.11894 0.059471 1.1689 0.31940

```

```

Line:Dam 4 0.64889 0.162222 3.1884 0.02113 *
Age      1 0.16462 0.164622 3.2356 0.07835 .
Weight    1 0.25828 0.258283 5.0764 0.02886 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

Line     0  

Sire     6 0.95299 0.15883 3.1217 0.01155 *  

Dam      2 0.32039 0.16019 3.1485 0.05190 .  

Line:Dam 4 0.46516 0.11629 2.2856 0.07373 .  

Age      1 0.34830 0.34830 6.8456 0.01185 *  

Weight    1 0.25828 0.25828 5.0764 0.02886 *  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

CAUTION: Singularity Exists !
      Df  Sum Sq Mean Sq F value Pr(>F)  

Line     0  

Sire     6 0.95299 0.15883 3.1217 0.01155 *  

Dam      2 0.12469 0.06234 1.2253 0.30268  

Line:Dam 4 0.46516 0.11629 2.2856 0.07373 .  

Age      1 0.34830 0.34830 6.8456 0.01185 *  

Weight    1 0.25828 0.25828 5.0764 0.02886 *  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(5) MODEL

```
GLM(Gain ~ Sire + Dam + Line:Dam, p101)
```

```

$ANOVA
Response : Gain
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      14 2.0743 0.148162 2.5856 0.006996 **
RESIDUALS   50 2.8651 0.057302
CORRECTED TOTAL 64 4.9394
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Gain Mean
0.4199453 9.927022 0.2393787 2.411385

```

\$`Type I`

```

          Df  Sum Sq  Mean Sq F value Pr(>F)
Sire      8 1.30644 0.163305  2.8499 0.01089 *
Dam       2 0.11894 0.059471  1.0379 0.36172
Dam:Line  4 0.64889 0.162222  2.8310 0.03412 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`:
          Df  Sum Sq  Mean Sq F value Pr(>F)
Sire      6 1.06000 0.176667  3.0831 0.01202 *
Dam       2 0.11894 0.059471  1.0379 0.36172
Dam:Line  4 0.64889 0.162222  2.8310 0.03412 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`:
CAUTION: Singularity Exists !
          Df  Sum Sq  Mean Sq F value Pr(>F)
Sire      6 1.06000 0.176667  3.0831 0.01202 *
Dam       2 0.02569 0.012844  0.2242 0.79999
Dam:Line  4 0.64889 0.162222  2.8310 0.03412 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

3 Snee EMS ANOVA 1974

Reference

- Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.

(6) MODEL

```
Snee = read.csv("C:/G/Rt/ANOVA/Snee_EMS_ANOVA1974.csv")
Snee = af(Snee, c("Machine", "Analyst", "Test", "Day"))
GLM(Y ~ Day/Machine/Analyst/Test, Snee)

$ANOVA
Response : Y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          167 751.27  4.4986
RESIDUALS       0    0.00
CORRECTED TOTAL 167 751.27

$Fitness
R-square Coef Var Root MSE   Y Mean
      1       NA     NA 8.736905

$`Type I`
              Df Sum Sq Mean Sq F value Pr(>F)
Day             41 365.58  8.9166
Day:Machine     42 196.59  4.6807
Day:Machine:Analyst 42 118.80  2.8285
Day:Machine:Analyst:Test 42  70.30  1.6739

$`Type II`
              Df Sum Sq Mean Sq F value Pr(>F)
Day             41 365.58  8.9166
Day:Machine     42 196.59  4.6807
Day:Machine:Analyst 42 118.80  2.8285
Day:Machine:Analyst:Test 42  70.30  1.6739

$`Type III`
              Df Sum Sq Mean Sq F value Pr(>F)
Day             41 359.44  8.7669
Day:Machine     42 199.40  4.7477
Day:Machine:Analyst 42 118.80  2.8285
Day:Machine:Analyst:Test 42  70.30  1.6739
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Day/Machine/Analyst/Test, Snee), type=3, singular.ok=TRUE)
# NOT WORKING
```

4 Goodnight

Reference

- Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User's Group, SAS Institute, Raleigh, N.C. 1976.

4.1 Type I SS

4.1.1 p7

(7) MODEL

```
p7 = read.csv("C:/G/Rt/ANOVA/Goodnight-p7.csv")
p7 = af(p7, c("A", "B"))
GLM(y ~ A + B + A:B, p7)
```

```
$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       3 13.6027 4.5342  2.807 0.1721
RESIDUALS    4  6.4613 1.6153
CORRECTED TOTAL 7 20.0639
```

```
$Fitness
R-square Coef Var Root MSE y Mean
0.6779647 23.22438 1.270954 5.4725
```

```
$`Type I` 
      Df  Sum Sq Mean Sq F value Pr(>F)
A       1 10.8113 10.8113 6.6929 0.06087 .
B       1  1.3122  1.3122 0.8123 0.41839
A:B     1  1.4792  1.4792 0.9157 0.39279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II` 
      Df  Sum Sq Mean Sq F value Pr(>F)
A       1 10.8113 10.8113 6.6929 0.06087 .
B       1  1.3122  1.3122 0.8123 0.41839
A:B     1  1.4792  1.4792 0.9157 0.39279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III` 
      Df  Sum Sq Mean Sq F value Pr(>F)
```

```

A     1 10.8113 10.8113  6.6929 0.06087 .
B     1 1.3122  1.3122  0.8123 0.41839
A:B   1 1.4792  1.4792  0.9157 0.39279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(8) MODEL

```
GLM(y ~ A + A:B + B, p7)
```

```
$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      3 13.6027  4.5342  2.807 0.1721
RESIDUALS  4  6.4613  1.6153
CORRECTED TOTAL 7 20.0639
```

```
$Fitness
R-square Coef Var Root MSE y Mean
0.6779647 23.22438 1.270954 5.4725
```

```
$`Type I`
          Df  Sum Sq Mean Sq F value Pr(>F)
A     1 10.8113 10.8113  6.6929 0.06087 .
A:B   2 2.7914  1.3957  0.8640 0.48764
B     0
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df  Sum Sq Mean Sq F value Pr(>F)
A     1 10.8113 10.8113  6.6929 0.06087 .
A:B   1 1.4792  1.4792  0.9157 0.39279
B     1 1.3122  1.3122  0.8123 0.41839
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df  Sum Sq Mean Sq F value Pr(>F)
A     1 10.8113 10.8113  6.6929 0.06087 .
A:B   1 1.4792  1.4792  0.9157 0.39279
B     1 1.3122  1.3122  0.8123 0.41839
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(9) MODEL

```
GLM(y ~ B + A + A:B, p7)
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		
CORRECTED TOTAL	7	20.0639			

\$Fitness

R-square	Coef	Var	Root	MSE	y	Mean
0.6779647	23.22438	1.270954	5.4725			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(10) MODEL

```
GLM(y ~ B + A:B + A, p7)
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		

CORRECTED TOTAL 7 20.0639

\$Fitness

	R-square	Coef	Var	Root	MSE	y	Mean
	0.6779647	23.22438	1.270954	5.4725			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.4184
B:A	2	12.2905	6.1452	3.8043	0.1187
A	0				

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
B:A	1	1.4792	1.4792	0.9157	0.39279
A	1	10.8113	10.8113	6.6929	0.06087
					.

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
B:A	1	1.4792	1.4792	0.9157	0.39279
A	1	10.8113	10.8113	6.6929	0.06087
					.

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(11) MODEL

GLM(y ~ A:B + A + B, p7)

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		
CORRECTED TOTAL	7	20.0639			

\$Fitness

	R-square	Coef	Var	Root	MSE	y	Mean
	0.6779647	23.22438	1.270954	5.4725			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A:B	3	13.603	4.5342	2.807	0.1721
A	0				

```

B      0

$`Type II`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   1  1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

B     1  1.3122  1.3122  0.8123 0.41839  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   1  1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

B     1  1.3122  1.3122  0.8123 0.41839  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(12) MODEL

```
GLM(y ~ A:B + A + B, p7)
```

```
$ANOVA  

Response : y  

  Df  Sum Sq Mean Sq F value Pr(>F)  

MODEL          3 13.6027  4.5342  2.807 0.1721  

RESIDUALS       4  6.4613  1.6153  

CORRECTED TOTAL 7 20.0639
```

```
$Fitness  

  R-square Coef Var Root MSE y Mean  

  0.6779647 23.22438 1.270954 5.4725
```

```
$`Type I`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   3 13.603  4.5342  2.807 0.1721  

A     0  

B     0
```

```
$`Type II`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   1  1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

B     1  1.3122  1.3122  0.8123 0.41839  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A:B  1 1.4792 1.4792 0.9157 0.39279  

A    1 10.8113 10.8113 6.6929 0.06087 .  

B    1 1.3122 1.3122 0.8123 0.41839  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

4.2 Type II SS

4.2.1 p14

(13) MODEL

```
GLM(y ~ A + B + A:B, p7[-8,]) # p16
```

```
$ANOVA  

Response : y  

  Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      3 12.7672 4.2557 2.0088 0.2906  

RESIDUALS   3  6.3555 2.1185  

CORRECTED TOTAL 6 19.1227
```

```
$Fitness  

  R-square Coef Var Root MSE   y Mean  

0.6676471 27.24211 1.455507 5.342857
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A    1 9.9567 9.9567 4.6999 0.1187  

B    1 1.9225 1.9225 0.9075 0.4111  

A:B  1 0.8880 0.8880 0.4192 0.5635
```

```
$`Type II`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A    1 11.1715 11.1715 5.2733 0.1053  

B    1  1.9225  1.9225 0.9075 0.4111  

A:B  1  0.8880  0.8880 0.4192 0.5635
```

```
$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A    1 9.5258 9.5258 4.4965 0.1241  

B    1 1.3690 1.3690 0.6462 0.4803  

A:B  1 0.8880 0.8880 0.4192 0.5635
```

4.2.2 p24

(14) MODEL

```
p24 = read.csv("C:/G/Rt/ANOVA/Goodnight-p24.csv")
p24 = af(p24, c("A", "B", "C"))
GLM(Y ~ A + B + C, p24) # p27

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       6 45.924  7.6540  9.1615 0.00499 ***
RESIDUALS    7  5.848  0.8354
CORRECTED TOTAL 13 51.772
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
   R-square Coef Var Root MSE  Mean
0.8870405 14.83986 0.9140295 6.159286

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
A 1 4.724  4.7235  5.6538 0.04904 *
B 3 37.998 12.6660 15.1606 0.00191 **
C 2  3.203  1.6013  1.9167 0.21686
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
A 0
B 2 0.4424  0.2212  0.2648 0.7747
C 2 3.2025  1.6013  1.9167 0.2169

$`Type III`
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)
A 0
B 2 0.4424  0.2212  0.2648 0.7747
C 2 3.2026  1.6013  1.9167 0.2169
```

4.3 Type III SS

4.3.1 p27

(15) MODEL

```

p27 = read.csv("C:/G/Rt/ANOVA/Goodnight-p27.csv")
p27 = af(p27, c("A", "B"))
GLM(y ~ A + B + A:B, p27) # p29

```

```

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 128.193 25.6386  53.469 6.77e-05 ***
RESIDUALS   6   2.877  0.4795
CORRECTED TOTAL 11 131.070
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE y Mean
0.9780499 7.413912 0.6924594 9.34

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
A      2 89.580 44.790 93.4102 3.013e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
A      2 126.778 63.389 132.1977 1.093e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
A      2 126.778 63.389 132.1977 1.093e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

4.3.2 p33

(16) MODEL

```
p33 = read.csv("C:/G/Rt/ANOVA/Goodnight-p33.csv")
p33 = af(p33, c("A", "B"))
GLM(y ~ A + B + A:B, p33) # p35
```

\$ANOVA
 Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	4	34.905	8.7261		
RESIDUALS	0	0.000			
CORRECTED TOTAL	4	34.905			

\$Fitness

R-square	Coef	Var	Root	MSE	y	Mean
1	NA	NA	NA	6.946		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	2	11.3739	5.6870		
B	1	23.5225	23.5225		
A:B	1	0.0081	0.0081		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	3.0276	3.0276		
B	1	23.5225	23.5225		
A:B	1	0.0081	0.0081		

\$`Type III`
 CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	3.0276	3.0276		
B	1	23.5225	23.5225		
A:B	1	0.0081	0.0081		

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(y ~ A + B + A:B, p33), type=3, singular.ok=TRUE) # NOT WORKING
```

5 SAS for Linear Models 4e

Reference

- Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.

5.1 Chapter 2

5.1.1 p5

(17) MODEL

```
p5 = read.table("C:/G/Rt/SAS4lm/p5.txt", head=TRUE)
GLM(COST ~ CATTLE, p5) # p6 Output 2.2

$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       1 6582.1  6582.1   59.34 6.083e-07 ***
RESIDUALS   17 1885.7    110.9
CORRECTED TOTAL 18 8467.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE COST Mean
0.7773107 29.84119 10.53198  35.29342

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE    1 6582.1  6582.1   59.34 6.083e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE    1 6582.1  6582.1   59.34 6.083e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE    1 6582.1  6582.1   59.34 6.083e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.1.2 p12

(18) MODEL

```
p12 = read.table("C:/G/Rt/SAS4lm/p12.txt", head=TRUE)
GLM(COST ~ CATTLE + CALVES + HOGS + SHEEP, p12)

$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       4 7936.7 1984.18   52.31 2.885e-08 ***
RESIDUALS   14  531.0   37.93
CORRECTED TOTAL 18 8467.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE COST Mean
0.9372871 17.4504 6.158842 35.29342

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 6582.1 6582.1 173.5265 2.801e-09 ***
CALVES     1 186.7  186.7  4.9213 0.0435698 *
HOGS       1 489.9  489.9 12.9145 0.0029351 **
SHEEP      1 678.1  678.1 17.8773 0.0008431 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2200.71 2200.71 58.0183 2.413e-06 ***
CALVES     1 136.08 136.08  3.5876 0.0790616 .
HOGS       1 113.66 113.66  2.9964 0.1054198
SHEEP      1 678.11 678.11 17.8773 0.0008431 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2200.71 2200.71 58.0183 2.413e-06 ***
CALVES     1 136.08 136.08  3.5876 0.0790616 .
HOGS       1 113.66 113.66  2.9964 0.1054198
SHEEP      1 678.11 678.11 17.8773 0.0008431 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(19) MODEL

```
GLM(COST ~ CATTLE + CALVES + SHEEP, p12)
```

```
$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       3 7823.1 2607.69  60.673 1.281e-08 ***
RESIDUALS   15  644.7   42.98
CORRECTED TOTAL 18 8467.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE COST Mean
0.9238649 18.57538 6.555887 35.29342

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 6582.1 6582.1 153.1443 2.835e-09 ***
CALVES     1  186.7   186.7   4.3432 0.0546701 .
SHEEP      1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES     1  260.6   260.6   6.0634 0.0263909 *
SHEEP      1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES     1  260.6   260.6   6.0634 0.0263909 *
SHEEP      1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(20) MODEL

```
GLM(COST ~ CATTLE + CALVES + offset(1*HOGS) + SHEEP, p12)
```

```
$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
```

```

MODEL           3 7823.1 2607.69 60.673 1.281e-08 ***
RESIDUALS      15 644.7   42.98
CORRECTED TOTAL 18 8467.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE COST Mean
0.9238649 18.57538 6.555887 35.29342

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE  1 6582.1 6582.1 153.1443 2.835e-09 ***
CALVES  1 186.7   186.7   4.3432 0.0546701 .
SHEEP   1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE  1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES  1 260.6   260.6   6.0634 0.0263909 *
SHEEP   1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE  1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES  1 260.6   260.6   6.0634 0.0263909 *
SHEEP   1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(21) MODEL

```
GLM(COST ~ CATTLE + CALVES + I(HOGS + SHEEP), p12)
```

```

$ANOVA
Response : COST
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       3 7936.7 2645.6 74.726 3.011e-09 ***
RESIDUALS   15 531.1   35.4
CORRECTED TOTAL 18 8467.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE COST Mean
0.937285 16.85896 5.950105 35.29342

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE        1 6582.1 6582.1 185.9151 7.406e-10 ***
CALVES        1 186.7 186.7  5.2726  0.03649 *
I(HOGS + SHEEP) 1 1168.0 1168.0 32.9896 3.883e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE        1 2215.48 2215.48 62.5775 9.887e-07 ***
CALVES        1 155.03 155.03  4.3788  0.0538 .
I(HOGS + SHEEP) 1 1167.96 1167.96 32.9896 3.883e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE        1 2215.48 2215.48 62.5775 9.887e-07 ***
CALVES        1 155.03 155.03  4.3788  0.0538 .
I(HOGS + SHEEP) 1 1167.96 1167.96 32.9896 3.883e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(22) MODEL

```
REG(COST ~ CATTLE + CALVES + I(HOGS + SHEEP) - 1, p12)
```

	Estimate	Std. Error	Df	t value	Pr(> t)
CATTLE	3.3000	0.38314	16	8.6131	2.100e-07 ***
CALVES	1.9672	0.59108	16	3.3281	0.004259 **
I(HOGS + SHEEP)	0.8068	0.13800	16	5.8466	2.479e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.2 Chapter 3

5.2.1 p63

(23) MODEL

```

p63w = read.table("C:/G/Rt/SAS4lm/p63.txt", header=TRUE)
p63l = reshape(p63w,
  direction = "long",
  varying = list(names(p63w)[2:9]),
  v.names = "fruitwt",
  idvar = c("irrig"),
  timevar = "bloc",
  times = 1:8)
p63l = af(p63l, c("bloc"))
GLM(fruitwt ~ bloc + irrig, p63l) # p64

```

```

$ANOVA
Response : fruitwt
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 445334   40485   12.04 6.643e-08 ***
RESIDUALS    28 94147    3362
CORRECTED TOTAL 39 539481
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE fruitwt Mean
0.8254864 21.71153 57.98607      267.075

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc     7 401308   57330 17.0503 1.452e-08 ***
irrig    4  44026   11006  3.2734  0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc     7 401308   57330 17.0503 1.452e-08 ***
irrig    4  44026   11006  3.2734  0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc     7 401308   57330 17.0503 1.452e-08 ***
irrig    4  44026   11006  3.2734  0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2.2 p72

(24) MODEL

```
p72 = read.table("C:/G/Rt/SAS4lm/p72.txt", header=TRUE)
p72 = af(p72, c("run", "pos", "mat"))
GLM(wtloss ~ run + pos + mat, p72) # p73
```

```
$ANOVA
Response : wtloss
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      9 7076.5  786.28  12.837 0.002828 ***
RESIDUALS   6  367.5   61.25
CORRECTED TOTAL 15 7444.0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE wtloss Mean
0.9506314 3.26774 7.826238      239.5

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3  986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3  986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3  986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

GLM(shrink ~ run + pos + mat, p72) # p73

$ANOVA
Response : shrink
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       9 265.75 29.528 9.8426 0.005775 ***
RESIDUALS     6 18.00  3.000
CORRECTED TOTAL 15 283.75
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE shrink Mean
0.9365639 3.675439 1.732051      47.125

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 33.25 11.083 3.6944 0.081254 .
pos   3 60.25 20.083 6.6944 0.024212 *
mat   3 172.25 57.417 19.1389 0.001786 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 33.25 11.083 3.6944 0.081254 .
pos   3 60.25 20.083 6.6944 0.024212 *
mat   3 172.25 57.417 19.1389 0.001786 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 33.25 11.083 3.6944 0.081254 .
pos   3 60.25 20.083 6.6944 0.024212 *
mat   3 172.25 57.417 19.1389 0.001786 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2.3 p75

(25) MODEL

```

p75w = read.table("C:/G/Rt/SAS4lm/p75.txt", header=TRUE)
p75l = reshape(p75w,
               direction = "long",

```

```

varying = list(names(p75w)[4:9]),
v.names = "Y",
idvar = c("method", "variety", "trt"),
timevar = "yield",
times = 1:6)
p751 = af(p751, c("variety", "yield"))
GLM(Y ~ method*variety, p751) # p78

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL     14 1339.0  95.645  4.8674 2.723e-06 ***
RESIDUALS   75 1473.8  19.650
CORRECTED TOTAL 89 2812.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE    Y Mean
0.4760484 24.04225 4.432857 18.43778

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
method     2 953.16  476.58 24.2531 7.525e-09 ***
variety    4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81  2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
method     2 953.16  476.58 24.2531 7.525e-09 ***
variety    4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81  2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
method     2 953.16  476.58 24.2531 7.525e-09 ***
variety    4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81  2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3 Chapter 4

5.3.1 p94

(26) MODEL

```
p94w = read.table("C:/G/Rt/SAS4lm/p94.txt", head=TRUE)
p94l = reshape(p94w,
  direction = "long",
  varying = list(names(p94w)[3:8]),
  v.names = "ct",
  idvar = c("package"),
  timevar = "sample",
  times = 1:6)
p94l$sampleA = floor((p94l$sample + 1)/2)
p94l$sampleB = 2 - (p94l$sample) %% 2
p94l$logct = log10(p94l$ct)
p94l = af(p94l, c("sample", "sampleA", "sampleB", "package"))
GLM(logct ~ package + sampleA %in% package, p94l) # p97
```

```
$ANOVA
Response : logct
            Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          59 50.463 0.85531  22.229 < 2.2e-16 ***
RESIDUALS      60  2.309 0.03848
CORRECTED TOTAL 119 52.772
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE logct Mean
0.9562528 6.432487 0.196156   3.049459

$`Type I`
            Df Sum Sq Mean Sq F value    Pr(>F)
package        19 30.529 1.60680  41.760 < 2.2e-16 ***
package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
            Df Sum Sq Mean Sq F value    Pr(>F)
package        19 30.529 1.60680  41.760 < 2.2e-16 ***
package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

package          19 30.529 1.60680  41.760 < 2.2e-16 ***  

package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.3.2 p116

(27) MODEL

```
GLM(Y ~ method + variety + method:variety, p751) # p116
```

```
$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL       14 1339.0  95.645  4.8674 2.723e-06 ***  

RESIDUALS   75 1473.8  19.650  

CORRECTED TOTAL 89 2812.8  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE    Y Mean  

  0.4760484 24.04225 4.432857 18.43778
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

method        2 953.16  476.58 24.2531 7.525e-09 ***  

variety       4 11.38    2.85  0.1448  0.96476  

method:variety 8 374.49   46.81  2.3822  0.02409 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

method        2 953.16  476.58 24.2531 7.525e-09 ***  

variety       4 11.38    2.85  0.1448  0.96476  

method:variety 8 374.49   46.81  2.3822  0.02409 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

method        2 953.16  476.58 24.2531 7.525e-09 ***  

variety       4 11.38    2.85  0.1448  0.96476
```

```

method:variety 8 374.49 46.81 2.3822 0.02409 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3.3 p122

(28) MODEL

```

p122 = read.table("C:/G/Rt/SAS4lm/p122.txt", header=TRUE)
p122 = af(p122, c("et", "wafer", "pos"))
GLM(resista ~ et + wafer %in% et + pos + et:pos, p122)

```

```

$ANOVA
Response : resista
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      23 9.3250 0.40544 3.6477 0.001263 **
RESIDUALS   24 2.6676 0.11115
CORRECTED TOTAL 47 11.9926
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE resista Mean
0.7775641 5.553811 0.3333906     6.002917

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
et      3 3.1122 1.03739 9.3333 0.0002851 ***
et:wafer 8 4.2745 0.53431 4.8071 0.0012742 **
pos      3 1.1289 0.37630 3.3855 0.0345139 *
et:pos    9 0.8095 0.08994 0.8092 0.6125279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
et      3 3.1122 1.03739 9.3333 0.0002851 ***
et:wafer 8 4.2745 0.53431 4.8071 0.0012742 **
pos      3 1.1289 0.37630 3.3855 0.0345139 *
et:pos    9 0.8095 0.08994 0.8092 0.6125279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
et      3 3.1122 1.03739 9.3333 0.0002851 ***

```

```

et:wafer  8 4.2745 0.53431  4.8071 0.0012742 ***
pos       3 1.1289 0.37630  3.3855 0.0345139 *
et:pos    9 0.8095 0.08994  0.8092 0.6125279
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3.4 p136

(29) MODEL

```

p136 = read.table("C:/G/Rt/SAS4lm/p136.txt", header=TRUE)
p136 = af(p136, "rep")
GLM(drywt ~ rep + cult + rep:cult + inoc + cult:inoc, p136)

```

\$ANOVA

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	11	157.208	14.2917	20.26	4.594e-06 ***
RESIDUALS	12	8.465	0.7054		
CORRECTED TOTAL	23	165.673			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	drywt	Mean
0.9489055	2.761285	0.8398909		30.41667		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	3	25.320	8.440	11.9646	0.0006428 ***
cult	1	2.407	2.407	3.4117	0.0895283 .
rep:cult	3	9.480	3.160	4.4796	0.0249095 *
inoc	2	118.176	59.088	83.7631	8.919e-08 ***
cult:inoc	2	1.826	0.913	1.2942	0.3097837

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	3	25.320	8.440	11.9646	0.0006428 ***
cult	1	2.407	2.407	3.4117	0.0895283 .
rep:cult	3	9.480	3.160	4.4796	0.0249095 *
inoc	2	118.176	59.088	83.7631	8.919e-08 ***
cult:inoc	2	1.826	0.913	1.2942	0.3097837

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep        3  25.320   8.440 11.9646 0.0006428 ***  

cult       1   2.407   2.407  3.4117 0.0895283 .  

rep:cult   3   9.480   3.160  4.4796 0.0249095 *  

inoc       2 118.176  59.088 83.7631 8.919e-08 ***  

cult:inoc  2   1.826   0.913  1.2942 0.3097837  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.4 Chapter 5

5.4.1 p142

(30) MODEL

```
p142 = read.table("C:/G/Rt/SAS4lm/p142.txt", header=TRUE, na.strings=".")  

p142 = af(p142, c("STUDY", "PATIENT"))  

GLM(FLUSH ~ STUDY + TRT, p142) # Incomplete data, 56 lines are truncated.
```

```
$ANOVA  

Response : FLUSH  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      5  3619.9  723.98   2.392 0.04607 *  

RESIDUALS   71 21489.2   302.67  

CORRECTED TOTAL 76 25109.1  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE FLUSH Mean  

  0.1441665 75.2251 17.39728 23.12697
```

```
$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

STUDY     4 3553.9  888.46   2.9355 0.02638 *  

TRT       1   66.0   66.04   0.2182 0.64185  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

STUDY     4 3599.4  899.85   2.9731 0.02496 *  

TRT       1   66.0   66.04   0.2182 0.64185  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

STUDY   4 3599.4 899.85 2.9731 0.02496 *  

TRT     1   66.0   66.04  0.2182 0.64185  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(31) MODEL

```
GLM(FLUSH ~ TRT + STUDY + TRT:STUDY, p142) # Different data
```

```

$ANOVA  

Response : FLUSH  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      9 4093.7 454.86 1.4501 0.1851  

RESIDUALS   67 21015.4 313.66  

CORRECTED TOTAL 76 25109.1

```

```

$Fitness  

R-square Coef Var Root MSE FLUSH Mean  

0.1630364 76.57962 17.71054 23.12697

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

TRT       1   20.5   20.49  0.0653 0.79906  

STUDY     4 3599.4 899.85  2.8688 0.02956 *  

TRT:STUDY 4  473.8 118.45  0.3776 0.82383  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

TRT       1   66.0   66.04  0.2105 0.64783  

STUDY     4 3599.4 899.85  2.8688 0.02956 *  

TRT:STUDY 4  473.8 118.45  0.3776 0.82383  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

TRT       1    1.9    1.93  0.0062 0.9377  

STUDY     4 3339.4 834.85  2.6616 0.0400 *  

TRT:STUDY 4  473.8 118.45  0.3776 0.8238  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.5 Chapter 6

5.5.1 p171

(32) MODEL

```
p171 = read.table("C:/G/Rt/SAS4lm/p171.txt", header=TRUE)
GLM(score2 ~ teach, p171) # p173 Output 6.2, p174 Output 6.5
```

```
$ANOVA
Response : score2
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      2   49.74  24.868  0.5598 0.5776
RESIDUALS  28 1243.94  44.426
CORRECTED TOTAL 30 1293.68
```

```
$Fitness
  R-square Coef Var Root MSE score2 Mean
0.03844533 9.062496 6.66532    73.54839
```

```
$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
teach  2 49.736  24.868  0.5598 0.5776
```

```
$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
teach  2 49.736  24.868  0.5598 0.5776
```

```
$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
teach  2 49.736  24.868  0.5598 0.5776
```

5.5.2 p188

(33) MODEL

```
p188 = read.table("C:/G/Rt/SAS4lm/p188.txt", header=TRUE)
p188 = af(p188, c("a", "b"))
GLM(y ~ a + b + a:b, p188) # p189
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 63.711 12.7422   5.866 0.005724 ***
RESIDUALS 12 26.067  2.1722
CORRECTED TOTAL 17 89.778
```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   y Mean
0.7096535 28.83612 1.473846 5.111111

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
a     1 7.803  7.8028  3.5921 0.082395 .
b     2 20.492 10.2459  4.7168 0.030798 *
a:b   2 35.416 17.7082  8.1521 0.005807 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
a     1 15.850 15.850  7.2968 0.019265 *
b     2 20.492 10.246  4.7168 0.030798 *
a:b   2 35.416 17.708  8.1521 0.005807 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
a     1 9.641  9.6407  4.4382 0.056865 .
b     2 30.866 15.4330  7.1047 0.009212 **
a:b   2 35.416 17.7082  8.1521 0.005807 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.5.3 p203

(34) MODEL

```
GLM(y ~ a + b + a:b,  p188[-8,])
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          4 45.816 11.4539  5.2729 0.01097 *
RESIDUALS      12 26.067  2.1722
CORRECTED TOTAL 16 71.882

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE   y Mean
  0.6373704 27.53339 1.473846 5.352941

$`Type I` 
  Df  Sum Sq Mean Sq F value    Pr(>F)
a    1  2.9252  2.9252  1.3466 0.268432
b    2 13.3224  6.6612  3.0665 0.083997 .
a:b  1 29.5681 29.5681 13.6119 0.003095 ** 
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df  Sum Sq Mean Sq F value    Pr(>F)
a    1  5.5652  5.5652  2.5620 0.135442
b    2 13.3224  6.6612  3.0665 0.083997 .
a:b  1 29.5681 29.5681 13.6119 0.003095 ** 
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df  Sum Sq Mean Sq F value    Pr(>F)
a    1  0.3507  0.3507  0.1615 0.694881
b    2 16.0733  8.0367  3.6997 0.056021 .
a:b  1 29.5681 29.5681 13.6119 0.003095 ** 
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.5.4 p215

(35) MODEL

```

p215 = read.table("C:/G/Rt/SAS4lm/p215.txt", header=TRUE)
p215 = af(p215, c("irrig", "reps"))
GLM(yield ~ irrig/reps + cult + irrig:cult, p215) # p216 Book is wrong.

```

```

$ANOVA
Response : yield
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        11  67.662  6.1511  0.6253 0.7636
RESIDUALS      6  59.023  9.8372
CORRECTED TOTAL 17 126.685

```

```

$Fitness
  R-square Coef Var Root MSE yield Mean
  0.5340937 10.1448 3.136435 30.91667

```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

irrig       2  7.320  3.6600  0.3721 0.7042  

irrig:reps  6 59.870  9.9783  1.0143 0.4933  

cult        1  0.467  0.4672  0.0475 0.8347  

irrig:cult  2  0.004  0.0022  0.0002 0.9998  

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

irrig       2  7.320  3.6600  0.3721 0.7042  

irrig:reps  6 59.870  9.9783  1.0143 0.4933  

cult        1  0.467  0.4672  0.0475 0.8347  

irrig:cult  2  0.004  0.0022  0.0002 0.9998  

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

irrig       2  7.320  3.6600  0.3721 0.7042  

irrig:reps  6 59.870  9.9783  1.0143 0.4933  

cult        1  0.467  0.4672  0.0475 0.8347  

irrig:cult  2  0.004  0.0022  0.0002 0.9998
```

```
# Compare with SAS output
```

(36) MODEL

```
GLM(yield ~ reps + irrig + reps:irrig + cult + cult:irrig, p215)
```

```
$ANOVA  

Response : yield  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL          11 67.662  6.1511  0.6253 0.7636  

RESIDUALS       6 59.023  9.8372  

CORRECTED TOTAL 17 126.685  

$Fitness  

  R-square Coef Var Root MSE yield Mean  

  0.5340937 10.1448 3.136435   30.91667  

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998  

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998
```

5.6 Chapter 7

5.6.1 p232

(37) MODEL

```
p232 = read.table("C:/G/Rt/SAS4lm/p232.txt", header=TRUE)
p232 = af(p232, c("trt", "rep"))
GLM(final ~ trt + initial, p232) # p233
```

```
$ANOVA
Response : final
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 354.45  70.889  235.05 5.493e-13 ***
RESIDUALS   14   4.22   0.302
CORRECTED TOTAL 19 358.67
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE final Mean
0.9882278 1.780438 0.5491762      30.845

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)
trt       4 198.41  49.602  164.47 1.340e-11 ***
initial   1 156.04 156.040  517.38 1.867e-12 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

trt      4  12.089   3.022  10.021 0.0004819 ***  

initial  1 156.040 156.040 517.384 1.867e-12 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

trt      4  12.089   3.022  10.021 0.0004819 ***  

initial  1 156.040 156.040 517.384 1.867e-12 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.2 p240

(38) MODEL

```

GLM(final ~ initial + trt + trt:initial, p232) # p240

```

\$ANOVA

Response : final

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	9	355.84	39.537	139.51	2.572e-09 ***
RESIDUALS	10	2.83	0.283		
CORRECTED TOTAL	19	358.67			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	final	Mean
0.9920985	1.725901	0.5323541		30.845		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
initial	1	342.36	342.36	1208.0336	9.211e-12 ***
trt	4	12.09	3.02	10.6645	0.001247 **
initial:trt	4	1.39	0.35	1.2247	0.360175

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
initial	1	156.040	156.040	550.5987	4.478e-10 ***
trt	4	12.089	3.022	10.6645	0.001247 **

```

initial:trt 4 1.388 0.347 1.2247 0.360175
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
initial     1 68.529 68.529 241.8091 2.472e-08 ***
trt        4  1.696   0.424   1.4963   0.2752
initial:trt 4  1.388   0.347   1.2247   0.3602
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.3 p241

(39) MODEL

```

p241 = read.table("C:/G/Rt/SAS4lm/p241.txt", header=TRUE)
p241 = af(p241, c("STORE", "DAY"))
GLM(Q1 ~ P1 + DAY + P1:DAY, p241) # p242

```

```

$ANOVA
Response : Q1
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 1111.52 101.048 4.6445 0.0008119 ***
RESIDUALS    24 522.15  21.756
CORRECTED TOTAL 35 1633.68
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  Q1 Mean
  0.6803814 45.65257 4.664374 10.21711

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
P1       1 516.59 516.59 23.7444 5.739e-05 ***
DAY      5 430.54  86.11  3.9578  0.009275 **
P1:DAY   5 164.39  32.88  1.5112  0.223566
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
P1       1 696.73 696.73 32.0243 7.925e-06 ***
DAY      5 430.54  86.11  3.9578  0.009275 **
P1:DAY   5 164.39  32.88  1.5112  0.223566

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
P1      1 554.79  554.79 25.4999 3.665e-05 ***
DAY     5 201.17   40.23  1.8493   0.1412
P1:DAY  5 164.39   32.88  1.5112   0.2236
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.4 p243

(40) MODEL

```
GLM(Q1 ~ DAY + DAY:P1, p241)
```

```

$ANOVA
Response : Q1
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 1111.52 101.048 4.6445 0.0008119 ***
RESIDUALS    24 522.15  21.756
CORRECTED TOTAL 35 1633.68
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE  Q1 Mean
0.6803814 45.65257 4.664374 10.21711

```

```

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
DAY      5 250.40  50.079  2.3018 0.0764717 .
DAY:P1   6 861.13 143.521  6.5967 0.0003239 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
DAY      5 250.40  50.079  2.3018 0.0764717 .
DAY:P1   6 861.13 143.521  6.5967 0.0003239 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)

```

```

DAY      5 201.17 40.234 1.8493 0.1411648
DAY:P1   6 861.13 143.521 6.5967 0.0003239 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
REG(Q1 ~ DAY + DAY:P1 - 1, p241) # Output 7.10
```

	Estimate	Std. Error	Df	t value	Pr(> t)
DAY1	18.675	14.4110	24	1.2959	0.2073286
DAY2	38.487	15.1094	24	2.5472	0.0176863 *
DAY3	45.330	26.1576	24	1.7329	0.0959384 .
DAY4	49.149	16.6092	24	2.9592	0.0068366 **
DAY5	77.899	27.5007	24	2.8326	0.0092034 **
DAY6	73.273	13.4837	24	5.4341	1.39e-05 ***
DAY1:P1	-0.220	0.2915	24	-0.7562	0.4568599
DAY2:P1	-0.624	0.2978	24	-2.0940	0.0470031 *
DAY3:P1	-0.611	0.5049	24	-1.2102	0.2379998
DAY4:P1	-0.796	0.3193	24	-2.4914	0.0200350 *
DAY5:P1	-1.196	0.5049	24	-2.3683	0.0262648 *
DAY6:P1	-1.225	0.2652	24	-4.6199	0.0001092 ***

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(41) MODEL

```
GLM(Q1 ~ P1 + DAY + P1:DAY, p241)
```

\$ANOVA

Response : Q1

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	11	1111.52	101.048	4.6445	0.0008119 ***
RESIDUALS	24	522.15	21.756		
CORRECTED TOTAL	35	1633.68			

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

R-square	Coef	Var	Root	MSE	Q1	Mean
0.6803814	45.65257	4.664374	10.21711			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P1	1	516.59	516.59	23.7444	5.739e-05 ***
DAY	5	430.54	86.11	3.9578	0.009275 **
P1:DAY	5	164.39	32.88	1.5112	0.223566

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

P1       1 696.73 696.73 32.0243 7.925e-06 ***  

DAY      5 430.54   86.11  3.9578  0.009275 **  

P1:DAY  5 164.39   32.88  1.5112  0.223566  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

P1       1 554.79 554.79 25.4999 3.665e-05 ***  

DAY      5 201.17   40.23  1.8493   0.1412  

P1:DAY  5 164.39   32.88  1.5112   0.2236  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(42) MODEL

```
GLM(Q1 ~ STORE + DAY + P1 + P2, p241)
```

```

$ANOVA  

Response : Q1  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      12 1225.37 102.114  5.7521 0.0001688 ***  

RESIDUALS  23  408.31  17.753  

CORRECTED TOTAL 35 1633.68  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

R-square Coef Var Root MSE  Q1 Mean  

0.7500678 41.23842 4.213375 10.21711

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

STORE     5 313.42   62.68  3.5310   0.01629 *  

DAY      5 250.40   50.08  2.8210   0.03957 *  

P1       1 622.01  622.01 35.0377 4.924e-06 ***  

P2       1  39.54   39.54  2.2274   0.14917  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

STORE     5 223.83   44.77  2.5217  0.058346 .

```

```

DAY      5 433.10   86.62  4.8793  0.003456 **
P1       1 538.17   538.17 30.3150 1.342e-05 ***
P2       1 39.54    39.54   2.2274  0.149171
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df Sum Sq Mean Sq F value    Pr(>F)
STORE   5 223.83   44.77  2.5217  0.058346 .
DAY     5 433.10   86.62  4.8793  0.003456 **
P1      1 538.17   538.17 30.3150 1.342e-05 ***
P2      1 39.54    39.54   2.2274  0.149171
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.5 p250

(43) MODEL

```

p250 = read.table("C:/G/Rt/SAS4lm/p250.txt", header=TRUE)
p250 = af(p250, c("variety", "spacing", "plant"))
GLM(lint ~ bollwt + variety + spacing + variety:spacing + variety:spacing:plant,
    p250) # p252 Output 7.18, Parameter is different due to different order

```

```

$ANOVA
Response : lint
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      8 31.160  3.8950  80.704 < 2.2e-16 ***
RESIDUALS   40 1.931   0.0483
CORRECTED TOTAL 48 33.091
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE lint Mean
  0.9416596 12.37325 0.2196884   1.77551

$`Type I` 
  Df Sum Sq Mean Sq F value    Pr(>F)
bollwt      1 29.0693 29.0693 602.3107 < 2.2e-16 ***
variety      1  1.2635  1.2635  26.1802 8.158e-06 ***
spacing      1  0.4666  0.4666   9.6689  0.003447 **
variety:spacing  1  0.0933  0.0933   1.9325  0.172169
variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`  

          Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt      1 11.1186 11.1186 230.3745 < 2.2e-16 ***  

variety     1  1.1973  1.1973  24.8084 1.259e-05 ***  

spacing     1  0.4666  0.4666   9.6689  0.003447 **  

variety:spacing  1  0.0933  0.0933   1.9325  0.172169  

variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

          Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt      1 11.1186 11.1186 230.3745 < 2.2e-16 ***  

variety     1  0.9424  0.9424  19.5269 7.379e-05 ***  

spacing     1  0.3748  0.3748   7.7666  0.008101 **  

variety:spacing  1  0.0479  0.0479   0.9915  0.325350  

variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.6.6 p254 Output 7.20

(44) MODEL

```
GLM(lint ~ bollwt + variety + spacing, p250)
```

```
$ANOVA  

Response : lint  

          Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      3 30.799 10.2665 201.65 < 2.2e-16 ***  

RESIDUALS  45  2.291  0.0509  

CORRECTED TOTAL 48 33.091  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE lint Mean  

0.9307624 12.70849 0.2256406 1.77551
```

```
$`Type I`  

          Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt    1 29.0693 29.0693 570.9531 < 2.2e-16 ***  

variety   1  1.2635  1.2635  24.8172 9.777e-06 ***  

spacing   1  0.4666  0.4666   9.1655  0.004072 **  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt   1 11.5717 11.5717 227.2815 < 2.2e-16 ***  

variety  1  1.1973  1.1973  23.5168 1.516e-05 ***  

spacing   1  0.4666  0.4666   9.1655  0.004072 **  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt   1 11.5717 11.5717 227.2815 < 2.2e-16 ***  

variety  1  1.1973  1.1973  23.5168 1.516e-05 ***  

spacing   1  0.4666  0.4666   9.1655  0.004072 **  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.7 p256

(45) MODEL

```

p256 = read.table("C:/G/Rt/SAS4lm/p256.txt", header=TRUE)
p256b = af(p256, c("bloc", "type", "logdose"))
GLM(y ~ bloc + type + logdose + type:logdose, p256b) # p258 Output 7.22

```

```

$ANOVA  

Response : y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      8  816.50 102.063  6.0641 0.0014 **  

RESIDUALS  15  252.46  16.831  

CORRECTED TOTAL 23 1068.96  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE   y Mean  

  0.7638277 7.464757 4.102506 54.95833

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04  12.042  0.7155 0.4109264  

logdose    2 121.58  60.792  3.6120 0.0524231 .  

type:logdose 2 144.08  72.042  4.2804 0.0338265 *  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc      3 538.79 179.597 10.6709 0.0005223 ***  

type      1 12.04 12.042  0.7155 0.4109264  

logdose    2 121.58 60.792  3.6120 0.0524231 .  

type:logdose 2 144.08 72.042  4.2804 0.0338265 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc      3 538.79 179.597 10.6709 0.0005223 ***  

type      1 12.04 12.042  0.7155 0.4109264  

logdose    2 121.58 60.792  3.6120 0.0524231 .  

type:logdose 2 144.08 72.042  4.2804 0.0338265 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.8 p261 Output 7.27

(46) MODEL

```

p256 = af(p256, c("bloc", "type"))
p256$logd2 = (p256$logdose)^2
GLM(y ~ bloc + type + logdose + logd2 + type:logdose + type:logd2, p256)

```

```

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL     8   816.50 102.063  6.0641 0.0014 **
RESIDUALS 15   252.46 16.831
CORRECTED TOTAL 23 1068.96
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE   y Mean
0.7638277 7.464757 4.102506 54.95833

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc      3 538.79 179.597 10.6709 0.0005223 ***  

type      1 12.04 12.042  0.7155 0.4109264  

logdose   1 115.56 115.562  6.8662 0.0193005 *

```

```

logd2           1   6.02   6.021  0.3577 0.5586917
type:logdose   1 138.06 138.062  8.2031 0.0118242 *
type:logd2     1   6.02   6.021  0.3577 0.5586917
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04 12.042  0.7155 0.4109264  

logdose    1   0.39   0.389  0.0231 0.8811262  

logd2      1   6.02   6.021  0.3577 0.5586917  

type:logdose 1   0.81   0.812  0.0483 0.8290541  

type:logd2   1   6.02   6.021  0.3577 0.5586917
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 28.12 28.125  1.6711 0.2156736  

logdose    1   0.39   0.389  0.0231 0.8811262  

logd2      1   6.02   6.021  0.3577 0.5586917  

type:logdose 1   0.81   0.812  0.0483 0.8290541  

type:logd2   1   6.02   6.021  0.3577 0.5586917
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.9 p262 Output 7.28

(47) MODEL

```
GLM(y ~ bloc + type + type:logdose, p256b)
```

```
$ANOVA  

Response : y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      8 816.50 102.063  6.0641 0.0014 **  

RESIDUALS  15 252.46 16.831  

CORRECTED TOTAL 23 1068.96
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE   y Mean  

0.7638277 7.464757 4.102506 54.95833
```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04 12.042  0.7155 0.4109264  

type:logdose 4 265.67 66.417  3.9462 0.0220552 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04 12.042  0.7155 0.4109264  

type:logdose 4 265.67 66.417  3.9462 0.0220552 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04 12.042  0.7155 0.4109264  

type:logdose 4 265.67 66.417  3.9462 0.0220552 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.7 Chapter 8

5.7.1 p269

(48) MODEL

```

p269 = read.csv("C:/G/Rt/SAS4lm/fev1uni.csv")
p269 = af(p269, c("drug", "hour", "patient"))
GLM(fev1 ~ drug + patient %in% drug + hour + drug:hour, p269) # p271 Output 8.3

```

```

$ANOVA
Response : fev1
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      92 296.65  3.2244  51.078 < 2.2e-16 ***
RESIDUALS   483 30.49  0.0631
CORRECTED TOTAL 575 327.14
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE fev1 Mean

```

```

0.9067963 8.138859 0.2512505 3.087049

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

drug        2 25.783 12.8913 204.212 < 2.2e-16 ***  

drug:patient 69 247.412  3.5857  56.801 < 2.2e-16 ***  

hour        7 17.170  2.4529  38.857 < 2.2e-16 ***  

drug:hour     14  6.280  0.4486   7.106 1.923e-13 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

drug        2 25.783 12.8913 204.212 < 2.2e-16 ***  

drug:patient 69 247.412  3.5857  56.801 < 2.2e-16 ***  

hour        7 17.170  2.4529  38.857 < 2.2e-16 ***  

drug:hour     14  6.280  0.4486   7.106 1.923e-13 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

drug        2 25.783 12.8913 204.212 < 2.2e-16 ***  

drug:patient 69 247.412  3.5857  56.801 < 2.2e-16 ***  

hour        7 17.170  2.4529  38.857 < 2.2e-16 ***  

drug:hour     14  6.280  0.4486   7.106 1.923e-13 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8 Chapter 11

5.8.1 p390

(49) MODEL

```

p390 = read.table("C:/G/Rt/SAS4lm/p390.txt", header=TRUE)
p390$ca = ifelse(p390$a == 0, -1, 1)
p390$cb = ifelse(p390$b == 0, -1, 1)
p390$cc = ifelse(p390$c == 0, -1, 1)
p390 = af(p390, c("rep", "blk", "a", "b", "c"))
GLM(y ~ rep blk + ca*cb*cc, p390)

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL       12 81.75  6.8125 33.601 6.618e-07 ***

```

```

RESIDUALS           11   2.23  0.2027
CORRECTED TOTAL  23  83.98
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var  Root MSE  y Mean
0.9734438 18.96878 0.4502714 2.37375

$`Type I`
      Df Sum Sq Mean Sq  F value    Pr(>F)
rep       2  0.051   0.025   0.1256  0.8832237
rep:blk   3  7.432   2.477  12.2194 0.0007966 ***
ca        1 21.075  21.075 103.9487 6.090e-07 ***
cb        1  0.005   0.005   0.0224  0.8837872
ca:cb    1  1.723   1.723   8.4969  0.0140640 *
cc        1 37.776  37.776 186.3209 3.063e-08 ***
ca:cc    1  2.318   2.318  11.4332  0.0061285 **
cb:cc    1 11.340  11.340  55.9328 1.232e-05 ***
ca:cb:cc 1  0.031   0.031   0.1511  0.7049490
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq  F value    Pr(>F)
rep       2  0.051   0.025   0.1256  0.883224
rep:blk   3  1.668   0.556   2.7416  0.093789 .
ca        1 21.075  21.075 103.9487 6.090e-07 ***
cb        1  0.005   0.005   0.0224  0.883787
ca:cb    1  1.723   1.723   8.4969  0.014064 *
cc        1 37.776  37.776 186.3209 3.063e-08 ***
ca:cc    1  2.318   2.318  11.4332  0.006129 **
cb:cc    1 11.340  11.340  55.9328 1.232e-05 ***
ca:cb:cc 1  0.031   0.031   0.1511  0.704949
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq  F value    Pr(>F)
rep       2  0.051   0.025   0.1256  0.883224
rep:blk   3  1.668   0.556   2.7416  0.093789 .
ca        1 21.075  21.075 103.9487 6.090e-07 ***
cb        1  0.005   0.005   0.0224  0.883787
ca:cb    1  1.723   1.723   8.4969  0.014064 *
cc        1 37.776  37.776 186.3209 3.063e-08 ***
ca:cc    1  2.318   2.318  11.4332  0.006129 **
cb:cc    1 11.340  11.340  55.9328 1.232e-05 ***
ca:cb:cc 1  0.031   0.031   0.1511  0.704949

```

```
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.8.2 p394

(50) MODEL

```
p394 = read.table("C:/G/Rt/SAS4lm/p394.txt", header=TRUE)
p394 = af(p394, c("a", "b", "c", "d"))
GLM(y ~ ca*cb*cc*cd, p394)
```

```
$ANOVA
Response : y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7 6.3559 0.90798
RESIDUALS      0 0.0000
CORRECTED TOTAL 7 6.3559
```

```
$Fitness
R-square Coef Var Root MSE  y Mean
      1       NA     NA 2.68875
```

```
$`Type I`
              Df Sum Sq Mean Sq F value Pr(>F)
ca            1 2.07061 2.07061
cb            1 0.59951 0.59951
ca:cb         1 0.00031 0.00031
cc            1 0.00551 0.00551
ca:cc         1 0.80011 0.80011
cb:cc         1 2.82031 2.82031
ca:cb:cc     1 0.05951 0.05951
cd            0
ca:cd         0
cb:cd         0
ca:cb:cd     0
cc:cd         0
ca:cc:cd     0
cb:cc:cd     0
ca:cb:cc:cd  0
```

```
$`Type II`
              Df Sum Sq Mean Sq F value Pr(>F)
ca            0
cb            0
ca:cb         0
cc            0
```

```

ca:cc      0
cb:cc      0
ca:cb:cc   0
cd         0
ca:cd      0
cb:cd      0
ca:cb:cd   0
cc:cd      0
ca:cc:cd   0
cb:cc:cd   0
ca:cb:cc:cd 0

$`Type III`  

CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)
ca        0
cb        0
ca:cb     0
cc        0
ca:cc     0
cb:cc     0
ca:cb:cc  0
cd        0
ca:cd     0
cb:cd     0
ca:cb:cd  0
cc:cd     0
ca:cc:cd  0
cb:cc:cd  0
ca:cb:cc:cd 0

```

(51) MODEL

```
GLM(y ~ a*b*c*d, p394)
```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       7 6.3559 0.90798
RESIDUALS    0 0.0000
CORRECTED TOTAL 7 6.3559

```

```

$Fitness
R-square Coef Var Root MSE  y Mean
      1       NA      NA 2.68875

```

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
a	1	2.07061	2.07061		
b	1	0.59951	0.59951		
a:b	1	0.00031	0.00031		
c	1	0.00551	0.00551		
a:c	1	0.80011	0.80011		
b:c	1	2.82031	2.82031		
a:b:c	1	0.05951	0.05951		
d	0				
a:d	0				
b:d	0				
a:b:d	0				
c:d	0				
a:c:d	0				
b:c:d	0				
a:b:c:d	0				

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
a	0				
b	0				
a:b	0				
c	0				
a:c	0				
b:c	0				
a:b:c	0				
d	0				
a:d	0				
b:d	0				
a:b:d	0				
c:d	0				
a:c:d	0				
b:c:d	0				
a:b:c:d	0				

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
a	0				
b	0				
a:b	0				
c	0				
a:c	0				
b:c	0				
a:b:c	0				
d	0				
a:d	0				
b:d	0				

```
a:b:d      0
c:d      0
a:c:d      0
b:c:d      0
a:b:c:d    0
```

5.8.3 p399

(52) MODEL

```
p399 = read.table("C:/G/Rt/SAS4lm/p399.txt", header=TRUE)
p399 = af(p399, c("blk", "trt"))
GLM(y ~ trt + blk, p399)

$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       8 281.127 35.141 40.822 0.005606 ***
RESIDUALS   3   2.583   0.861
CORRECTED TOTAL 11 283.710
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE y Mean
0.9908974 9.516011 0.927811   9.75

$`Type I`
          Df  Sum Sq Mean Sq F value    Pr(>F)
trt       3 102.26 34.086 39.596 0.006515 ***
blk       5 178.87 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df  Sum Sq Mean Sq F value    Pr(>F)
trt       3 59.018 19.673 22.853 0.014388 *
blk       5 178.871 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df  Sum Sq Mean Sq F value    Pr(>F)
trt       3 59.017 19.672 22.853 0.014388 *
blk       5 178.871 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.8.4 p403

(53) MODEL

```
p403 = read.table("C:/G/Rt/SAS4lm/p403.txt", header=TRUE)
p403 = af(p403, c("PATIENT", "VISIT"))
GLM(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT, p403)
```

\$ANOVA

Response : HR

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	29	6408.7	220.99	3.912	3.127e-05 ***
RESIDUALS	42	2372.6	56.49		
CORRECTED TOTAL	71	8781.3			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	HR	Mean
0.7298134	9.301326	7.515988	80.80556			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SEQUENCE	5	508.9	101.79	1.8019	0.133346
SEQUENCE:PATIENT	18	4692.3	260.69	4.6147	2.21e-05 ***
VISIT	2	146.8	73.39	1.2991	0.283499
DRUG	2	668.8	334.39	5.9194	0.005435 **
RESIDS	1	391.0	391.02	6.9219	0.011854 *
RESIDT	1	0.8	0.84	0.0149	0.903511

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SEQUENCE	5	701.2	140.237	2.4825	0.04665 *
SEQUENCE:PATIENT	18	4692.3	260.685	4.6147	2.21e-05 ***
VISIT	2	146.8	73.389	1.2991	0.28350
DRUG	2	344.0	171.975	3.0443	0.05826 .
RESIDS	1	309.2	309.174	5.4731	0.02414 *
RESIDT	1	0.8	0.840	0.0149	0.90351

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SEQUENCE	5	701.2	140.237	2.4825	0.04665 *
SEQUENCE:PATIENT	18	4692.3	260.685	4.6147	2.21e-05 ***

```

VISIT          2   146.8  73.389  1.2991  0.28350
DRUG          2   343.9  171.975  3.0443  0.05826 .
RESIDS        1   309.2  309.174  5.4731  0.02414 *
RESIDT        1     0.8    0.840  0.0149  0.90351
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT,
         p403), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: HR
      Sum Sq Df F values    Pr(>F)
SEQUENCE       0.0  0
VISIT          146.8  2   1.2991  0.28350
DRUG           344.0  2   3.0443  0.05826 .
RESIDS         309.2  1   5.4731  0.02414 *
RESIDT         0.8   1   0.0149  0.90351
SEQUENCE:PATIENT 4692.3 18   4.6147  2.21e-05 ***
Residuals      2372.6 42
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.5 p409 11.5

(54) MODEL

```

p409 = read.table("C:/G/Rt/SAS4lm/p409.txt", header=TRUE)
GLM(TS ~ SOURCE*AMT, p409) # p410 Output 11.21

```

```

$ANOVA
Response : TS
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      5  258.727  51.745  263.71 1.785e-09 ***
RESIDUALS  9    1.766   0.196
CORRECTED TOTAL 14  260.493
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$Fitness

```

R-square Coef Var Root MSE TS Mean
0.9932206 2.762805 0.4429698 16.03333

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
SOURCE      2 98.001 49.001 249.720 1.306e-08 ***
AMT         1 138.245 138.245 704.534 7.392e-10 ***
SOURCE:AMT  2 22.481 11.240  57.284 7.595e-06 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
SOURCE      2 98.001 49.001 249.720 1.306e-08 ***
AMT         1 138.245 138.245 704.534 7.392e-10 ***
SOURCE:AMT  2 22.481 11.240  57.284 7.595e-06 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
SOURCE      2 0.070  0.035  0.179     0.839
AMT         1 138.245 138.245 704.534 7.392e-10 ***
SOURCE:AMT  2 22.481 11.240  57.284 7.595e-06 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.6 p412

(55) MODEL

```

p412 = read.table("C:/G/Rt/SAS4lm/p412.txt", header=TRUE)
GLM(ts ~ source:amt, p412) # p413 Output 11.24

```

```

$ANOVA
Response : ts
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      3 393.01 131.002 903.34 < 2.2e-16 ***
RESIDUALS  16   2.32   0.145
CORRECTED TOTAL 19 395.33
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE ts Mean
0.9941306 2.619986 0.380815 14.535

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

source:amt   3 393.01     131  903.34 < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

source:amt   3 393.01     131  903.34 < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

source:amt   3 393.01     131  903.34 < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.7 p414

(56) MODEL

```

p414 = read.table("C:/G/Rt/SAS4lm/p414.txt", header=TRUE)
p414 = af(p414, c("lackofit"))
GLM(loglivcu ~ level + lackofit, p414) # p415 Output 11.26

```

```

$ANOVA  

Response : loglivcu  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL          3 5.2310 1.74365 155.47 5.018e-14 ***  

RESIDUALS       20 0.2243 0.01122  

CORRECTED TOTAL 23 5.4553  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$Fitness  

  R-square Coef Var Root MSE loglivcu Mean  

  0.9588819 6.051026 0.1059034      1.750172  

  

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

level          1 4.9859 4.9859 444.555 3.997e-15 ***  

lackofit       2 0.2450 0.1225 10.924 0.0006216 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

level      0  

lackofit  2 0.24504 0.12252  10.924 0.0006216 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

level      0  

lackofit  2 0.24504 0.12252  10.924 0.0006216 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.8 p417

(57) MODEL

```

p417 = read.table("C:/G/Rt/SAS4lm/p417.txt", header=TRUE)
p417 = af(p417, c("TRT", "POT", "PLANT"))
GLM(Y ~ TRT + POT %in% TRT, p417) # p418 Output 11.28

```

```

$ANOVA  

Response : Y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL       7 267.226 38.175 12.433 7.522e-05 ***  

RESIDUALS   13 39.917  3.071  

CORRECTED TOTAL 20 307.143  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE    Y Mean  

0.8700388 11.35742 1.752288 15.42857

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

TRT        2 236.921 118.460 38.580 3.412e-06 ***  

TRT:POT   5  30.306   6.061   1.974     0.1499  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)

```

```

TRT      2 236.921 118.460 38.580 3.412e-06 ***
TRT:POT 5 30.306   6.061    1.974    0.1499
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

TRT      2 200.111 100.055 32.586 8.626e-06 ***  

TRT:POT 5 30.306   6.061    1.974    0.1499
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ TRT + POT %in% TRT, p417), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: Y  

      Sum Sq Df F values Pr(>F)  

TRT      22.310 1 7.266 0.01835 *  

TRT:POT 30.306 5 1.974 0.14991  

Residuals 39.917 13
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.9 p431

(58) MODEL

```

p431 = read.table("C:/G/Rt/SAS4lm/p431.txt", header=TRUE)
p431 = af(p431, c("line", "sire", "agedam", "steerno"))
GLM(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431)

```

```

$ANOVA
Response : avdlygn
      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      16 2.5275 0.157966 3.1437 0.001091 **  

RESIDUALS   48 2.4119 0.050248  

CORRECTED TOTAL 64 4.9394
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE avdlygn Mean
0.511696 9.295956 0.2241612      2.411385

$`Type I` 
      Df  Sum Sq Mean Sq F value Pr(>F)
line       2 0.38009 0.190046  3.7821 0.02983 *
line:sire   6 0.92634 0.154391  3.0726 0.01260 *
agedam     2 0.11894 0.059471  1.1835 0.31497
line:agedam 4 0.64889 0.162222  3.2284 0.02000 *
age         1 0.18349 0.183487  3.6516 0.06200 .
intlwt      1 0.26970 0.269704  5.3674 0.02483 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df  Sum Sq Mean Sq F value Pr(>F)
line       2 0.05526 0.02763  0.5498 0.580636
line:sire   6 0.97389 0.16231  3.2303 0.009543 **
agedam     2 0.33106 0.16553  3.2943 0.045640 *
line:agedam 4 0.45343 0.11336  2.2560 0.076821 .
age         1 0.38128 0.38128  7.5878 0.008277 **
intlwt      1 0.26970 0.26970  5.3674 0.024830 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df  Sum Sq Mean Sq F value Pr(>F)
line       2 0.13620 0.06810  1.3553 0.267560
line:sire   6 0.97389 0.16231  3.2303 0.009543 **
agedam     2 0.13011 0.06505  1.2946 0.283392
line:agedam 4 0.45343 0.11336  2.2560 0.076821 .
age         1 0.38128 0.38128  7.5878 0.008277 **
intlwt      1 0.26970 0.26970  5.3674 0.024830 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

p433 Output 11.40

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlwt, p431),
      type=3, singular.ok=TRUE) # NOT OK for line

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: avdlygn
          Sum Sq Df F values   Pr(>F)
line      0.00000  0
agedam   0.13011  2  1.2946  0.283392
age       0.38128  1  7.5878  0.008277 ** 
intlw    0.26970  1  5.3674  0.024830 *  
line:sire 0.97389  6  3.2303  0.009543 ** 
line:agedam 0.45343  4  2.2560  0.076821 .
Residuals 2.41192 48
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(59) MODEL

```
GLM(avdlygn ~ sire + agedam, p431) # # p434 Output 11.41
```

```

$ANOVA
Response : avdlygn
          Df Sum Sq Mean Sq F value   Pr(>F)
MODEL      10 1.4254 0.142538  2.1904 0.03237 *
RESIDUALS  54 3.5140 0.065074
CORRECTED TOTAL 64 4.9394
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE avdlygn Mean
0.2885747 10.57882 0.2550961      2.411385

```

```

$`Type I` 
          Df Sum Sq Mean Sq F value   Pr(>F)
sire      8 1.30644 0.163305  2.5095 0.02138 *
agedam   2 0.11894 0.059471  0.9139 0.40707
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II` 
          Df Sum Sq Mean Sq F value   Pr(>F)
sire      8 1.33017 0.166271  2.5551 0.01937 *
agedam   2 0.11894 0.059471  0.9139 0.40707
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III` 
          Df Sum Sq Mean Sq F value   Pr(>F)
sire      8 1.33017 0.166271  2.5551 0.01937 *

```

```

agedam 2 0.11894 0.059471 0.9139 0.40707
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.10 p437 ABSORB option in SAS

(60) MODEL

```
GLM(avdlygn ~ line + sire + agedam + line:agedam + age + intlw, p431)
```

```

$ANOVA
Response : avdlygn
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      16 2.5275 0.157966 3.1437 0.001091 **
RESIDUALS   48 2.4119 0.050248
CORRECTED TOTAL 64 4.9394
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE avdlygn Mean
0.511696 9.295956 0.2241612      2.411385

```

```

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
line      2 0.38009 0.190046 3.7821 0.02983 *
sire      6 0.92634 0.154391 3.0726 0.01260 *
agedam    2 0.11894 0.059471 1.1835 0.31497
line:agedam 4 0.64889 0.162222 3.2284 0.02000 *
age       1 0.18349 0.183487 3.6516 0.06200 .
intlw     1 0.26970 0.269704 5.3674 0.02483 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
line      0
sire      6 0.97389 0.16231 3.2303 0.009543 **
agedam    2 0.33106 0.16553 3.2943 0.045640 *
line:agedam 4 0.45343 0.11336 2.2560 0.076821 .
age       1 0.38128 0.38128 7.5878 0.008277 **
intlw     1 0.26970 0.26970 5.3674 0.024830 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type III`
```

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
line	0				
sire	6	0.97389	0.16231	3.2303	0.009543 **
agedam	2	0.13011	0.06505	1.2946	0.283392
line:agedam	4	0.45343	0.11336	2.2560	0.076821 .
age	1	0.38128	0.38128	7.5878	0.008277 **
intlwt	1	0.26970	0.26970	5.3674	0.024830 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

p437 Output 11.43

6 Sahai - Unbalanced

Reference

- Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.

6.1 Table 11.2

(61) MODEL

```
T11.2 = read.table("C:/G/Rt/ANOVA/T11.2.txt")
colnames(T11.2) = c("Group", "Y")
T11.2 = af(T11.2, "Group")
GLM(Y ~ Group, T11.2) # p115

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       4  80.401 20.1003  5.9884 0.0004103 ***
RESIDUALS   59 198.036  3.3565
CORRECTED TOTAL 63 278.438
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  Y Mean
  0.2887583 2.855667 1.832089 64.15625

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4  80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4  80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4  80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

6.2 Table 12.6

(62) MODEL

```
T12.6 = read.table("C:/G/Rt/ANOVA/T12.6.txt")
colnames(T12.6) = c("Location", "Family", "Y")
T12.6 = af(T12.6, c("Location", "Family"))
GLM(Y ~ Location + Family, T12.6) # p184

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       7 1.6144 0.230636  8.9562 7.223e-07 ***
RESIDUALS   45 1.1588 0.025752
CORRECTED TOTAL 52 2.7733
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE    Y Mean
0.5821469 25.55532 0.160473 0.6279434

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.74036 0.24679  9.5833 5.219e-05 ***
Family    4 0.87410 0.21852  8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.83765 0.27921 10.8426 1.753e-05 ***
Family    4 0.87410 0.21852  8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.83765 0.27921 10.8426 1.753e-05 ***
Family    4 0.87410 0.21852  8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

6.3 Table 13.6

(63) MODEL

```

T13.6 = read.table("C:/G/Rt/ANOVA/T13.6.txt")
colnames(T13.6) = c("Site", "Worker", "Y")
T13.6 = af(T13.6, c("Site", "Worker"))
GLM(Y ~ Site + Worker + Site:Worker, T13.6)

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 2643.11 240.283  60.323 < 2.2e-16 ***
RESIDUALS   35 139.42   3.983
CORRECTED TOTAL 46 2782.52
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   Y Mean
0.9498962 2.370629 1.995817 84.18936

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site       2 1281.55  640.77 160.866 < 2.2e-16 ***
Worker     3 399.27  133.09  33.412 2.234e-10 ***
Site:Worker 6 962.29  160.38  40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site       2 1322.24  661.12 165.973 < 2.2e-16 ***
Worker     3 399.27  133.09  33.412 2.234e-10 ***
Site:Worker 6 962.29  160.38  40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site       2 804.83  402.42 101.026 2.887e-15 ***
Worker     3 430.88  143.63  36.058 8.310e-11 ***
Site:Worker 6 962.29  160.38  40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.4 Table 14.2

(64) MODEL

```

T14.2 = read.csv("C:/G/Rt/ANOVA/T14.2.csv")
T14.2 = T14.2[!is.na(T14.2$Y),]
T14.2 = af(T14.2, c("Day", "Machine", "Operator"))
GLM(Y ~ Day + Machine + Operator, T14.2)

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       7   6345.4  906.48  8.1297 5.931e-08 ***
RESIDUALS  110  12265.3  111.50
CORRECTED TOTAL 117  18610.6
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE   Y Mean
0.340954 5.495791 10.55946 192.1373

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Day       2  3737.8 1868.90 16.7611 4.426e-07 ***
Machine   2  2440.7 1220.33 10.9445 4.625e-05 ***
Operator  3   166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Day       2  3795.1 1897.56 17.0181 3.636e-07 ***
Machine   2  2464.8 1232.39 11.0526 4.227e-05 ***
Operator  3   166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Day       2  3795.1 1897.56 17.0181 3.636e-07 ***
Machine   2  2464.8 1232.39 11.0526 4.227e-05 ***
Operator  3   166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.5 Table 15.3

(65) MODEL

```

T15.3 = read.table("C:/G/Rt/ANOVA/T15.3.txt")
colnames(T15.3) = c("Dam", "Sire", "pH")
T15.3 = af(T15.3, c("Dam", "Sire"))
GLM(pH ~ Dam/Sire, T15.3) # p301

$ANOVA
Response : pH
      Df  Sum Sq  Mean Sq F value Pr(>F)
MODEL     36 0.25804 0.0071678 2.8977 7.2e-06 ***
RESIDUALS 123 0.30425 0.0024736
CORRECTED TOTAL 159 0.56229
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square   Coef Var   Root MSE  pH Mean
0.4589074 0.6676053 0.04973534 7.449813

$`Type I`
      Df  Sum Sq  Mean Sq F value Pr(>F)
Dam     14 0.178017 0.0127155 5.1405 1.563e-07 ***
Dam:Sire 22 0.080024 0.0036374 1.4705 0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq  Mean Sq F value Pr(>F)
Dam     14 0.178017 0.0127155 5.1405 1.563e-07 ***
Dam:Sire 22 0.080024 0.0036374 1.4705 0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq  Mean Sq F value Pr(>F)
Dam     14 0.179405 0.0128146 5.1805 1.347e-07 ***
Dam:Sire 22 0.080024 0.0036374 1.4705 0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(pH ~ Dam/Sire, T15.3), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: pH
      Sum Sq Df F values    Pr(>F)
Dam       0.081011   6  5.4584 4.898e-05 ***
Dam:Sire  0.080024  22  1.4705  0.09662 .
Residuals 0.304253 123
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.6 Table 16.3

(66) MODEL

```

T16.3 = read.csv("C:/G/Rt/ANOVA/T16.3.csv")
colnames(T16.3) = c("Plot", "Sample", "Subsample", "Residue")
T16.3 = af(T16.3, c("Plot", "Sample", "Subsample"))
GLM(Residue ~ Plot/Sample/Subsample, T16.3) # p344

```

```

$ANOVA
Response : Residue
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      54 3.1897 0.059069  5.8842 1.476e-05 ***
RESIDUALS   22 0.2208 0.010039
CORRECTED TOTAL 76 3.4106
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Residue Mean
0.9352456 19.94535 0.100193     0.5023377

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
Plot        10 1.84041 0.184041 18.3332 1.929e-08 ***
Plot:Sample  22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
Plot        10 1.84041 0.184041 18.3332 1.929e-08 ***
Plot:Sample  22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type III`  

          Df  Sum Sq  Mean Sq F value    Pr(>F)  

Plot           10 1.78686 0.178686 17.7998 2.547e-08 ***  

Plot:Sample     22 0.99175 0.045079  4.4906 0.0004209 ***  

Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts = c("contr.sum", "contr.poly"))  

Anova(lm(Residue ~ Plot/Sample/Subsample, T16.3), type=3, singular.ok=TRUE)
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```
Response: Residue  

          Sum Sq Df F values   Pr(>F)  

Plot           0.00000 0  

Plot:Sample     0.36613 11  3.3156 0.00805 **  

Plot:Sample:Subsample 0.35758 22  1.6191 0.13306  

Residuals      0.22085 22  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
# NOT OK
```

7 Federer - Variations

Reference

- Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.

7.1 Example 1.1

(67) MODEL

```
ex1.1 = read.table("C:/G/Rt/Split/Ex1.1-spex1.txt", header=TRUE)
ex1.1 = af(ex1.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex1.1)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      27 4905.7 181.694   10.75 1.994e-10 ***
RESIDUALS   36  608.5  16.902
CORRECTED TOTAL 63 5514.2
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
      R-square Coef Var Root MSE    Y Mean
0.8896527 6.215594 4.111227 66.14375

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3 223.8  74.60  4.4138  0.00963 **
A      3 194.6  64.85  3.8370  0.01756 *
R:A     9 158.2  17.58  1.0402  0.42842
B      3 4107.4 1369.13 81.0030 4.441e-16 ***
A:B     9 221.7  24.64  1.4577  0.20117
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3 223.8  74.60  4.4138  0.00963 **
A      3 194.6  64.85  3.8370  0.01756 *
R:A     9 158.2  17.58  1.0402  0.42842
B      3 4107.4 1369.13 81.0030 4.441e-16 ***
A:B     9 221.7  24.64  1.4577  0.20117
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     3   223.8   74.60   4.4138   0.00963 **  

A     3   194.6   64.85   3.8370   0.01756 *  

R:A   9   158.2   17.58   1.0402   0.42842  

B     3 4107.4 1369.13 81.0030 4.441e-16 ***  

A:B   9   221.7   24.64   1.4577   0.20117  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.2 Example 1.2

(68) MODEL

```

ex1.2 = read.table("C:/G/Rt/Split/Ex1.2-spex2.txt", header=TRUE)
ex1.2 = af(ex1.2, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex1.2)

```

```

$ANOVA  

Response : Y  

  Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL          47 35573  756.88  31.243 < 2.2e-16 ***  

RESIDUALS      48   1163   24.23  

CORRECTED TOTAL 95 36736  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE    Y Mean  

  0.9683464 19.45279  4.92196 25.30208

```

```

$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     2    38.6    19.3   0.7963 0.4568480  

A     7   763.2   109.0   4.5003 0.0006418 ***  

R:A  14  1377.2    98.4   4.0608 0.0001343 ***  

B     3 30774.3 10258.1 423.4386 < 2.2e-16 ***  

A:B  21  2620.1   124.8   5.1502 1.327e-06 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     2    38.6    19.3   0.7963 0.4568480

```

```

A     7    763.2   109.0   4.5003 0.0006418 ***
R:A  14   1377.2    98.4   4.0608 0.0001343 ***
B     3 30774.3 10258.1 423.4386 < 2.2e-16 ***
A:B  21   2620.1   124.8   5.1502 1.327e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df  Sum Sq Mean Sq F value    Pr(>F)
R     2     38.6    19.3   0.7963 0.4568480
A     7    763.2   109.0   4.5003 0.0006418 ***
R:A  14   1377.2    98.4   4.0608 0.0001343 ***
B     3 30774.3 10258.1 423.4386 < 2.2e-16 ***
A:B  21   2620.1   124.8   5.1502 1.327e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.3 Example 2.1

(69) MODEL

```

ex2.1 = read.table("C:/G/Rt/Split/sbex.txt", header=TRUE)
colnames(ex2.1) = c("Y", "R", "A", "B")
ex2.1 = af(ex2.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + R:B + A:B, ex2.1)

$ANOVA
Response : Y
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        41 274.750  6.7012  5.1475 0.0002305 ***
RESIDUALS     18  23.433  1.3019
CORRECTED TOTAL 59 298.183
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE    Y Mean
  0.921413 2.501251 1.140987 45.61667

$`Type I`
  Df  Sum Sq Mean Sq F value    Pr(>F)
R     1   2.817  2.8167  2.1636 0.1585807
A     9 77.683   8.6315  6.6302 0.0003456 ***
R:A  9 81.017   9.0019  6.9147 0.0002658 ***
B     2 35.433  17.7167 13.6088 0.0002510 ***
R:B  2 16.233   8.1167  6.2347 0.0087635 **

```

```

A:B 18 61.567 3.4204 2.6273 0.0236253 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     1  2.817  2.8167  2.1636 0.1585807  

A     9 77.683  8.6315  6.6302 0.0003456 ***  

R:A   9 81.017  9.0019  6.9147 0.0002658 ***  

B     2 35.433 17.7167 13.6088 0.0002510 ***  

R:B   2 16.233  8.1167  6.2347 0.0087635 **  

A:B 18 61.567  3.4204  2.6273 0.0236253 *  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     1  2.817  2.8167  2.1636 0.1585807  

A     9 77.683  8.6315  6.6302 0.0003456 ***  

R:A   9 81.017  9.0019  6.9147 0.0002658 ***  

B     2 35.433 17.7167 13.6088 0.0002510 ***  

R:B   2 16.233  8.1167  6.2347 0.0087635 **  

A:B 18 61.567  3.4204  2.6273 0.0236253 *  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.4 Example 2.2

(70) MODEL

```

ex2.2 = read.table("C:/G/Rt/Split/sbex2_2.txt", header=TRUE)
ex2.2 = af(ex2.2, c("Row", "Column", "R", "S"))
GLM(Y ~ Column + R + R:Column + S + S:Column + R:S, ex2.2)

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      51 10328 202.51 0.8112 0.7688
RESIDUALS  48 11982 249.63
CORRECTED TOTAL 99 22310

```

```

$Fitness
  R-square Coef Var Root MSE   Y Mean
0.4629279 1.579816 15.79971 1000.098

```

\$`Type I`

```

          Df Sum Sq Mean Sq F value Pr(>F)
Column     4 1318.6 329.66 1.3206 0.2758
R          4 1159.8 289.94 1.1615 0.3396
Column:R 16 2808.6 175.54 0.7032 0.7766
S          3 351.9 117.29 0.4699 0.7047
Column:S 12 3863.3 321.94 1.2897 0.2555
R:S        12  826.0   68.83 0.2757 0.9906

```

\$`Type II`

```

          Df Sum Sq Mean Sq F value Pr(>F)
Column     4 1318.6 329.66 1.3206 0.2758
R          4 1159.8 289.94 1.1615 0.3396
Column:R 16 2808.6 175.54 0.7032 0.7766
S          3 351.9 117.29 0.4699 0.7047
Column:S 12 3863.3 321.94 1.2897 0.2555
R:S        12  826.0   68.83 0.2757 0.9906

```

\$`Type III`

```

          Df Sum Sq Mean Sq F value Pr(>F)
Column     4 1318.6 329.66 1.3206 0.2758
R          4 1159.8 289.94 1.1615 0.3396
Column:R 16 2808.6 175.54 0.7032 0.7766
S          3 351.9 117.29 0.4699 0.7047
Column:S 12 3863.3 321.94 1.2897 0.2555
R:S        12  826.0   68.83 0.2757 0.9906

```

(71) MODEL

```
GLM(Y ~ Row + R + Row:R + S + Column:S + R:S + Column:R:S, ex2.2)
```

\$ANOVA

```

Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      99 22310 225.36
RESIDUALS    0      0
CORRECTED TOTAL 99 22310

```

\$Fitness

R-square	Coef	Var	Root	MSE	Y	Mean
1	NA		NA	1000.098		

\$`Type I`

```

          Df Sum Sq Mean Sq F value Pr(>F)
Row        4   147.4   36.86
R          4 1159.8 289.94
Row:R     16 3979.8 248.74
S          3   351.9   117.29

```

```

S:Column 12 3863.3 321.94
R:S       12  826.0   68.83
R:S:Column 48 11982.3 249.63

$`Type II` 
      Df Sum Sq Mean Sq F value Pr(>F)
Row        0
R          4 1159.8 289.94
Row:R      0
S          3  351.9 117.29
S:Column 12 3863.3 321.94
R:S       12  826.0   68.83
R:S:Column 48 11982.3 249.63

```

```

$`Type III` 
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)
Row        0
R          4 1159.8 289.94
Row:R      0
S          3  351.9 117.29
S:Column 12 3863.3 321.94
R:S       12  826.0   68.83
R:S:Column 48 11982.3 249.63

```

(72) MODEL

```
GLM(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      99 22310 225.36
RESIDUALS    0      0
CORRECTED TOTAL 99 22310
```

```
$Fitness
R-square Coef Var Root MSE   Y Mean
      1       NA     NA 1000.098
```

```
$`Type I` 
      Df Sum Sq Mean Sq F value Pr(>F)
Row        4 147.4   36.86
R          4 1159.8 289.94
S          3  351.9 117.29
R:S       12  826.0   68.83
Row:R     16 3979.8 248.74
```

```

S:Column 12 3863.3 321.94
R:S:Column 48 11982.3 249.63

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Row 0  

R 4 1159.8 289.94  

S 3 351.9 117.29  

R:S 12 826.0 68.83  

Row:R 0  

S:Column 12 3863.3 321.94
R:S:Column 48 11982.3 249.63

$`Type III`  

CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)  

Row 0  

R 4 1159.8 289.94  

S 3 351.9 117.29  

R:S 12 826.0 68.83  

Row:R 0  

S:Column 12 3863.3 321.94
R:S:Column 48 11982.3 249.63

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2), type=3,
      singular.ok=TRUE) # NOT WORKING

```

7.5 Example 3.1

(73) MODEL

```

ex3.1 = read.table("C:/G/Rt/Split/spedsite.txt", header=TRUE)
ex3.1 = af(ex3.1, c("Site", "A", "B", "C", "Block"))
GLM(Yield ~ Site + Site:Block + A + B + A:B + A:Site + B:Site + A:B:Site +
     A:B:Site:Block + C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site +
     A:B:C:Site, ex3.1)

```

```

$ANOVA
Response : Yield
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      239 2724374186 11399055 23.682 < 2.2e-16 ***
RESIDUALS   240 115521933   481341
CORRECTED TOTAL 479 2839896119
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Yield Mean
  0.9593218 8.368195 693.7877    8290.769

$`Type I`
      Df   Sum Sq  Mean Sq  F value  Pr(>F)
Site        3 621230991 207076997 430.2082 < 2e-16 ***
Site:Block  8 1305369943 163171243 338.9928 < 2e-16 ***
A           1 1333205   1333205   2.7698  0.09737 .
B           4 47928577  11982144  24.8932 < 2e-16 ***
A:B         4 14849     3712    0.0077  0.99988
Site:A      3 33010     11003    0.0229  0.99531
Site:B      12 37932     3161    0.0066  1.00000
Site:A:B    12 11494     958    0.0020  1.00000
Site:Block:A:B 72 8239680   114440   0.2378  1.00000
C           3 739890389 246630130 512.3809 < 2e-16 ***
A:C         3 3233      1078    0.0022  0.99985
B:C         12 34961     2913    0.0061  1.00000
A:B:C       12 11077     923    0.0019  1.00000
Site:C      9 25983     2887    0.0060  1.00000
Site:A:C    9 22227     2470    0.0051  1.00000
Site:B:C    36 88610     2461    0.0051  1.00000
Site:A:B:C  36 98025     2723    0.0057  1.00000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df   Sum Sq  Mean Sq  F value  Pr(>F)
Site        3 621230991 207076997 430.2082 < 2e-16 ***
Site:Block  8 1305369943 163171243 338.9928 < 2e-16 ***
A           1 1333205   1333205   2.7698  0.09737 .
B           4 47928577  11982144  24.8932 < 2e-16 ***
A:B         4 14849     3712    0.0077  0.99988
Site:A      3 33010     11003    0.0229  0.99531
Site:B      12 37932     3161    0.0066  1.00000
Site:A:B    12 11494     958    0.0020  1.00000
Site:Block:A:B 72 8239680   114440   0.2378  1.00000
C           3 739890389 246630130 512.3809 < 2e-16 ***
A:C         3 3233      1078    0.0022  0.99985
B:C         12 34961     2913    0.0061  1.00000
A:B:C       12 11077     923    0.0019  1.00000
Site:C      9 25983     2887    0.0060  1.00000
Site:A:C    9 22227     2470    0.0051  1.00000
Site:B:C    36 88610     2461    0.0051  1.00000
Site:A:B:C  36 98025     2723    0.0057  1.00000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type III`  

          Df Sum Sq Mean Sq F value Pr(>F)  

Site        3 621230991 207076997 430.2082 < 2e-16 ***  

Site:Block  8 1305369943 163171243 338.9928 < 2e-16 ***  

A           1 1333205 1333205  2.7698 0.09737 .  

B           4 47928577 11982144 24.8932 < 2e-16 ***  

A:B         4 14849    3712  0.0077 0.99988  

Site:A      3 33010    11003  0.0229 0.99531  

Site:B      12 37932     3161  0.0066 1.00000  

Site:A:B    12 11494     958  0.0020 1.00000  

Site:Block:A:B 72 8239680 114440  0.2378 1.00000  

C           3 739890389 246630130 512.3809 < 2e-16 ***  

A:C         3 3233     1078  0.0022 0.99985  

B:C         12 34961     2913  0.0061 1.00000  

A:B:C       12 11077     923  0.0019 1.00000  

Site:C      9 25983     2887  0.0060 1.00000  

Site:A:C    9 22227     2470  0.0051 1.00000  

Site:B:C    36 88610     2461  0.0051 1.00000  

Site:A:B:C  36 98025     2723  0.0057 1.00000  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(74) MODEL

```
ex3.1a = read.table("C:/G/Rt/Split/Ex3.1-example.txt", header=TRUE)  

ex3.1a = af(ex3.1a, c("row", "P", "column", "R", "S"))  

GLM(height ~ P + column + column:P + R + P:R + column:R + column:R:P + S +  

  P:S + column:S + column:S:P + R:S + R:S:column + R:S:P + R:S:P:column, ex3.1a)
```

```
$ANOVA  

Response : height  

          Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      199 7534.8 37.863  

RESIDUALS   0     0.0  

CORRECTED TOTAL 199 7534.8
```

```
$Fitness  

R-square Coef Var Root MSE height Mean  

1       NA     NA    93.965
```

```
$`Type I`  

          Df Sum Sq Mean Sq F value Pr(>F)  

P           1 253.1 253.125  

column      4 109.4 27.357  

P:column    4 207.9 51.987  

R           4  90.6 22.657
```

P:R	4	505.0	126.238
column:R	16	3357.8	209.864
P:column:R	16	1442.6	90.163
S	3	16.4	5.458
P:S	3	14.3	4.765
column:S	12	265.4	22.121
P:column:S	12	96.5	8.044
R:S	12	195.1	16.254
column:R:S	48	365.5	7.615
P:R:S	12	100.3	8.361
P:column:R:S	48	514.7	10.723

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	253.1	253.125		
column	4	109.4	27.358		
P:column	4	208.0	51.988		
R	4	90.6	22.657		
P:R	4	504.9	126.237		
column:R	16	3357.8	209.864		
P:column:R	16	1442.6	90.162		
S	3	16.4	5.458		
P:S	3	14.3	4.765		
column:S	12	265.5	22.121		
P:column:S	12	96.5	8.044		
R:S	12	195.0	16.254		
column:R:S	48	365.5	7.615		
P:R:S	12	100.3	8.361		
P:column:R:S	48	514.7	10.723		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	253.1	253.125		
column	4	109.4	27.358		
P:column	4	208.0	51.988		
R	4	90.6	22.657		
P:R	4	505.0	126.238		
column:R	16	3357.8	209.864		
P:column:R	16	1442.6	90.163		
S	3	16.4	5.458		
P:S	3	14.3	4.765		
column:S	12	265.4	22.121		
P:column:S	12	96.5	8.044		
R:S	12	195.0	16.254		
column:R:S	48	365.5	7.615		
P:R:S	12	100.3	8.361		
P:column:R:S	48	514.7	10.723		

(75) MODEL

```
GLM(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
     S:R:P + R:S:P:row, ex3.1a)
```

\$ANOVA

```
Response : height
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          199 7534.8 37.863
RESIDUALS      0    0.0
CORRECTED TOTAL 199 7534.8
```

\$Fitness

```
R-square Coef Var Root MSE height Mean
        1       NA       NA      93.965
```

\$`Type I`

```
Df Sum Sq Mean Sq F value Pr(>F)
row      4 2017.03 504.26
R        4   90.63  22.66
P        1 253.12 253.12
S        3   16.38   5.46
R:S      12 195.05 16.25
row:P    4   167.25 41.81
R:P      4 504.95 126.24
row:R:P  32 2933.52 91.67
P:S      3   14.29   4.76
row:P:S  24 234.68  9.78
R:P:S    12 100.33   8.36
row:R:P:S 96 1007.52 10.49
```

\$`Type II`

```
Df Sum Sq Mean Sq F value Pr(>F)
row      4 2017.03 504.26
R        4   90.63  22.66
P        1 253.12 253.12
S        3   16.38   5.46
R:S      12 195.05 16.25
row:P    4   167.25 41.81
R:P      4 504.95 126.24
row:R:P  32 2933.52 91.67
P:S      3   14.29   4.76
row:P:S  24 234.68  9.78
R:P:S    12 100.33   8.36
row:R:P:S 96 1007.52 10.49
```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
row	4	2017.03	504.26		
R	4	90.63	22.66		
P	1	253.12	253.12		
S	3	16.38	5.46		
R:S	12	195.05	16.25		
row:P	4	167.25	41.81		
R:P	4	504.95	126.24		
row:R:P	32	2933.52	91.67		
P:S	3	14.30	4.77		
row:P:S	24	234.68	9.78		
R:P:S	12	100.33	8.36		
row:R:P:S	96	1007.52	10.50		

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
S:P:row + S:R:P + R:S:P:row, ex3.1a), type=3, singular.ok=TRUE)
# NOT WORKING
```

```
alias(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
S:R:P + R:S:P:row, ex3.1a) # NO ALIAS
```

Model :
height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
S:P:row + S:R:P + R:S:P:row

(76) MODEL

- p94 Appendix 3.1

```
ex3.1b = read.table("C:/G/Rt/Split/spexvar3.txt", header=TRUE)
ex3.1b = af(ex3.1b, c("rep", "var", "nit", "row", "col"))
GLM(yield ~ rep + var + rep:var + nit + var:nit, ex3.1b)
```

```
$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      26  44017 1692.97  9.5603 4.779e-11 ***
RESIDUALS   45   7969  177.08
CORRECTED TOTAL 71  51986
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE yield Mean
```

```

0.8467134 12.79887 13.30727    103.9722

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      5 15875.3  3175.1 17.9297 9.525e-10 ***  

var      2   1786.4    893.2  5.0438  0.010557 *  

rep:var 10  6013.3   601.3  3.3957  0.002251 **  

nit      3 20020.5  6673.5 37.6856 2.458e-12 ***  

var:nit  6   321.7    53.6  0.3028  0.932199  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      5 15875.3  3175.1 17.9297 9.525e-10 ***  

var      2   1786.4    893.2  5.0438  0.010557 *  

rep:var 10  6013.3   601.3  3.3957  0.002251 **  

nit      3 20020.5  6673.5 37.6856 2.458e-12 ***  

var:nit  6   321.7    53.6  0.3028  0.932199  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      5 15875.3  3175.1 17.9297 9.525e-10 ***  

var      2   1786.4    893.2  5.0438  0.010557 *  

rep:var 10  6013.3   601.3  3.3957  0.002251 **  

nit      3 20020.5  6673.5 37.6856 2.458e-12 ***  

var:nit  6   321.7    53.6  0.3028  0.932199  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(77) MODEL

```
GLM(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b)
```

```
$ANOVA  

Response : yield  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL       37  48090  1299.7 11.341 6.734e-11 ***  

RESIDUALS    34   3896   114.6  

CORRECTED TOTAL 71  51986  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

R-square Coef Var Root MSE yield Mean
```

```

0.9250491 10.29615 10.70513   103.9722

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      5 15875.3  3175.1 27.7056 4.391e-11 ***  

var      2  1786.4   893.2  7.7939 0.0016359 **  

rep:var 10  6013.3   601.3  5.2472 0.0001207 ***  

nit      3 20020.5  6673.5 58.2331 1.754e-13 ***  

var:nit  6   321.7    53.6  0.4679 0.8271333  

row      9   900.9   100.1  0.8734 0.5575581  

col      2  3171.5  1585.7 13.8373 4.012e-05 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      2 5942.5  2971.3 25.9273 1.449e-07 ***  

var      2 2799.8  1399.9 12.2155 0.0001005 ***  

rep:var  4   997.8   249.4  2.1767 0.0926008 .  

nit      3 12559.3  4186.4 36.5308 9.683e-11 ***  

var:nit  6   477.8    79.6  0.6949 0.6553307  

row      9   945.0   105.0  0.9162 0.5230151  

col      2  3171.5  1585.7 13.8373 4.012e-05 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep      2 5942.5  2971.3 25.9273 1.449e-07 ***  

var      2 2799.8  1399.9 12.2155 0.0001005 ***  

rep:var  4   997.8   249.4  2.1767 0.0926008 .  

nit      3 11977.9  3992.6 34.8397 1.775e-10 ***  

var:nit  6   477.8    79.6  0.6949 0.6553307  

row      9   945.0   105.0  0.9162 0.5230151  

col      2  3171.5  1585.7 13.8373 4.012e-05 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b),
      type=3, singular.ok=TRUE) # NOT OK for var

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: yield
            Sum Sq Df F values    Pr(>F)
rep      5942.5  2 25.9273 1.449e-07 ***
var        0.0  0
nit     11977.9  3 34.8397 1.775e-10 ***
row      945.0  9  0.9162    0.5230
col     3171.5  2 13.8373 4.012e-05 ***
rep:var   997.8  4  2.1767    0.0926 .
var:nit   477.8  6  0.6949    0.6553
Residuals 3896.4 34
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.6 Example 4.1

(78) MODEL

```

ex4.1 = read.table("C:/G/Rt/Split/Ex4.1-example.txt", header=TRUE)
ex4.1 = af(ex4.1, c("row", "P", "column", "R", "S"))
GLM(height ~ P + column + column:P + R + P:R + column:R + column:R:P + S +
     P:S + column:S + column:S:P + R:S + R:S:column + R:S:P + R:S:P:column, ex4.1)

```

```

$ANOVA
Response : height
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          199 1710.2 8.5937
RESIDUALS       0    0.0
CORRECTED TOTAL 199 1710.2

```

```

$Fitness
R-square Coef Var Root MSE height Mean
1         NA    NA    6.815

```

```

$`Type I`
            Df Sum Sq Mean Sq F value Pr(>F)
P             1 28.12 28.1250
column        4 34.33 8.5825
P:column      4 91.45 22.8625
R              4 31.03 7.7575
P:R            4 48.95 12.2375
column:R      16 467.92 29.2450
P:column:R    16 350.10 21.8813
S              3  3.77 1.2583
P:S            3  3.29 1.0983
column:S      12 74.55 6.2125

```

P:column:S	12	47.03	3.9192
R:S	12	36.65	3.0542
column:R:S	48	197.40	4.1125
P:R:S	12	26.33	2.1942
P:column:R:S	48	269.22	5.6087

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	28.13	28.1250		
column	4	34.33	8.5825		
P:column	4	91.45	22.8625		
R	4	31.03	7.7575		
P:R	4	48.95	12.2375		
column:R	16	467.92	29.2450		
P:column:R	16	350.10	21.8812		
S	3	3.77	1.2583		
P:S	3	3.30	1.0983		
column:S	12	74.55	6.2125		
P:column:S	12	47.03	3.9192		
R:S	12	36.65	3.0542		
column:R:S	48	197.40	4.1125		
P:R:S	12	26.33	2.1942		
P:column:R:S	48	269.22	5.6087		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	28.12	28.1250		
column	4	34.33	8.5825		
P:column	4	91.45	22.8625		
R	4	31.03	7.7575		
P:R	4	48.95	12.2375		
column:R	16	467.92	29.2450		
P:column:R	16	350.10	21.8813		
S	3	3.77	1.2583		
P:S	3	3.29	1.0983		
column:S	12	74.55	6.2125		
P:column:S	12	47.03	3.9192		
R:S	12	36.65	3.0542		
column:R:S	48	197.40	4.1125		
P:R:S	12	26.33	2.1942		
P:column:R:S	48	269.22	5.6088		

(79) MODEL

```
GLM(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
S:R:P + R:S:P:row, ex4.1)
```

\$ANOVA

```

Response : height
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL           199 1710.2 8.5937
RESIDUALS        0    0.0
CORRECTED TOTAL 199 1710.2

```

```

$Fitness
R-square Coef Var Root MSE height Mean
      1       NA       NA      6.815

```

```

$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
row        4 309.43 77.357
R          4 31.03 7.758
P          1 28.12 28.125
S          3 3.77 1.258
R:S        12 36.65 3.054
row:P      4 130.25 32.563
R:P        4 48.95 12.237
row:R:P    32 504.12 15.754
P:S        3 3.29 1.098
row:P:S   24 171.28 7.137
R:P:S     12 26.33 2.194
row:R:P:S 96 416.92 4.343

```

```

$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
row        4 309.43 77.357
R          4 31.03 7.757
P          1 28.12 28.125
S          3 3.78 1.258
R:S        12 36.65 3.054
row:P      4 130.25 32.563
R:P        4 48.95 12.238
row:R:P    32 504.12 15.754
P:S        3 3.30 1.098
row:P:S   24 171.28 7.137
R:P:S     12 26.33 2.194
row:R:P:S 96 416.92 4.343

```

```

$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
row        4 309.43 77.358
R          4 31.03 7.757
P          1 28.13 28.125
S          3 3.78 1.258
R:S        12 36.65 3.054
row:P      4 130.25 32.563

```

```

R:P          4  48.95 12.237
row:R:P     32 504.12 15.754
P:S          3   3.30  1.098
row:P:S    24 171.28 7.137
R:P:S       12  26.33 2.194
row:R:P:S  96 416.92 4.343

```

7.7 Example 5.1

(80) MODEL

```

ex5.1 = read.table("C:/G/Rt/Split/sbsp.txt", header=TRUE)
ex5.1 = af(ex5.1, c("R", "A", "C", "B", "Tx"))
GLM(Y ~ R + A + R:A + C + B + C:B + Tx + B:Tx, ex5.1)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      20 193.583 9.6792 9.4176 2.969e-05 ***
RESIDUALS   15 15.417  1.0278
CORRECTED TOTAL 35 209.000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE Y Mean
0.926236 18.43261 1.013794      5.5

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
R      2 33.500 16.7500 16.2973 0.0001734 ***
A      1 16.000 16.0000 15.5676 0.0012951 **
R:A    2 32.167 16.0833 15.6486 0.0002133 ***
C      2   0.500  0.2500  0.2432 0.7871141
B      1   1.778  1.7778  1.7297 0.2081966
C:B    2   0.389  0.1944  0.1892 0.8295745
Tx     5 103.333 20.6667 20.1081 3.63e-06 ***
B:Tx   5   5.917  1.1833  1.1514 0.3770453
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
R      2 23.047 11.5236 11.2122 0.0010520 **
A      1 12.375 12.3751 12.0406 0.0034285 **
R:A    2 27.164 13.5819 13.2148 0.0004907 ***

```

```

C      2    0.500  0.2500  0.2432  0.7871141
B      1    1.778  1.7778  1.7297  0.2081966
C:B    2    0.389  0.1944  0.1892  0.8295745
Tx     5  103.333 20.6667 20.1081  3.63e-06 ***
B:Tx   5    5.917  1.1833  1.1514  0.3770453
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

R     2   22.451 11.2254 10.9220 0.0011828 **  

A     1   15.001 15.0013 14.5958 0.0016719 **  

R:A    2   27.164 13.5819 13.2148 0.0004907 ***  

C     2    0.500  0.2500  0.2432  0.7871141  

B     1    1.778  1.7778  1.7297  0.2081966  

C:B    2    0.389  0.1944  0.1892  0.8295745  

Tx     5  103.333 20.6667 20.1081  3.63e-06 ***  

B:Tx   5    5.917  1.1833  1.1514  0.3770453
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(81) MODEL

```
GLM(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx, ex5.1)
```

```
$ANOVA  

Response : Y  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL        20 194.188  9.7094  9.8323 2.254e-05 ***  

RESIDUALS    15  14.813  0.9875  

CORRECTED TOTAL 35 209.000
---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE Y Mean  

0.9291268 18.06782 0.9937303      5.5
```

```
$`Type I`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

R     2   33.500 16.7500 16.9620 0.0001410 ***  

A     1   16.000 16.0000 16.2025 0.0011013 **  

R:A    2   32.167 16.0833 16.2869 0.0001739 ***  

C     2    0.500  0.2500  0.2532  0.7795913  

B     1    1.778  1.7778  1.8003  0.1996385  

C:B    2    0.389  0.1944  0.1969  0.8233570  

Tx     5  103.333 20.6667 20.9283  2.813e-06 ***
```

```

A:Tx 5 6.521 1.3042 1.3207 0.3078554
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R     2 33.500 16.7500 16.9620 0.0001410 ***  

A     1 16.000 16.0000 16.2025 0.0011013 **  

R:A   2 32.167 16.0833 16.2869 0.0001739 ***  

C     2  0.807  0.4037  0.4088 0.6716130  

B     1  1.757  1.7574  1.7797 0.2020905  

C:B   2  0.030  0.0150  0.0152 0.9849064  

Tx    5 103.333 20.6667 20.9283 2.813e-06 ***  

A:Tx 5 6.521 1.3042 1.3207 0.3078554
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R     2 33.500 16.7500 16.9620 0.0001410 ***  

A     1 16.000 16.0000 16.2025 0.0011013 **  

R:A   2 32.167 16.0833 16.2869 0.0001739 ***  

C     2  0.780  0.3902  0.3952 0.6803789  

B     1  1.776  1.7756  1.7980 0.1999029  

C:B   2  0.030  0.0150  0.0152 0.9849064  

Tx    5 103.333 20.6667 20.9283 2.813e-06 ***  

A:Tx 5 6.521 1.3042 1.3207 0.3078554
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(82) MODEL

```
GLM(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

```
$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      24 196.238 8.1766 7.0476 0.0008758 ***  

RESIDUALS    11 12.762 1.1602  

CORRECTED TOTAL 35 209.000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE Y Mean  

0.9389372 19.58405 1.077122      5.5
```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R     2   33.500 16.7500 14.4373 0.0008391 ***  

A     1   16.000 16.0000 13.7908 0.0034197 **  

R:A   2   32.167 16.0833 13.8626 0.0009856 ***  

C     2    0.500  0.2500  0.2155 0.8094766  

B     1    1.778  1.7778  1.5323 0.2415358  

C:B   2    0.389  0.1944  0.1676 0.8478141  

Tx    5 103.333 20.6667 17.8131 6.055e-05 ***  

A:Tx  5    6.521  1.3042  1.1241 0.4027183  

B:Tx  4    2.050  0.5126  0.4418 0.7761730  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R     2   23.116 11.5581 9.9622 0.003396 **  

A     1   12.375 12.3751 10.6664 0.007519 **  

R:A   2   27.426 13.7132 11.8197 0.001820 **  

C     2    0.970  0.4850  0.4180 0.668392  

B     1    1.757  1.7574  1.5148 0.244080  

C:B   2    0.085  0.0424  0.0366 0.964202  

Tx    5 103.333 20.6667 17.8131 6.055e-05 ***  

A:Tx  4    2.655  0.6636  0.5720 0.688652  

B:Tx  4    2.050  0.5126  0.4418 0.776173  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R     2   22.186 11.0928 9.5611 0.003924 **  

A     1   15.185 15.1853 13.0886 0.004042 **  

R:A   2   27.426 13.7132 11.8197 0.001820 **  

C     2    1.010  0.5049  0.4352 0.657839  

B     1    1.792  1.7922  1.5448 0.239751  

C:B   2    0.085  0.0424  0.0366 0.964202  

Tx    5 103.333 20.6667 17.8131 6.055e-05 ***  

A:Tx  4    2.655  0.6636  0.5720 0.688652  

B:Tx  4    2.050  0.5126  0.4418 0.776173  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
alias(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

```
Model :  
Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx
```

```

Complete :
      (Intercept) R1     R2     A1     C1     C2     B1     Tx1    Tx2    Tx3    Tx4    Tx5    R1:A1
B1:Tx5      0       0     -1/5     0       0     -1/5     0       0     0       0     0       0
      R2:A1  C1:B1  C2:B1  A1:Tx1  A1:Tx2  A1:Tx3  A1:Tx4  A1:Tx5  B1:Tx1  B1:Tx2  B1:Tx3
B1:Tx5      0       0     1/5     1/5     1/5     1/5     -1     1/5     1/5     1/5     1/5
      B1:Tx4
B1:Tx5     1/5

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F value	Pr(>F)							
R	22.186	2	9.5611	0.003924 **							
A	0.000	0									
C	1.010	2	0.4352	0.657839							
B	0.000	0									
Tx	103.333	5	17.8131	6.055e-05 ***							
R:A	27.426	2	11.8197	0.001820 **							
C:B	0.085	2	0.0366	0.964202							
A:Tx	2.655	4	0.5720	0.688652							
B:Tx	2.050	4	0.4418	0.776173							
Residuals	12.762	11									

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

(83) MODEL

```
GLM(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

```

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      28 204.2  7.2929 10.635 0.001719 **
RESIDUALS   7   4.8  0.6857
CORRECTED TOTAL 35 209.0
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Y Mean
  0.9770335 15.05598 0.8280787      5.5

$`Type I` 
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 33.500 16.7500 24.4271 0.0006969 ***
A     1 16.000 16.0000 23.3333 0.0018985 **
R:A   2 32.167 16.0833 23.4549 0.0007889 ***
C     2  0.500  0.2500  0.3646 0.7069339
B     1  1.778  1.7778  2.5926 0.1513998
C:B   2  0.389  0.1944  0.2836 0.7613494
Tx    5 103.333 20.6667 30.1389 0.0001357 ***
A:Tx  5  6.521  1.3042  1.9019 0.2123307
B:Tx  4  2.050  0.5126  0.7475 0.5896365
A:B:Tx 4  7.962  1.9905  2.9029 0.1038803
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 31.838 15.9191 23.2153 0.0008139 ***
A     1 12.375 12.3751 18.0470 0.0038017 **
R:A   1  2.017  2.0174  2.9420 0.1300172
C     2  0.500  0.2500  0.3645 0.7069558
B     1  1.757  1.7574  2.5629 0.1534298
C:B   1  0.644  0.6445  0.9399 0.3646045
Tx    5 103.333 20.6667 30.1389 0.0001357 ***
A:Tx  4  2.655  0.6636  0.9678 0.4812226
B:Tx  4  2.050  0.5126  0.7475 0.5896365
A:B:Tx 4  7.962  1.9905  2.9029 0.1038803
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 28.112 14.0562 20.4986 0.0011846 **
A     1 14.655 14.6551 21.3720 0.0024176 **
R:A   1  2.017  2.0174  2.9420 0.1300172
C     2  0.471  0.2356  0.3436 0.7205632
B     1  1.769  1.7694  2.5804 0.1522328
C:B   1  0.644  0.6445  0.9399 0.3646045
Tx    5 103.815 20.7630 30.2793 0.0001336 ***
A:Tx  4  2.951  0.7378  1.0760 0.4358837
B:Tx  4  3.553  0.8882  1.2954 0.3579988
A:B:Tx 4  7.962  1.9905  2.9029 0.1038803
---

```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
alias(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

Model :

```
Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx
```

Complete :

	(Intercept)	R1	R2	A1	C1	C2	B1	Tx1	Tx2	Tx3	Tx4	Tx5
B1:Tx5	0	0	0	-1/5	0	0	-1/5	0	0	0	0	0
A1:B1:Tx5	-1/6	0	0	0	0	0	0	1/6	1/6	1/6	1/6	-5/6
A1:B1:Tx6	0	2/3	0	4/45	2/3	-2/3	4/45	-1/3	1/3	-1/3	0	0
R1:A1	R2:A1	C1:B1	C2:B1	A1:Tx1	A1:Tx2	A1:Tx3	A1:Tx4	A1:Tx5	B1:Tx1			
B1:Tx5	0	0	0	0	1/5	1/5	1/5	1/5	-1	1/5		
A1:B1:Tx5	0	0	0	0	0	0	0	0	0	0		
A1:B1:Tx6	-2/9	4/9	-2/9	-2/9	-1/5	-1/5	-1/5	4/5	0	-1/5		
B1:Tx2	B1:Tx3	B1:Tx4	A1:B1:Tx1	A1:B1:Tx2	A1:B1:Tx3	A1:B1:Tx4						
B1:Tx5	1/5	1/5	1/5	0	0	0	0	0	0	0		
A1:B1:Tx5	0	0	0	0	0	0	0	0	0	0		
A1:B1:Tx6	-1/5	-1/5	4/5	1	-1		1		0			

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)
R	11.643	1	16.9793	0.004456 **
A	0.000	0		
C	0.002	1	0.0025	0.961483
B	0.000	0		
Tx	89.178	3	43.3503	6.87e-05 ***
R:A	2.017	1	2.9420	0.130017
C:B	0.644	1	0.9399	0.364604
A:Tx	0.543	3	0.2640	0.849381
B:Tx	3.384	3	1.6451	0.264128
A:B:Tx	7.962	4	2.9029	0.103880
Residuals	4.800	7		

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.8 Example 7.1

(84) MODEL

```
ex7.1 = read.table("C:/G/Rt/Split/asped.txt", header=TRUE)
ex7.1 = af(ex7.1, c("R", "G", "F"))
GLM(Y ~ R + G + R:G + F + F:G, ex7.1)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      95 577.83  6.0824  5.3082 1.068e-05 ***
RESIDUALS   24  27.50  1.1458
CORRECTED TOTAL 119 605.33
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
  R-square Coef Var Root MSE Y Mean
0.9545699 17.335 1.070436 6.175
```

```
$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3  84.76 28.2528 24.6570 1.655e-07 ***
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9  11.75  1.3056  1.1394    0.3749
F      2  59.85 29.9250 26.1164 9.481e-07 ***
G:F   54  77.98  1.4441  1.2603    0.2718
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3   5.75  1.9167  1.6727    0.1994
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9  11.75  1.3056  1.1394    0.3749
F      2  59.85 29.9250 26.1164 9.481e-07 ***
G:F   54  77.98  1.4441  1.2603    0.2718
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3   5.75  1.9167  1.6727    0.1994
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9  11.75  1.3056  1.1394    0.3749
F      2  50.51 25.2525 22.0385 3.686e-06 ***
```

```

G:F 54  77.98  1.4441  1.2603    0.2718
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + G + R:G + F + F:G, ex7.1), type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y
          Sum Sq Df F values   Pr(>F)
R           0.000  0
G        202.417  3 58.8848 3.258e-11 ***
F         50.505  2 22.0385 3.686e-06 ***
R:G       11.750  9  1.1394    0.3749
G:F       77.983 54  1.2603    0.2718
Residuals 27.500 24
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.9 Example 7.2

(85) MODEL

```

ex7.2 = read.table("C:/G/Rt/Split/aspedt.txt", header=TRUE)
ex7.2 = af(ex7.2, c("R", "T", "G"))
GLM(Y ~ R + T + R:T + G + G:T, ex7.2)

```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value   Pr(>F)
MODEL      99 538.70  5.4415  5.1892 1.286e-05 ***
RESIDUALS  24  25.17  1.0486
CORRECTED TOTAL 123 563.87
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE   Y Mean
  0.955368 16.97569 1.024017 6.032258

```

```
$`Type I`
```

```

      Df  Sum Sq Mean Sq F value    Pr(>F)
R      3   73.255 24.4183 23.2863 2.752e-07 ***
T      3   32.000 10.6667 10.1722 0.0001645 ***
R:T    9   28.402  3.1558  3.0095 0.0149568 *
G     21 309.908 14.7575 14.0734 7.158e-09 ***
T:G   63   95.140  1.5102  1.4401 0.1617931
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
R      3   4.229  1.4097  1.3444 0.2834998
T      3   32.000 10.6667 10.1722 0.0001645 ***
R:T    9   10.854  1.2060  1.1501 0.3684706
G     21 309.908 14.7575 14.0734 7.158e-09 ***
T:G   63   95.140  1.5102  1.4401 0.1617931
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
R      3   4.229  1.4097  1.3444 0.283500
T      3   22.668  7.5559  7.2056 0.001299 **
R:T    9   10.854  1.2060  1.1501 0.368471
G     21 309.908 14.7575 14.0734 7.158e-09 ***
T:G   63   95.140  1.5102  1.4401 0.161793
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.10 Example 7.3

(86) MODEL

```

ex7.3 = read.table("C:/G/Rt/Split/assped.txt", header=TRUE)
ex7.3 = af(ex7.3, c("R", "T", "G", "F"))
GLM(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL          155 656.12  4.2330 13.446 3.997e-14 ***
RESIDUALS       36  11.33  0.3148
CORRECTED TOTAL 191 667.45
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE Y Mean
0.98302 8.95495 0.5610836 6.265625

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
R       3 27.06   9.019  28.6489 1.203e-09 ***
T       1 10.55  10.547  33.5018 1.334e-06 ***
R:T     3  2.97   0.991   3.1489  0.036705 *
G      22 389.01  17.682  56.1668 < 2.2e-16 ***
T:G    22 18.42   0.837   2.6601  0.004445 **
R:T:G  12  8.78   0.731   2.3235  0.025315 *
F       2 164.28  82.141  260.9173 < 2.2e-16 ***
T:F    2   0.84   0.422   1.3401  0.274574
G:F    44 23.47   0.533   1.6943  0.053191 .
T:G:F  44 10.74   0.244   0.7753  0.790640
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
R       3 12.49   4.162  13.2206 5.655e-06 ***
T       1 10.55  10.547  33.5018 1.334e-06 ***
R:T     3  1.15   0.384   1.2206  0.316281
G      22 389.01  17.682  56.1668 < 2.2e-16 ***
T:G    22 18.42   0.837   2.6601  0.004445 **
R:T:G  12  8.78   0.731   2.3235  0.025315 *
F       2 164.28  82.141  260.9173 < 2.2e-16 ***
T:F    2   0.84   0.422   1.3401  0.274574
G:F    44 23.47   0.533   1.6943  0.053191 .
T:G:F  44 10.74   0.244   0.7753  0.790640
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
R       3 12.49   4.162  13.2206 5.655e-06 ***
T       1 11.16  11.158  35.4430 8.021e-07 ***
R:T     3  1.15   0.384   1.2206  0.316281
G      22 389.01  17.682  56.1668 < 2.2e-16 ***
T:G    22 18.42   0.837   2.6601  0.004445 **
R:T:G  12  8.78   0.731   2.3235  0.025315 *
F       2 120.56  60.282  191.4828 < 2.2e-16 ***
T:F    2   0.82   0.411   1.3060  0.283432
G:F    44 23.47   0.533   1.6943  0.053191 .
T:G:F  44 10.74   0.244   0.7753  0.790640
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3),
      type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)							
R	0.000	0									
T	0.000	0									
G	73.444	2	116.6471 < 2.2e-16 ***								
F	120.563	2	191.4828 < 2.2e-16 ***								
R:T	0.000	0									
T:G	5.778	2	9.1765 0.0006018 ***								
T:F	0.822	2	1.3060 0.2834316								
G:F	23.469	44	1.6943 0.0531910 .								
R:T:G	8.778	12	2.3235 0.0253153 *								
T:G:F	10.740	44	0.7753 0.7906401								
Residuals	11.333	36									

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

7.11 Example 8.1

(87) MODEL

```

ex8.1 = read.table("C:/G/Rt/Split/asbed.txt", header=TRUE)
ex8.1 = af(ex8.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1)

```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL       104 3951.8 37.999
RESIDUALS    0     0.0
CORRECTED TOTAL 104 3951.8

$Fitness
  R-square Coef Var Root MSE  Y Mean
        1       NA      NA 10.0381

$`Type I`
```

```

Df Sum Sq Mean Sq F value Pr(>F)
R     2 1787.68 893.84
A    12  601.24 50.10
R:A    6   24.93  4.16
B     8  156.87 19.61
R:B    4  319.87 79.97
A:B   60 1012.26 16.87
R:A:B 12   49.00  4.08

```

\$`Type II`

```

Df Sum Sq Mean Sq F value Pr(>F)
R     2 372.22 186.111
A    12  601.24 50.103
R:A    6   50.00  8.333
B     8  156.87 19.609
R:B    4   87.44 21.861
A:B   60 1012.26 16.871
R:A:B 12   49.00  4.083

```

\$`Type III`

```

Df Sum Sq Mean Sq F value Pr(>F)
R     2 372.22 186.111
A    12  572.31 47.692
R:A    6   50.00  8.333
B     8  185.85 23.231
R:B    4   87.44 21.861
A:B   60 1012.26 16.871
R:A:B 12   49.00  4.083

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1), type="III",
singular.ok=TRUE) # NOT WORKING

```

7.12 Example 9.1

(88) MODEL

```

ex9.1 = read.table("C:/G/Rt/Split/Ex9.1-spex1.txt", header=TRUE)
ex9.1 = af(ex9.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex9.1)

```

```

$ANOVA
Response : Y
Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      27 4920.8 182.251 10.594 5.927e-10 ***
RESIDUALS  34  584.9  17.203

```

```

CORRECTED TOTAL 61 5505.6
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   Y Mean
  0.8937663 6.265396 4.147591 66.19839

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
R     3 218.7 72.89 4.2369 0.01199 *
A     3 194.9 64.96 3.7760 0.01930 *
R:A   9 186.9 20.76 1.2070 0.32287
B     3 4087.4 1362.47 79.2018 1.998e-15 ***
A:B   9 233.0 25.88 1.5047 0.18602
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
R     3 157.8 52.61 3.0583 0.04134 *
A     3 227.2 75.73 4.4020 0.01014 *
R:A   9  94.5 10.50 0.6106 0.77932
B     3 4087.4 1362.47 79.2018 1.998e-15 ***
A:B   9 233.0 25.88 1.5047 0.18602
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
R     3 171.0 57.01 3.3138 0.03143 *
A     3 209.7 69.92 4.0643 0.01431 *
R:A   9  94.5 10.50 0.6106 0.77932
B     3 4089.9 1363.29 79.2493 1.998e-15 ***
A:B   9 233.0 25.88 1.5047 0.18602
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.13 Example 9.2

(89) MODEL

```

ex9.2 = read.table("C:/G/Rt/Split/Ex9.2-sbex.txt", header=TRUE)
ex9.2 = af(ex9.2, c("rep", "hyb", "gen"))
GLM(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2)

```

\$ANOVA

```

Response : yield
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        40 247.813 6.1953 4.4606 0.001119 **
RESIDUALS     16 22.222 1.3889
CORRECTED TOTAL 56 270.035
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE yield Mean
0.9177062 2.574747 1.178511   45.77193

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
rep       1  0.239  0.2388 0.1719 0.6839085
hyb       9 66.796  7.4218 5.3437 0.0018370 **
rep:hyb   8 67.000  8.3750 6.0300 0.0011569 **
gen       2 36.351 18.1754 13.0863 0.0004293 ***
rep:gen   2 16.923  8.4616 6.0924 0.0107858 *
hyb:gen  18 60.504  3.3613 2.4201 0.0408545 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
rep       1  0.167  0.1667 0.1200 0.7335481
hyb       9 66.796  7.4218 5.3437 0.0018370 **
rep:hyb   8 67.000  8.3750 6.0300 0.0011569 **
gen       2 36.351 18.1754 13.0863 0.0004293 ***
rep:gen   2 12.111  6.0556 4.3600 0.0308015 *
hyb:gen  18 60.504  3.3613 2.4201 0.0408545 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
rep       1  0.167  0.1667 0.1200 0.7335481
hyb       9 66.796  7.4218 5.3437 0.0018370 **
rep:hyb   8 67.000  8.3750 6.0300 0.0011569 **
gen       2 30.671 15.3356 11.0416 0.0009707 ***
rep:gen   2 12.111  6.0556 4.3600 0.0308015 *
hyb:gen  18 60.504  3.3613 2.4201 0.0408545 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2), type=3),

```

```
singular.ok=TRUE) # NOT OK
```

```
Note: model has aliased coefficients
      sums of squares computed by model comparison
```

Anova Table (Type III tests)

```
Response: yield
          Sum Sq Df F values    Pr(>F)
rep        0.000  0
hyb       66.704  8 6.0033 0.0011847 **
gen       30.671  2 11.0416 0.0009707 ***
rep:hyb   67.000  8 6.0300 0.0011569 **
rep:gen   12.111  2 4.3600 0.0308015 *
hyb:gen   60.504 18 2.4201 0.0408545 *
Residuals 22.222 16
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.14 Example 10.1

(90) MODEL

```
ex10.1 = read.table("C:/G/Rt/Split/Ex10.1-new.txt", header=TRUE)
ex10.1 = af(ex10.1, c("Site", "Block", "A", "B", "C"))
f10.1 = Yield ~ Site/Block + A/Site + B/Site + A:B + A:B:Site + A:B:Site:Block +
         C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site + A:B:C:Site
GLM(f10.1, ex10.1)
```

```
$ANOVA
Response : Yield
          Df     Sum Sq Mean Sq F value    Pr(>F)
MODEL      239 1639561484 6860090    2162 < 2.2e-16 ***
RESIDUALS  240    761522    3173
CORRECTED TOTAL 479 1640323006
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
R-square  Coef Var Root MSE Yield Mean
0.9995357 0.5651396 56.32947 9967.354
```

```
$`Type I`
          Df     Sum Sq Mean Sq F value    Pr(>F)
Site      3      552717   184239 5.8064e+01 < 2e-16 ***
```

Site:Block	8	7062320	882790	2.7822e+02	< 2e-16	***
A	4	1387680917	346920229	1.0933e+05	< 2e-16	***
Site:A	12	34068	2839	8.9470e-01	0.55301	
B	1	100939695	100939695	3.1812e+04	< 2e-16	***
Site:B	3	1618	539	1.6990e-01	0.91662	
A:B	4	31444008	7861002	2.4775e+03	< 2e-16	***
Site:A:B	12	33737	2811	8.8600e-01	0.56185	
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155	
C	3	19356264	6452088	2.0334e+03	< 2e-16	***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16	***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16	***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16	***
Site:C	9	47625	5292	1.6677e+00	0.09747	.
Site:A:C	36	104110	2892	9.1140e-01	0.61768	
Site:B:C	9	61111	6790	2.1400e+00	0.02701	*
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
Site	3	552717	184239	5.8064e+01	< 2e-16	***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16	***
A	4	1387680917	346920229	1.0933e+05	< 2e-16	***
Site:A	12	34068	2839	8.9470e-01	0.55301	
B	1	100939695	100939695	3.1812e+04	< 2e-16	***
Site:B	3	1618	539	1.6990e-01	0.91662	
A:B	4	31444008	7861002	2.4775e+03	< 2e-16	***
Site:A:B	12	33737	2811	8.8600e-01	0.56185	
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155	
C	3	19356264	6452088	2.0334e+03	< 2e-16	***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16	***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16	***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16	***
Site:C	9	47625	5292	1.6677e+00	0.09747	.
Site:A:C	36	104110	2892	9.1140e-01	0.61768	
Site:B:C	9	61111	6790	2.1400e+00	0.02701	*
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
Site	3	552717	184239	5.8064e+01	< 2e-16	***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16	***
A	4	1387680917	346920229	1.0933e+05	< 2e-16	***
Site:A	12	34068	2839	8.9470e-01	0.55301	
B	1	100939695	100939695	3.1812e+04	< 2e-16	***

Site:B	3	1618	539	1.6990e-01	0.91662						
A:B	4	31444008	7861002	2.4775e+03	< 2e-16 ***						
Site:A:B	12	33737	2811	8.8600e-01	0.56185						
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155						
C	3	19356264	6452088	2.0334e+03	< 2e-16 ***						
A:C	12	26075792	2172983	6.8483e+02	< 2e-16 ***						
B:C	3	23901388	7967129	2.5109e+03	< 2e-16 ***						
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16 ***						
Site:C	9	47625	5292	1.6677e+00	0.09747 .						
Site:A:C	36	104110	2892	9.1140e-01	0.61768						
Site:B:C	9	61111	6790	2.1400e+00	0.02701 *						
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(f10.1, ex10.1), type=3, singular.ok=TRUE) # NOT OK for Site:Block
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Yield

	Sum Sq	Df	F values	Pr(>F)
Site	552717	3	5.8064e+01	< 2e-16 ***
A	1387680917	4	1.0933e+05	< 2e-16 ***
B	100939695	1	3.1812e+04	< 2e-16 ***
C	19356264	3	2.0334e+03	< 2e-16 ***
Site:Block	0	0		
Site:A	34068	12	8.9470e-01	0.55301
Site:B	1618	3	1.6990e-01	0.91662
A:B	31444008	4	2.4775e+03	< 2e-16 ***
A:C	26075792	12	6.8483e+02	< 2e-16 ***
B:C	23901388	3	2.5109e+03	< 2e-16 ***
Site:C	47625	9	1.6677e+00	0.09747 .
Site:A:B	33737	12	8.8600e-01	0.56185
A:B:C	41996729	12	1.1030e+03	< 2e-16 ***
Site:A:C	104110	36	9.1140e-01	0.61768
Site:B:C	61111	9	2.1400e+00	0.02701 *
Site:Block:A:B	186911	72	8.1810e-01	0.84155
Site:A:B:C	82475	36	7.2200e-01	0.87941
Residuals	761522	240		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

7.15 Example 10.2

(91) MODEL

```
ex10.2 = read.table("C:/G/Rt/Split/Ex10.2-spbsite.txt", header=TRUE)
ex10.2 = af(ex10.2, c("Site", "Block", "A", "B"))
GLM(Yield ~ Site + Site:Block + A + A:Site + A:Site:Block + B + B:Site +
     B:Site:Block + A:B + A:B:Site, ex10.2)
```

\$ANOVA

Response : Yield

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	227	6370995084	28066058	10814	< 2.2e-16 ***
RESIDUALS	252	654049	2595		
CORRECTED TOTAL	479	6371649132			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	Yield	Mean
0.9998974	0.4596687	50.94537		11083.06		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	2	523573968	261786984	1.0086e+05	< 2.2e-16 ***
Site:Block	9	3756646710	417405190	1.6082e+05	< 2.2e-16 ***
A	4	29288163	7322041	2.8211e+03	< 2.2e-16 ***
Site:A	8	247899	30987	1.1939e+01	1.998e-14 ***
Site:Block:A	36	1783391	49539	1.9087e+01	< 2.2e-16 ***
B	7	1937592291	276798899	1.0665e+05	< 2.2e-16 ***
Site:B	14	15903698	1135978	4.3768e+02	< 2.2e-16 ***
Site:Block:B	63	105727288	1678211	6.4660e+02	< 2.2e-16 ***
A:B	28	91141	3255	1.2541e+00	0.1838
Site:A:B	56	140534	2510	9.6690e-01	0.5461

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	2	523573968	261786984	1.0086e+05	< 2.2e-16 ***
Site:Block	9	3756646710	417405190	1.6082e+05	< 2.2e-16 ***
A	4	29288163	7322041	2.8211e+03	< 2.2e-16 ***
Site:A	8	247899	30987	1.1939e+01	1.998e-14 ***
Site:Block:A	36	1783391	49539	1.9087e+01	< 2.2e-16 ***
B	7	1937592291	276798899	1.0665e+05	< 2.2e-16 ***
Site:B	14	15903698	1135978	4.3768e+02	< 2.2e-16 ***
Site:Block:B	63	105727288	1678211	6.4660e+02	< 2.2e-16 ***

```

A:B          28      91141      3255 1.2541e+00      0.1838
Site:A:B     56      140534      2510 9.6690e-01      0.5461
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df   Sum Sq  Mean Sq   F value   Pr(>F)
Site       2 523573968 261786984 1.0086e+05 < 2.2e-16 ***
Site:Block 9 3756646710 417405190 1.6082e+05 < 2.2e-16 ***
A          4 29288163  7322041 2.8211e+03 < 2.2e-16 ***
Site:A      8 247899    30987 1.1939e+01 1.998e-14 ***
Site:Block:A 36 1783391    49539 1.9087e+01 < 2.2e-16 ***
B          7 1937592291 276798899 1.0665e+05 < 2.2e-16 ***
Site:B      14 15903698   1135978 4.3768e+02 < 2.2e-16 ***
Site:Block:B 63 105727288   1678211 6.4660e+02 < 2.2e-16 ***
A:B         28      91141      3255 1.2541e+00      0.1838
Site:A:B     56      140534      2510 9.6690e-01      0.5461
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.16 Example 11.1

(92) MODEL

```

ex11.1 = read.table("C:/G/Rt/Split/Ex11.1-cov.txt", header=TRUE)
ex11.1 = af(ex11.1, c("R", "T", "S"))
GLM(Y ~ R + T + R:T + S + S:T, ex11.1)

```

```

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value   Pr(>F)
MODEL      11      328 29.8182 3.1948 0.02875 *
RESIDUALS  12      112  9.3333
CORRECTED TOTAL 23      440
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Y Mean
0.7454545 43.64358 3.05505      7

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value   Pr(>F)
R      2      48      24  2.5714 0.11765
T      1      24      24  2.5714 0.13479
R:T    2      16      8  0.8571 0.44880

```

```

S     3      156      52  5.5714 0.01251 *
T:S   3       84      28  3.0000 0.07277 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value Pr(>F)  

R     2     48     24  2.5714 0.11765  

T     1     24     24  2.5714 0.13479  

R:T   2     16      8  0.8571 0.44880  

S     3     156      52  5.5714 0.01251 *  

T:S   3     84     28  3.0000 0.07277 .  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

R     2     48     24  2.5714 0.11765  

T     1     24     24  2.5714 0.13479  

R:T   2     16      8  0.8571 0.44880  

S     3     156      52  5.5714 0.01251 *  

T:S   3     84     28  3.0000 0.07277 .  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(93) MODEL

```
GLM(Z ~ R + T + R:T + S + S:T, ex11.1)
```

```
$ANOVA  

Response : Z  

  Df Sum Sq Mean Sq F value Pr(>F)  

MODEL          11     46  4.1818  2.5091 0.06452 .  

RESIDUALS      12     20  1.6667  

CORRECTED TOTAL 23     66  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE Z Mean  

  0.6969697 51.63978 1.290994    2.5
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

R     2      9     4.5     2.7 0.1076  

T     1      6     6.0     3.6 0.0821 .  

R:T   2      1     0.5     0.3 0.7462
```

```

S      3      9      3.0      1.8 0.2008
T:S    3     21      7.0      4.2 0.0301 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df Sum Sq Mean Sq F value Pr(>F)
R      2      9      4.5      2.7 0.1076
T      1      6      6.0      3.6 0.0821 .
R:T    2      1      0.5      0.3 0.7462
S      3      9      3.0      1.8 0.2008
T:S    3     21      7.0      4.2 0.0301 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df Sum Sq Mean Sq F value Pr(>F)
R      2      9      4.5      2.7 0.1076
T      1      6      6.0      3.6 0.0821 .
R:T    2      1      0.5      0.3 0.7462
S      3      9      3.0      1.8 0.2008
T:S    3     21      7.0      4.2 0.0301 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(94) MODEL

```
GLM(Y ~ R + T + R:T + S + S:T + Z, ex11.1)
```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL       12 342.45 28.5375   3.218 0.03116 *
RESIDUALS    11  97.55  8.8682
CORRECTED TOTAL 23 440.00
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE Y Mean
0.7782955 42.54213 2.977949      7

```

```

$`Type I` 
  Df Sum Sq Mean Sq F value Pr(>F)
R      2  48.00  24.00  2.7063 0.11071
T      1  24.00  24.00  2.7063 0.12820
R:T    2  16.00   8.00  0.9021 0.43373

```

```

S      3 156.00   52.00  5.8637 0.01211 *
T:S    3  84.00   28.00  3.1574 0.06828 .
Z      1  14.45   14.45  1.6294 0.22807
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
R      2 18.300  9.1500  1.0318 0.38844
T      1  2.679  2.6786  0.3020 0.59359
R:T    2  9.450  4.7250  0.5328 0.60137
S      3 79.196 26.3985  2.9768 0.07822 .
T:S    3 37.474 12.4915  1.4086 0.29234
Z      1 14.450 14.4500  1.6294 0.22807
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
R      2 20.209 10.1043  1.1394 0.35505
T      1  6.104  6.1038  0.6883 0.42439
R:T    2  9.450  4.7250  0.5328 0.60137
S      3 84.243 28.0810  3.1665 0.06782 .
T:S    3 37.474 12.4915  1.4086 0.29234
Z      1 14.450 14.4500  1.6294 0.22807
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.17 Example 11.2

(95) MODEL

```

ex11.2a = read.table("C:/G/Rt/Split/Ex11.2-sp3.txt", header=TRUE)
ex11.2a = af(ex11.2a, "A")
ex11.2a$MY = (ex11.2a$Y1 + ex11.2a$Y2)/sqrt(2)
ex11.2a$Z = 2*ex11.2a$Z/sqrt(2)
GLM(MY ~ Z + A, ex11.2a)

```

```

$ANOVA
Response : MY
  Df  Sum Sq Mean Sq F value Pr(>F)
MODEL        2 234.639 117.32  9.5696 0.01953 *
RESIDUALS     5  61.298   12.26
CORRECTED TOTAL 7 295.937
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE  MY Mean
  0.7928678 17.45091 3.501377 20.06415

$`Type I`
  Df  Sum Sq Mean Sq F value  Pr(>F)
Z  1 190.148 190.148 15.5101 0.01098 *
A  1  44.492  44.492  3.6291 0.11512
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq Mean Sq F value  Pr(>F)
Z  1 166.577 166.577 13.5874 0.0142 *
A  1  44.492  44.492  3.6291 0.1151
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df  Sum Sq Mean Sq F value  Pr(>F)
Z  1 166.577 166.577 13.5874 0.0142 *
A  1  44.492  44.492  3.6291 0.1151
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(96) MODEL

```

ex11.2b = read.table("C:/G/Rt/Split/Ex11.2-two.txt", header=TRUE)
ex11.2b = af(ex11.2b, c("sub", "A", "B"))
GLM(Y ~ A + A:sub + B + A:B, ex11.2b)

```

```

$ANOVA
Response : Y
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      9 382.06  42.451  39.954 0.0001135 ***
RESIDUALS   6   6.38   1.062
CORRECTED TOTAL 15 388.44
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE  Y Mean
  0.9835881 7.265384 1.030776 14.1875

$`Type I`
  Df  Sum Sq Mean Sq F value    Pr(>F)

```

```

A      1 68.062 68.062 64.0588 0.0002029 ***
A:sub  6 227.875 37.979 35.7451 0.0001934 ***
B      1 85.562 85.562 80.5294 0.0001070 ***
A:B    1  0.562   0.562  0.5294  0.4942562
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

Df  Sum Sq Mean Sq F value    Pr(>F)  

A      1 68.062 68.062 64.0588 0.0002029 ***  

A:sub  6 227.875 37.979 35.7451 0.0001934 ***  

B      1 85.562 85.562 80.5294 0.0001070 ***  

A:B    1  0.562   0.562  0.5294  0.4942562
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

Df  Sum Sq Mean Sq F value    Pr(>F)  

A      1 68.062 68.062 64.0588 0.0002029 ***  

A:sub  6 227.875 37.979 35.7451 0.0001934 ***  

B      1 85.562 85.562 80.5294 0.0001070 ***  

A:B    1  0.562   0.562  0.5294  0.4942562
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(97) MODEL

```

ex11.2c = read.table("C:/G/Rt/Split/Ex11.2-spcov2.txt", header=TRUE)
ex11.2c = af(ex11.2c, c("block", "whole", "split"))
GLM(Y ~ block + whole + block:whole + split:white, ex11.2c)

```

```

$ANOVA
Response : Y
Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        11     328 29.8182  3.1948 0.02875 *
RESIDUALS    12     112  9.3333
CORRECTED TOTAL 23     440
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE Y Mean
0.7454545 43.64358 3.05505      7

```

```

$`Type I`  

Df  Sum Sq Mean Sq F value    Pr(>F)
block        2      48       24  2.5714 0.11765

```

```

whole      1     24      24  2.5714 0.13479
block:whole 2     16      8   0.8571 0.44880
split      3    156      52  5.5714 0.01251 *
whole:split 3     84      28  3.0000 0.07277 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block      2     48      24  2.5714 0.11765  

whole      1     24      24  2.5714 0.13479  

block:whole 2     16      8   0.8571 0.44880  

split      3    156      52  5.5714 0.01251 *
whole:split 3     84      28  3.0000 0.07277 .

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block      2     48      24  2.5714 0.11765  

whole      1     24      24  2.5714 0.13479  

block:whole 2     16      8   0.8571 0.44880  

split      3    156      52  5.5714 0.01251 *
whole:split 3     84      28  3.0000 0.07277 .

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(98) MODEL

```
GLM(Z ~ block + whole + block:whole + split + split:white, ex11.2c)
```

```

$ANOVA  

Response : Z  

      Df Sum Sq Mean Sq      F value      Pr(>F)  

MODEL       11     38  3.4545 3.5903e+15 < 2.2e-16 ***  

RESIDUALS    12      0  0.0000  

CORRECTED TOTAL 23     38  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

R-square   Coef Var      Root MSE Z Mean  

1 8.86264e-07 3.101924e-08      3.5

```

```

$`Type I`  

      Df Sum Sq Mean Sq      F value      Pr(>F)  

block       2 36.000 18.0000 1.8707e+16 <2e-16 ***

```

```

whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***
block:whole 2  1.333  0.6667 6.9286e+14 <2e-16 ***
split      3  0.000  0.0000 0.0000e+00      1
whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq   F value Pr(>F)  

block      2 36.000 18.0000 1.8707e+16 <2e-16 ***  

whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***  

block:whole 2  1.333  0.6667 6.9286e+14 <2e-16 ***  

split      3  0.000  0.0000 0.0000e+00      1  

whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq   F value Pr(>F)  

block      2 36.000 18.0000 1.8707e+16 <2e-16 ***  

whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***  

block:whole 2  1.333  0.6667 6.9286e+14 <2e-16 ***  

split      3  0.000  0.0000 0.0000e+00      1  

whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(99) MODEL

```
GLM(Y ~ block + whole + block:white + split:white + Z, ex11.2c)
```

```
$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      11     328 29.8182  3.1948 0.02875 *  

RESIDUALS    12     112  9.3333  

CORRECTED TOTAL 23     440
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE Y Mean  

0.7454545 43.64358 3.05505      7
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block      2     48       24  2.5714 0.11765
```

```

whole      1     24      24  2.5714 0.13479
block:whole 2     16       8  0.8571 0.44880
split      3    156      52  5.5714 0.01251 *
whole:split 3     84      28  3.0000 0.07277 .
Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

block      2  13.286   6.643  0.7117 0.51039  

whole      1  16.000  16.000  1.7143 0.21495  

block:whole 1  16.000  16.000  1.7143 0.21495  

split      3 156.000   52.000  5.5714 0.01251 *  

whole:split 3  84.000   28.000  3.0000 0.07277 .  

Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value Pr(>F)  

block      2  13.286   6.643  0.7117 0.51039  

whole      1  16.000  16.000  1.7143 0.21495  

block:whole 1  16.000  16.000  1.7143 0.21495  

split      3 156.000   52.000  5.5714 0.01251 *  

whole:split 3  84.000   28.000  3.0000 0.07277 .  

Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.18 Example 11.3

(100) MODEL

```

ex11.3 = read.table("C:/G/Rt/Split/Ex11.3-sbcov.txt", header=TRUE)
ex11.3 = af(ex11.3, c("block", "A", "B"))
GLM(Y ~ block + A + block:A + B + block:B + A:B, ex11.3)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      17 16.833   0.9902  1.9804 0.2038
RESIDUALS      6  3.000   0.5000
CORRECTED TOTAL 23 19.833

```

```

$Fitness
  R-square Coef Var Root MSE   Y Mean
  0.8487395 24.24366 0.7071068 2.916667

$`Type I` 
      Df Sum Sq Mean Sq F value Pr(>F)
block    3 4.5000 1.5000 3.0000 0.11696
A        1 1.5000 1.5000 3.0000 0.13397
block:A  3 0.5000 0.1667 0.3333 0.80220
B        2 8.3333 4.1667 8.3333 0.01855 *
block:B  6 1.0000 0.1667 0.3333 0.89648
A:B     2 1.0000 0.5000 1.0000 0.42188
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value Pr(>F)
block    3 4.5000 1.5000 3.0000 0.11696
A        1 1.5000 1.5000 3.0000 0.13397
block:A  3 0.5000 0.1667 0.3333 0.80220
B        2 8.3333 4.1667 8.3333 0.01855 *
block:B  6 1.0000 0.1667 0.3333 0.89648
A:B     2 1.0000 0.5000 1.0000 0.42188
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value Pr(>F)
block    3 4.5000 1.5000 3.0000 0.11696
A        1 1.5000 1.5000 3.0000 0.13397
block:A  3 0.5000 0.1667 0.3333 0.80220
B        2 8.3333 4.1667 8.3333 0.01855 *
block:B  6 1.0000 0.1667 0.3333 0.89648
A:B     2 1.0000 0.5000 1.0000 0.42188
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(101) MODEL

```
GLM(Z ~ block + A + block:A + B + block:B + A:B, ex11.3)
```

```

$ANOVA
Response : Z
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       17 31.167 1.83333    3.3 0.07324 .
RESIDUALS   6  3.333 0.55556
CORRECTED TOTAL 23 34.500

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE Z Mean
  0.9033816 42.59177 0.745356   1.75

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
block   3 6.8333 2.2778     4.1 0.06689 .
A       1 6.0000 6.0000    10.8 0.01669 *
block:A 3 1.6667 0.5556     1.0 0.45472
B       2 13.0000 6.5000    11.7 0.00850 **
block:B 6 3.6667 0.6111     1.1 0.45542
A:B     2 0.0000 0.0000     0.0 1.00000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
block   3 6.8333 2.2778     4.1 0.06689 .
A       1 6.0000 6.0000    10.8 0.01669 *
block:A 3 1.6667 0.5556     1.0 0.45472
B       2 13.0000 6.5000    11.7 0.00850 **
block:B 6 3.6667 0.6111     1.1 0.45542
A:B     2 0.0000 0.0000     0.0 1.00000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
block   3 6.8333 2.2778     4.1 0.06689 .
A       1 6.0000 6.0000    10.8 0.01669 *
block:A 3 1.6667 0.5556     1.0 0.45472
B       2 13.0000 6.5000    11.7 0.00850 **
block:B 6 3.6667 0.6111     1.1 0.45542
A:B     2 0.0000 0.0000     0.0 1.00000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(102) MODEL

```
GLM(Y ~ block + A + block:A + B + block:B + A:B + Z, ex11.3)
```

```
$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
```

MODEL	18	17.8417	0.99120	2.4884	0.1589
RESIDUALS	5	1.9917	0.39833		
CORRECTED TOTAL	23	19.8333			

\$Fitness

R-square	Coef	Var	Root	MSE	Y Mean
0.8995798	21.63897	0.6311365	2.916667		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	3	4.5000	1.5000	3.7657	0.09378 .
A	1	1.5000	1.5000	3.7657	0.10999
block:A	3	0.5000	0.1667	0.4184	0.74788
B	2	8.3333	4.1667	10.4603	0.01634 *
block:B	6	1.0000	0.1667	0.4184	0.84059
A:B	2	1.0000	0.5000	1.2552	0.36163
Z	1	1.0083	1.0083	2.5314	0.17248

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	3	3.6203	1.20678	3.0296	0.1319
A	1	0.0000	0.00000	0.0000	1.0000
block:A	3	0.2583	0.08611	0.2162	0.8813
B	2	1.0317	0.51587	1.2951	0.3522
block:B	6	0.4210	0.07017	0.1762	0.9717
A:B	2	1.0000	0.50000	1.2552	0.3616
Z	1	1.0083	1.00833	2.5314	0.1725

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	3	3.6613	1.22045	3.0639	0.1297
A	1	0.0054	0.00536	0.0134	0.9122
block:A	3	0.2583	0.08611	0.2162	0.8813
B	2	0.7685	0.38427	0.9647	0.4423
block:B	6	0.4210	0.07017	0.1762	0.9717
A:B	2	1.0000	0.50000	1.2552	0.3616
Z	1	1.0083	1.00833	2.5314	0.1725

8 Hinkelmann & Kempthorne - Volume 1

Reference

- Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.

8.1 Chapter 6

8.1.1 p202

(103) MODEL

```
v1p202 = read.table("C:/G/Rt/Kemp/v1p202.txt", head=TRUE)
v1p202 = af(v1p202,c("brand"))
GLM(miles ~ brand, v1p202) # OK

$ANOVA
Response : miles
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL     4 47.234  11.809  15.661 0.004924 ***
RESIDUALS   5  3.770    0.754
CORRECTED TOTAL  9 51.004
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE miles Mean
0.9260842 3.309191 0.8683317      26.24

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand    4 47.234  11.809  15.661 0.004924 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand    4 47.234  11.809  15.661 0.004924 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand    4 47.234  11.809  15.661 0.004924 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.1.2 p205

(104) MODEL

```
v1p205 = read.table("C:/G/Rt/Kemp/v1p205.txt", head=TRUE)
v1p205 = af(v1p205,c("brand", "car"))
GLM(miles ~ brand + car %in% brand, v1p205) # OK

$ANOVA
Response : miles
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      9 140.05 15.561   80.21 1.017e-13 ***
RESIDUALS  20   3.88   0.194
CORRECTED TOTAL 29 143.93
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE miles Mean
0.9730418 1.683265 0.4404543   26.16667

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361    7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361    7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361    7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.2 Chapter 7

8.2.1 p232

(105) MODEL

```

v1p232 = read.table("C:/G/Rt/Kemp/v1p232.txt", head=TRUE)
v1p232 = af(v1p232,c("trt"))
GLM(yield ~ trt, v1p232) # OK

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       4 59.174 14.793 28.781 0.0012 **
RESIDUALS   5  2.570  0.514
CORRECTED TOTAL 9 61.744
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE yield Mean
0.9583765 4.497729 0.7169379      15.94

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.2.2 p235

(106) MODEL

```

v1p235 = read.table("C:/G/Rt/Kemp/v1p235.txt", head=TRUE)
v1p235 = af(v1p235,c("density"))
GLM(yield ~ density, v1p235) # OK

```

```

$ANOVA
Response : yield

```

```

          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        4 88.007 22.0017 32.198 1.095e-05 ***
RESIDUALS    10  6.833  0.6833
CORRECTED TOTAL 14 94.840
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE yield Mean
0.9279488 5.040486 0.8266398      16.4

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
density     4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
density     4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
density     4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.3 Chapter 8

8.3.1 p265

(107) MODEL

```
v1p265 = read.table("C:/G/Rt/Kemp/v1p265.txt", head=TRUE)
v1p265 = af(v1p265,c("trt"))
GLM(y ~ trt + x, v1p265) # OK
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        3 84.678 28.2260 36.866 4.941e-06 ***
RESIDUALS    11  8.422  0.7656
CORRECTED TOTAL 14 93.100
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE y Mean
  0.9095378 9.722312 0.8750081      9

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  2 66.868 33.434 43.668 5.858e-06 ***
x    1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  2 83.147 41.573 54.299 1.996e-06 ***
x    1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  2 83.147 41.573 54.299 1.996e-06 ***
x    1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.3.2 p272

(108) MODEL

```
GLM(y ~ trt + x %in% trt, v1p265) # OK
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 85.711 17.142 20.881 0.0001046 ***
RESIDUALS   9  7.389  0.821
CORRECTED TOTAL 14 93.100
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE y Mean
  0.9206374 10.06744 0.9060697      9

```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

trt     2 66.868 33.434 40.7254 3.092e-05 ***  

trt:x  3 18.843  6.281  7.6509  0.007578 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

trt     2 66.868 33.434 40.7254 3.092e-05 ***  

trt:x  3 18.843  6.281  7.6509  0.007578 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

  Df   Sum Sq Mean Sq F value    Pr(>F)  

trt     2  6.1392  3.0696  3.7390  0.065769 .  

trt:x  3 18.8433  6.2811  7.6509  0.007578 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.3.3 p273

(109) MODEL

```
GLM(y ~ trt + x + x %in% trt, v1p265) # OK
```

```
$ANOVA  

Response : y  

  Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      5 85.711 17.142 20.881 0.0001046 ***  

RESIDUALS  9  7.389  0.821  

CORRECTED TOTAL 14 93.100  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE y Mean  

  0.9206374 10.06744 0.9060697      9
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

trt     2 66.868 33.434 40.7254 3.092e-05 ***  

x       1 17.810 17.810 21.6940  0.001189 **  

trt:x  2  1.033  0.517  0.6294  0.554843  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

trt     2 83.147 41.573 50.6397 1.267e-05 ***  

x       1 17.810 17.810 21.6940  0.001189 **  

trt:x  2  1.033   0.517  0.6294  0.554843  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

trt     2 6.1392 3.0696 3.7390 0.065769 .  

x       1 17.2071 17.2071 20.9597 0.001331 **  

trt:x  2  1.0334 0.5167  0.6294  0.554843  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4 Chapter 9

8.4.1 p344

(110) MODEL

```

v1p344 = read.table("C:/G/Rt/Kemp/v1p344.txt", head=TRUE)
v1p344 = af(v1p344,c("diet", "litter"))
GLM(gain ~ litter + diet, v1p344)

```

```

$ANOVA
Response : gain
      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      9 4915.6 546.18 15.544 3.363e-07 ***  

RESIDUALS  20 702.8  35.14  

CORRECTED TOTAL 29 5618.4  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE gain Mean
0.874919 8.677219 5.927698 68.31333

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

litter    5 4438.0   887.6 25.2608 5.298e-08 ***  

diet      4  477.6   119.4  3.3981  0.02824 *  

---  


```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

litter   5 4438.0   887.6 25.2608 5.298e-08 ***  

diet     4  477.6   119.4  3.3981  0.02824 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

litter   5 4438.0   887.6 25.2608 5.298e-08 ***  

diet     4  477.6   119.4  3.3981  0.02824 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4.2 p349

(111) MODEL

```

v1p349 = read.table("C:/G/Rt/Kemp/v1p349.txt", head=TRUE)
v1p349 = af(v1p349,c("subject", "exercise"))
GLM(diast ~ subject + exercise + subject:exercise, v1p349) # OK

```

```

$ANOVA  

Response : diast  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL       14 1541.5 110.105 28.475 2.953e-08 ***  

RESIDUALS    15   58.0   3.867  

CORRECTED TOTAL 29 1599.5  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE diast Mean  

  0.9637379 1.461633 1.966384   134.5333

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

subject      4 905.13 226.283 58.5216 5.672e-09 ***  

exercise     2 591.27 295.633 76.4569 1.357e-08 ***  

subject:exercise 8  45.07   5.633  1.4569     0.2522  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type II`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
subject        4 905.13 226.283 58.5216 5.672e-09 ***
exercise       2 591.27 295.633 76.4569 1.357e-08 ***
subject:exercise 8  45.07   5.633  1.4569     0.2522
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
          Df Sum Sq Mean Sq F value    Pr(>F)
subject        4 905.13 226.283 58.5216 5.672e-09 ***
exercise       2 591.27 295.633 76.4569 1.357e-08 ***
subject:exercise 8  45.07   5.633  1.4569     0.2522
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4.3 p354

(112) MODEL

```

v1p354 = read.table("C:/G/Rt/Kemp/v1p354.txt", head=TRUE)
v1p354 = af(v1p354,c("loc", "block", "HSF"))
GLM(height ~ loc + block %in% loc + HSF + loc:HSF + block:loc:HSF, v1p354) # OK

```

\$ANOVA

Response : height

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	23	40782	1773.12	80.444 < 2.2e-16	***
RESIDUALS	24	529	22.04		
CORRECTED TOTAL	47	41311			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	height	Mean
0.9871946	2.228571	4.694855		210.6667		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
loc	1	20336.3	20336.3	922.6314 < 2.2e-16	***
loc:block	6	1462.3	243.7	11.0573 6.408e-06	***
HSF	2	12170.7	6085.3	276.0832 < 2.2e-16	***
loc:HSF	2	6511.2	3255.6	147.7013 3.242e-14	***
loc:block:HSF	12	301.2	25.1	1.1386 0.3769	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

loc        1 20336.3 20336.3 922.6314 < 2.2e-16 ***  

loc:block   6  1462.3   243.7 11.0573 6.408e-06 ***  

HSF        2 12170.7  6085.3 276.0832 < 2.2e-16 ***  

loc:HSF     2  6511.2  3255.6 147.7013 3.242e-14 ***  

loc:block:HSF 12   301.2    25.1   1.1386    0.3769  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

loc        1 20336.3 20336.3 922.6314 < 2.2e-16 ***  

loc:block   6  1462.3   243.7 11.0573 6.408e-06 ***  

HSF        2 12170.7  6085.3 276.0832 < 2.2e-16 ***  

loc:HSF     2  6511.2  3255.6 147.7013 3.242e-14 ***  

loc:block:HSF 12   301.2    25.1   1.1386    0.3769  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.4.4 p357

(113) MODEL

```
v1p357 = read.table("C:/G/Rt/Kemp/v1p357.txt", head=TRUE)
v1p357 = af(v1p357,c("var", "N"))
GLM(y ~ var + N + var:N, v1p357) # OK
```

```
$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       9 4465.5  496.16 14.116 0.000142 ***
RESIDUALS   10  351.5   35.15
CORRECTED TOTAL 19 4817.0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
R-square Coef Var Root MSE y Mean
0.9270285 4.310246 5.928744 137.55
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)
var     1  140.5  140.45  3.9957  0.073519 .
N      4 3393.7  848.42 24.1373 4.027e-05 ***
var:N  4  931.3  232.82  6.6238  0.007152 **
```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
var     1 140.5 140.45  3.9957  0.073519 .
N       4 3393.7 848.43 24.1373 4.027e-05 ***
var:N   4  931.3 232.82  6.6238  0.007152 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
var     1 140.5 140.45  3.9957  0.073519 .
N       4 3393.7 848.42 24.1373 4.027e-05 ***
var:N   4  931.3 232.83  6.6238  0.007152 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4.5 p361

(114) MODEL

```

v1p361 = read.table("C:/G/Rt/Kemp/v1p361.txt", head=TRUE)
v1p361 = af(v1p361,c("block", "trt"))
GLM(y ~ block + trt, v1p361) # OK

```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      4 241.33 60.333 40.222 0.1176
RESIDUALS  1   1.50   1.500
CORRECTED TOTAL 5 242.83

```

```

$Fitness
  R-square Coef Var Root MSE   y Mean
  0.9938229 6.175184 1.224745 19.83333

```

```

$`Type I`
  Df Sum Sq Mean Sq F value  Pr(>F)
block   2 24.333 12.167  8.1111 0.24097
trt     2 217.000 108.500 72.3333 0.08286 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
```

```

Df Sum Sq Mean Sq F value Pr(>F)
block  2     108     54.0  36.000 0.11704
trt    2     217    108.5  72.333 0.08286 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
Df Sum Sq Mean Sq F value Pr(>F)
block  2     108     54.0  36.000 0.11704
trt    2     217    108.5  72.333 0.08286 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

y = model.frame(y ~ block + trt, v1p361)[,1]
x = ModelMatrix(y ~ block + trt, v1p361)
rx = lfit(x, y)
K = cbind(rep(1, 3), matrix(1/3, nrow=3, ncol=3), diag(3)) ; K

```

	[,1]	[,2]	[,3]	[,4]	[,5]	[,6]	[,7]
[1,]	1	0.3333333	0.3333333	0.3333333	1	0	0
[2,]	1	0.3333333	0.3333333	0.3333333	0	1	0
[3,]	1	0.3333333	0.3333333	0.3333333	0	0	1

est(K, x\$X, rx)

	Estimate	Lower CL	Upper CL	Std. Error	t value	Df	Pr(> t)
[1,]	29.5	17.334735	41.66526	0.9574271	30.81175	1	0.02065434
[2,]	16.5	4.334735	28.66526	0.9574271	17.23369	1	0.03689905
[3,]	13.5	1.334735	25.66526	0.9574271	14.10029	1	0.04507394

attr(,"Estimability")
[1] TRUE TRUE TRUE

8.5 Chapter 10

8.5.1 p405

(115) MODEL

```

v1p405 = read.table("C:/G/Rt/Kemp/v1p405.txt", head=TRUE)
v1p405 = af(v1p405,c("trt", "Row", "Col"))
GLM(y ~ Row + Col + trt, v1p405) # OK

```

```

$ANOVA
Response : y
Df Sum Sq Mean Sq F value Pr(>F)

```

```

MODEL           12 4094.7  341.23  2.3416 0.07739 .
RESIDUALS      12 1748.7  145.73
CORRECTED TOTAL 24 5843.4
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE y Mean
0.7007379 12.93584 12.07173 93.32

$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
Row  4 514.24 128.56  0.8822 0.50328
Col  4 1711.44 427.86  2.9360 0.06611 .
trt  4 1869.04 467.26  3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
Row  4 514.24 128.56  0.8822 0.50328
Col  4 1711.44 427.86  2.9360 0.06611 .
trt  4 1869.04 467.26  3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
Row  4 514.24 128.56  0.8822 0.50328
Col  4 1711.44 427.86  2.9360 0.06611 .
trt  4 1869.04 467.26  3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.5.2 p408

(116) MODEL

```

v1p408 = read.table("C:/G/Rt/Kemp/v1p408.txt", head=TRUE)
v1p408 = af(v1p408,c("breed", "farm", "wclass", "dosage"))
GLM(response ~ breed + breed:farm + wclass + dosage + breed:dosage, v1p408) # OK

```

```

$ANOVA
Response : response
Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       16 4470.2 279.391 140.87 2.039e-13 ***

```

```

RESIDUALS      15   29.7   1.983
CORRECTED TOTAL 31 4500.0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE response Mean
  0.9933889 0.904211 1.408309           155.75

$`Type I`
    Df Sum Sq Mean Sq   F value   Pr(>F)
breed       1 3280.5 3280.5 1654.0336 < 2.2e-16 ***
breed:farm  6    9.0    1.5    0.7563   0.6146
wclass      3  466.8  155.6   78.4454 2.142e-09 ***
dosage      3  580.2  193.4   97.5210 4.596e-10 ***
breed:dosage 3  133.8   44.6   22.4790 8.366e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
    Df Sum Sq Mean Sq   F value   Pr(>F)
breed       1 3280.5 3280.5 1654.0336 < 2.2e-16 ***
breed:farm  6    9.0    1.5    0.7563   0.6146
wclass      3  466.7  155.6   78.4454 2.142e-09 ***
dosage      3  580.2  193.4   97.5210 4.596e-10 ***
breed:dosage 3  133.8   44.6   22.4790 8.366e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
    Df Sum Sq Mean Sq   F value   Pr(>F)
breed       1 3280.5 3280.5 1654.0336 < 2.2e-16 ***
breed:farm  6    9.0    1.5    0.7563   0.6146
wclass      3  466.8  155.6   78.4454 2.142e-09 ***
dosage      3  580.3  193.4   97.5210 4.596e-10 ***
breed:dosage 3  133.7   44.6   22.4790 8.366e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.5.3 p410

(117) MODEL

```

v1p410 = read.table("C:/G/Rt/Kemp/v1p410.txt", head=TRUE)
v1p410$carry = ifelse(v1p410$carry == 0, 3, v1p410$carry)
v1p410 = af(v1p410,c("period", "sequence", "steer", "trt", "carry"))
GLM(y ~ period + sequence + steer:sequence + trt + carry, v1p410) # OK

```

```

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       17 1302.51  76.618  8.7402 1.572e-05 ***
RESIDUALS    18 157.79   8.766
CORRECTED TOTAL 35 1460.31
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   y Mean
  0.8919461 5.654535 2.960778 52.36111

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
period       2 292.06 146.028 16.6580 8.038e-05 ***
sequence     5 326.47  65.294  7.4484 0.0006072 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt          2 549.06 274.528 31.3166 1.377e-06 ***
carry         2 16.43   8.215  0.9372 0.4100385
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
period       2 172.31  86.154  9.8279 0.0013030 **
sequence     5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt          2 440.61 220.304 25.1311 6.164e-06 ***
carry         2 16.43   8.215  0.9372 0.4100385
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
period       2 172.31  86.154  9.8279 0.0013030 **
sequence     5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt          2 440.61 220.304 25.1311 6.164e-06 ***
carry         2 16.43   8.215  0.9372 0.4100385
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(y ~ period + sequence + steer:sequence + trt + carry, v1p410), type=3,
      singular.ok=TRUE) # NOT OK for sequence

```

Note: model has aliased coefficients

```
sums of squares computed by model comparison
```

```
Anova Table (Type III tests)
```

```
Response: y
```

	Sum Sq	Df	F value	Pr(>F)
period	172.31	2	9.8279	0.001303 **
sequence	0.00	0		
trt	440.61	2	25.1311	6.164e-06 ***
carry	16.43	2	0.9372	0.410038
sequence:steer	118.50	6	2.2530	0.084912 .
Residuals	157.79	18		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8.6 Chapter 11

8.6.1 p432

```
(118) MODEL
```

```
v1p432 = read.table("C:/G/Rt/Kemp/v1p432.txt", head=TRUE)
v1p432 = af(v1p432,c("V", "Block", "A", "B", "C"))
GLM(Y ~ V + Block:V + A + B + A:B + V:A + V:B + V:A:B + Block:A:V + Block:B:V,
v1p432) # OK
```

```
$ANOVA
```

```
Response : Y
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	94	261663	2783.65	30.584	2.065e-14 ***
RESIDUALS	25	2275	91.02		
CORRECTED TOTAL	119	263939			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$Fitness
```

R-square	Coef	Var	Root	MSE	Y	Mean
0.991379	1.556578	9.540266			612.9	

```
$`Type I`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	282.2094 < 2.2e-16 ***	
V:Block	25	50019	2001	21.9825 1.588e-11 ***	
A	1	18451	18451	202.7233 1.692e-13 ***	
B	1	78541	78541	862.9280 < 2.2e-16 ***	
A:B	1	108	108	1.1899 0.28575	

```

V:A      4   3751     938  10.3023 4.532e-05 ***
V:B      4   307      77   0.8421   0.51168
V:A:B    4   1495     374   4.1058   0.01081 *
V:Block:A 25  3416     137   1.5011   0.15818
V:Block:B 25  2833     113   1.2451   0.29390
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

V       4 102743  25686 282.2094 < 2.2e-16 ***  

V:Block 25 50019   2001 21.9825 1.588e-11 ***  

A       1 18451   18451 202.7233 1.692e-13 ***  

B       1 78541   78541 862.9280 < 2.2e-16 ***  

A:B     1   108     108   1.1899  0.28575  

V:A     4   3751     938  10.3023 4.532e-05 ***  

V:B     4   307      77   0.8421   0.51168  

V:A:B    4   1495     374   4.1058   0.01081 *  

V:Block:A 25  3416     137   1.5011   0.15818  

V:Block:B 25  2833     113   1.2451   0.29390
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

V       4 102743  25686 282.2094 < 2.2e-16 ***  

V:Block 25 50019   2001 21.9825 1.588e-11 ***  

A       1 18451   18451 202.7233 1.692e-13 ***  

B       1 78541   78541 862.9280 < 2.2e-16 ***  

A:B     1   108     108   1.1899  0.28575  

V:A     4   3751     938  10.3023 4.532e-05 ***  

V:B     4   307      77   0.8421   0.51168  

V:A:B    4   1495     374   4.1058   0.01081 *  

V:Block:A 25  3416     137   1.5011   0.15818  

V:Block:B 25  2833     113   1.2451   0.29390
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.6.2 p434

(119) MODEL

```
GLM(Y ~ V + Block:V + A + B + A:B + V:A + V:B + V:A:B, v1p432) # OK
```

```
$ANOVA  
Response : Y
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	44	255415	5804.9	51.075	< 2.2e-16 ***
RESIDUALS	75	8524	113.7		
CORRECTED TOTAL	119	263939			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	Y Mean
0.9677043	1.739417	10.66088	612.9		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
A	1	18451	18451	162.3447	< 2.2e-16 ***
B	1	78541	78541	691.0494	< 2.2e-16 ***
A:B	1	108	108	0.9529	0.33212
V:A	4	3751	938	8.2503	1.435e-05 ***
V:B	4	307	77	0.6744	0.61182
V:A:B	4	1495	374	3.2880	0.01541 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
A	1	18451	18451	162.3447	< 2.2e-16 ***
B	1	78541	78541	691.0494	< 2.2e-16 ***
A:B	1	108	108	0.9529	0.33212
V:A	4	3751	938	8.2503	1.435e-05 ***
V:B	4	307	77	0.6744	0.61182
V:A:B	4	1495	374	3.2880	0.01541 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
A	1	18451	18451	162.3447	< 2.2e-16 ***
B	1	78541	78541	691.0494	< 2.2e-16 ***
A:B	1	108	108	0.9529	0.33212
V:A	4	3751	938	8.2503	1.435e-05 ***
V:B	4	307	77	0.6744	0.61182
V:A:B	4	1495	374	3.2880	0.01541 *

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.6.3 p438

(120) MODEL

```
GLM(Y ~ V + Block:V + C + V:C, v1p432) # OK
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	44	255415	5804.9	51.075	< 2.2e-16 ***
RESIDUALS	75	8524	113.7		
CORRECTED TOTAL	119	263939			

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

R-square	Coef	Var	Root	MSE	Y Mean
0.9677043	1.739417	10.66088	612.9		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
C	3	97100	32367	284.7823	< 2.2e-16 ***
V:C	12	5552	463	4.0709	7.23e-05 ***

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
C	3	97100	32367	284.7823	< 2.2e-16 ***
V:C	12	5552	463	4.0709	7.23e-05 ***

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
C	3	97100	32367	284.7823	< 2.2e-16 ***
V:C	12	5552	463	4.0709	7.23e-05 ***

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.6.4 p444

(121) MODEL

```
v1p444 = v1p432[v1p432$Block==5,]
GLM(Y ~ V + A + B + A:B + V:A, v1p444) # OK
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 39278  3570.8 59.787 1.897e-06 ***
RESIDUALS     8   478    59.7
CORRECTED TOTAL 19 39756
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE Y Mean
  0.9879817 1.225336 7.728195 630.7

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***
A      1 3380.0 3380.0 56.5927 6.780e-05 ***
B      1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B    1   115.2   115.2   1.9288  0.202326
V:A    4  2450.5   612.6   10.2574  0.003081 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***
A      1 3380.0 3380.0 56.5927 6.780e-05 ***
B      1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B    1   115.2   115.2   1.9288  0.202326
V:A    4  2450.5   612.6   10.2574  0.003081 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***
A      1 3380.0 3380.0 56.5927 6.780e-05 ***
B      1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B    1   115.2   115.2   1.9288  0.202326
V:A    4  2450.5   612.6   10.2574  0.003081 **
```

```
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.6.5 p482

(122) MODEL

```
v1p482 = read.table("C:/G/Rt/Kemp/v1p482.txt", head=TRUE)
v1p482 = af(v1p482,c("block", "A", "B"))
GLM(y ~ block + A + B + A:B, v1p482) # OK
```

```
$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL     8 156.88 19.6094 9.8871 9.377e-05 ***
RESIDUALS 15 29.75  1.9833
CORRECTED TOTAL 23 186.62
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
R-square Coef Var Root MSE y Mean
0.8405894 17.88328 1.408309 7.875
```

```
$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 108.38 21.675 10.9286 0.0001415 ***
A       1    4.00   4.000  2.0168 0.1760166
B       1   42.25  42.250 21.3025 0.0003365 ***
A:B     1    2.25   2.250  1.1345 0.3036727
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 31.417   6.283   3.1681 0.0377804 *
A       1   4.000   4.000   2.0168 0.1760166
B       1  42.250  42.250  21.3025 0.0003365 ***
A:B     1   2.250   2.250   1.1345 0.3036727
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 31.417   6.283   3.1681 0.0377804 *
A       1   4.000   4.000   2.0168 0.1760166
```

```

B      1 42.250 42.250 21.3025 0.0003365 ***
A:B    1  2.250   2.250  1.1345 0.3036727
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.7 Chapter 12

8.7.1 p525

(123) MODEL

```

v1p525 = read.table("C:/G/Rt/Kemp/v1p525.txt", head=TRUE)
REG(y ~ x1 + x2 + x3, v1p525)

```

```

Estimate Std. Error Df t value Pr(>|t|)
(Intercept) 14.2125    0.10383 12 136.8787 < 2.2e-16 ***
x1          0.7875    0.10383 12   7.5843 6.465e-06 ***
x2          1.3875    0.10383 12  13.3628 1.446e-08 ***
x3          1.6625    0.10383 12  16.0113 1.839e-09 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
GLM(y ~ x1 + x2 + x3, v1p525) # OK
```

```

$ANOVA
Response : y
Df Sum Sq Mean Sq F value Pr(>F)
MODEL       3 84.948 28.3158 164.15 5.26e-10 ***
RESIDUALS   12  2.070  0.1725
CORRECTED TOTAL 15 87.018
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE y Mean
0.9762117 2.922295 0.4153312 14.2125

```

```

$`Type I`
Df Sum Sq Mean Sq F value     Pr(>F)
x1  1  9.923  9.923  57.522 6.465e-06 ***
x2  1 30.803 30.803 178.565 1.446e-08 ***
x3  1 44.223 44.223 256.362 1.839e-09 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

x1  1  9.923   9.923  57.522 6.465e-06 ***  

x2  1 30.803  30.803 178.565 1.446e-08 ***  

x3  1 44.223  44.223 256.362 1.839e-09 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

x1  1  9.923   9.923  57.522 6.465e-06 ***  

x2  1 30.803  30.803 178.565 1.446e-08 ***  

x3  1 44.223  44.223 256.362 1.839e-09 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.7.2 p527

(124) MODEL

```
v1p527 = read.table("C:/G/Rt/Kemp/v1p527.txt", head=TRUE)  

GLM(y ~ A + B, v1p527) # OK
```

```
$ANOVA  

Response : y  

  Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      2  22.99  11.4952  4.8917 0.04686 *  

RESIDUALS  7  16.45   2.3499  

CORRECTED TOTAL 9  39.44  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE y Mean  

  0.5829197 29.47989 1.532954     5.2
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

A  1 10.364  10.364  4.4103 0.07386 .  

B  1 12.626  12.626  5.3730 0.05355 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

A  1 10.364  10.364  4.4103 0.07386 .
```

```

B 1 12.626 12.626 5.3730 0.05355 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 10.364 10.364 4.4103 0.07386 .
B 1 12.626 12.626 5.3730 0.05355 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.7.3 p529

(125) MODEL

```
v1p529 = read.table("C:/G/Rt/Kemp/v1p529.txt", head=TRUE)
GLM(y ~ A + B + I(A*A) + I(B*B) + I(A*B), v1p529) # OK
```

```
$ANOVA
Response : y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      5 35.713 7.1427 6.7928 0.01857 *
RESIDUALS   6  6.309 1.0515
CORRECTED TOTAL 11 42.023
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
  R-square Coef Var Root MSE y Mean
  0.8498641 19.4395 1.025434 5.275
```

```
$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
A      1 11.6012 11.6012 11.0329 0.01597 *
B      1 12.6263 12.6263 12.0077 0.01338 *
I(A * A) 1 1.7167 1.7167 1.6326 0.24855
I(B * B) 1 5.3593 5.3593 5.0967 0.06476 .
I(A * B) 1 4.4100 4.4100 4.1940 0.08649 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
A      1 11.6012 11.6012 11.0329 0.01597 *
B      1 12.6263 12.6263 12.0077 0.01338 *
I(A * A) 1 5.5468 5.5468 5.2750 0.06137 .
```

```

I(B * B) 1 5.3593 5.3593 5.0967 0.06476 .
I(A * B) 1 4.4100 4.4100 4.1940 0.08649 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

A       1 11.6012 11.6012 11.0329 0.01597 *  

B       1 12.6263 12.6263 12.0077 0.01338 *  

I(A * A) 1 5.5468 5.5468 5.2750 0.06137 .  

I(B * B) 1 5.3593 5.3593 5.0967 0.06476 .  

I(A * B) 1 4.4100 4.4100 4.1940 0.08649 .  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.8 Chapter 13

8.8.1 p563

(126) MODEL

```

v1p563 = read.table("C:/G/Rt/Kemp/v1p563.txt", head=TRUE)
v1p563 = af(v1p563, c("rep", "A", "B"))
GLM(y ~ rep + A + rep:A + B + A:B, v1p563) # OK

```

```

$ANOVA  

Response : y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      14 2097.08 149.792 17.228 8.385e-05 ***  

RESIDUALS    9   78.25   8.694  

CORRECTED TOTAL 23 2175.33  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE   y Mean  

  0.9640285 9.460859 2.948634 31.16667

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

rep      3 1241.00 413.67 47.5783 7.606e-06 ***  

A       2  353.08 176.54 20.3051 0.0004613 ***  

rep:A    6  192.25  32.04  3.6853 0.0393557 *  

B       1  216.00 216.00 24.8435 0.0007550 ***  

A:B     2   94.75  47.38  5.4489 0.0281496 *
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

rep     3 1241.00  413.67 47.5783 7.606e-06 ***  

A       2   353.08  176.54 20.3051 0.0004613 ***  

rep:A   6   192.25   32.04  3.6853 0.0393557 *  

B       1   216.00  216.00 24.8435 0.0007550 ***  

A:B     2    94.75   47.38  5.4489 0.0281496 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

rep     3 1241.00  413.67 47.5783 7.606e-06 ***  

A       2   353.08  176.54 20.3051 0.0004613 ***  

rep:A   6   192.25   32.04  3.6853 0.0393557 *  

B       1   216.00  216.00 24.8435 0.0007550 ***  

A:B     2    94.75   47.38  5.4489 0.0281496 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.8.2 p566

(127) MODEL

```

v1p566 = read.table("C:/G/Rt/Kemp/v1p566.txt", head=TRUE)
v1p566 = af(v1p566, c("subject", "A", "B"))
GLM(y ~ A + B + A:B, v1p566) # OK

```

```

$ANOVA  

Response : y  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      5 1469.58  293.92     86.2 5.592e-09 ***  

RESIDUALS   12   40.92    3.41  

CORRECTED TOTAL 17 1510.50  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE    y Mean  

0.9729118 5.153144 1.846543 35.83333

```

```

$`Type I`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

A       2 1390.04  695.02 203.8350 5.466e-10 ***

```

```

B     1    76.06    76.06   22.3055 0.0004945 ***
A:B   2     3.49     1.74    0.5112 0.6122667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq Mean Sq F value    Pr(>F)
A     2 1390.04 695.02 203.8350 5.466e-10 ***
B     1    76.06    76.06   22.3055 0.0004945 ***
A:B   2     3.49     1.74    0.5112 0.6122667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df  Sum Sq Mean Sq F value    Pr(>F)
A     2 1390.04 695.02 203.8350 5.466e-10 ***
B     1    79.00    79.00   23.1700 0.0004237 ***
A:B   2     3.49     1.74    0.5112 0.6122667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.9 Chapter 14

8.9.1 p581

(128) MODEL

```

v1p581 = read.table("C:/G/Rt/Kemp/v1p581.txt", head=TRUE)
v1p581 = af(v1p581, c("drug", "person", "time"))
GLM(rate ~ drug + person:drug + time + drug:time, v1p581) # OK

```

```

$ANOVA
Response : rate
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      23 2449.5 106.500 12.733 3.469e-11 ***
RESIDUALS  36  301.1   8.364
CORRECTED TOTAL 59 2750.6
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE rate Mean
0.890533 3.722058 2.892039      77.7

```

```

$`Type I`
  Df  Sum Sq Mean Sq F value    Pr(>F)

```

```
drug          2 337.60 168.800 20.1820 1.323e-06 ***
drug:person 12 1498.50 124.875 14.9303 1.501e-10 ***
time         3 256.33  85.444 10.2159 5.230e-05 ***
drug:time    6 357.07  59.511  7.1152 4.707e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
drug	2	337.60	168.800	20.1820	1.323e-06 ***
drug:person	12	1498.50	124.875	14.9303	1.501e-10 ***
time	3	256.33	85.444	10.2159	5.230e-05 ***
drug:time	6	357.07	59.511	7.1152	4.707e-05 ***

```
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
drug	2	337.60	168.800	20.1820	1.323e-06 ***
drug:person	12	1498.50	124.875	14.9303	1.501e-10 ***
time	3	256.33	85.444	10.2159	5.230e-05 ***
drug:time	6	357.07	59.511	7.1152	4.707e-05 ***

```
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9 Hinkelmann & Kempthorne - Volume 2

Reference - Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 2 Advanced Experimental Design. 2e. John Wiley & Sons Inc. 2008.

9.1 Chapter 1

9.1.1 p53

(129) MODEL

```
v2p53 = read.table("C:/G/Rt/Kemp/v2p53.txt", head=TRUE)
v2p53 = af(v2p53, c("TRT", "BLOCK"))
GLM(Y ~ BLOCK + TRT, v2p53) # OK
```

\$ANOVA
Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	7	518.21	74.030	8.1408	0.1137
RESIDUALS	2	18.19	9.094		
CORRECTED TOTAL	9	536.40			

\$Fitness
R-square Coef Var Root MSE Y Mean
0.9660934 15.54425 3.015585 19.4

\$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
BLOCK 4 261.40 65.350 7.1863 0.12587
TRT 3 256.81 85.604 9.4135 0.09755 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
BLOCK 4 79.146 19.786 2.1758 0.33880
TRT 3 256.812 85.604 9.4135 0.09755 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
BLOCK 4 79.146 19.786 2.1758 0.33880
TRT 3 256.813 85.604 9.4135 0.09755 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

9.1.2 p62

(130) MODEL

```
GLM(Y ~ TRT + BLOCK, v2p53) # OK
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	7	518.21	74.030	8.1408	0.1137
RESIDUALS	2	18.19	9.094		
CORRECTED TOTAL	9	536.40			

\$Fitness

R-square	Coef	Var	Root	MSE	Y Mean
0.9660934	15.54425	3.015585		19.4	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
TRT	3	439.07	146.356	16.0941	0.05907 .
BLOCK	4	79.15	19.786	2.1758	0.33880

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
TRT	3	256.812	85.604	9.4135	0.09755 .
BLOCK	4	79.146	19.786	2.1758	0.33880

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
TRT	3	256.813	85.604	9.4135	0.09755 .
BLOCK	4	79.146	19.786	2.1758	0.33880

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

9.2 Chapter 2

9.2.1 p82

(131) MODEL

```

v2p82 = read.table("C:/G/Rt/Kemp/v2p82.txt", head=TRUE)
v2p82 = af(v2p82, c("B", "Tx"))
GLM(Y ~ B + Tx, v2p82) # OK

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      14  889.11  63.508  6.3183 0.000518 ***
RESIDUALS   15  150.77  10.052
CORRECTED TOTAL 29 1039.89
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   Y Mean
0.8550104 8.241975 3.170413 38.46667

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
B     9  730.39  81.154  8.0738 0.0002454 ***
Tx    5  158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
B     9  595.74  66.193  6.5854 0.0007602 ***
Tx    5  158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
B     9  595.74  66.193  6.5854 0.0007602 ***
Tx    5  158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.2.2 p87

(132) MODEL

```

v2p87 = read.table("C:/G/Rt/Kemp/v2p87.txt", head=TRUE)
GLM(y ~ x1 + x2 + x3 + x4 + x5 + x6, v2p87) # OK

```

\$ANOVA

```

Response : y
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      5  1613.25  322.65  2.2332 0.2282
RESIDUALS   4   577.91  144.48
CORRECTED TOTAL 9  2191.16

$Fitness
  R-square Coef Var Root MSE y Mean
0.7362523 10.41587 12.01991 115.4

$`Type I`
  Df  Sum Sq Mean Sq F value Pr(>F)
x1  1 1044.48 1044.48  7.2293 0.05473 .
x2  1   89.79   89.79  0.6215 0.47459
x3  1   10.45   10.45  0.0724 0.80124
x4  1  407.08  407.08  2.8176 0.16854
x5  1   61.44   61.44  0.4253 0.54990
x6  0
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq Mean Sq F value Pr(>F)
x1  0
x2  0
x3  0
x4  0
x5  0
x6  0

$`Type III`
CAUTION: Singularity Exists !
  Df  Sum Sq Mean Sq F value Pr(>F)
x1  0
x2  0
x3  0
x4  0
x5  0
x6  0

```

9.3 Chapter 6

9.3.1 p217

(133) MODEL

```
v2p217 = read.table("C:/G/Rt/Kemp/v2p217.txt", head=TRUE)
v2p217 = af(v2p217, c("R", "C", "Tx"))
GLM(Y ~ R + C + Tx, v2p217) # OK
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      22 4305.1 195.687 7.5094 0.0002682 ***
RESIDUALS   13 338.8  26.059
CORRECTED TOTAL 35 4643.9
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  Y Mean
  0.9270507 18.86789 5.104813 27.05556

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
R     3 3951.4 1317.15 50.5446 1.998e-07 ***
C     8 168.9   21.11  0.8101   0.6062
Tx   11 184.8   16.80  0.6446   0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
R     3 3403.5 1134.51 43.5360 4.83e-07 ***
C     8 112.4   14.05  0.5390   0.8077
Tx   11 184.8   16.80  0.6446   0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
R     3 3403.5 1134.51 43.5360 4.83e-07 ***
C     8 112.4   14.05  0.5390   0.8077
Tx   11 184.8   16.80  0.6446   0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.3.2 p234

(134) MODEL

```
v2p234 = read.table("C:/G/Rt/Kemp/v2p234.txt", head=TRUE)
v2p234 = af(v2p234, c("R", "C", "Tx"))
GLM(Y ~ C + R + Tx, v2p234) # OK
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      13 426.50 32.808 7.0936 0.1302
RESIDUALS   2   9.25  4.625
CORRECTED TOTAL 15 435.75

$Fitness
  R-square Coef Var Root MSE Y Mean
0.9787722 7.259346 2.150581 29.625

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
C   3 16.25  5.417  1.1712 0.49129
R   3 357.25 119.083 25.7477 0.03762 *
Tx  7 53.00  7.571  1.6371 0.43052
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
C   3 10.25  3.417  0.7387 0.6189
R   3 285.50 95.167 20.5766 0.0467 *
Tx  7 53.00  7.571  1.6371 0.4305
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
C   3 10.25  3.417  0.7387 0.6189
R   3 285.50 95.167 20.5766 0.0467 *
Tx  7 53.00  7.571  1.6371 0.4305
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.4 Chapter 7

9.4.1 p268

(135) MODEL

```
v2p268 = read.table("C:/G/Rt/Kemp/v2p268.txt", head=TRUE)
v2p268 = af(v2p268, c("A", "B", "C"))
GLM(y ~ block + A*B*C, v2p268) # OK
```

\$ANOVA
 Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	8	1026.00	128.250	24.981	0.0001765 ***
RESIDUALS	7	35.94	5.134		
CORRECTED TOTAL	15	1061.94			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	y	Mean
0.9661586	8.863833	2.265817	25.5625			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	715.56	715.56	139.3791	7.093e-06 ***
A	1	68.06	68.06	13.2574	0.0082753 **
B	1	0.06	0.06	0.0122	0.9152401
A:B	1	0.56	0.56	0.1096	0.7503276
C	1	232.56	232.56	45.2991	0.0002698 ***
A:C	1	0.06	0.06	0.0122	0.9152401
B:C	1	7.56	7.56	1.4730	0.2642229
A:B:C	1	1.56	1.56	0.3043	0.5983312

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	715.56	715.56	139.3791	7.093e-06 ***
A	1	68.06	68.06	13.2574	0.0082753 **
B	1	0.06	0.06	0.0122	0.9152401
A:B	1	0.56	0.56	0.1096	0.7503276
C	1	232.56	232.56	45.2991	0.0002698 ***
A:C	1	0.06	0.06	0.0122	0.9152401
B:C	1	7.56	7.56	1.4730	0.2642229
A:B:C	1	1.56	1.56	0.3043	0.5983312

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	715.56	715.56	139.3791	7.093e-06 ***
A	1	68.06	68.06	13.2574	0.0082753 **

```

B      1   0.06   0.06   0.0122  0.9152401
A:B    1   0.56   0.56   0.1096  0.7503276
C      1 232.56  232.56  45.2991  0.0002698 ***
A:C    1   0.06   0.06   0.0122  0.9152401
B:C    1   7.56   7.56   1.4730  0.2642229
A:B:C  1   1.56   1.56   0.3043  0.5983312
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.4.2 p273

(136) MODEL

```

v2p273 = read.table("C:/G/Rt/Kemp/v2p273.txt", head=TRUE)
v2p273 = af(v2p273, c("block", "A", "B", "C"))
GLM(y ~ block + A*B*C + block:A:B:C, v2p273) # OK

```

```

$ANOVA
Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      15 2245.0 149.665 129.44 8.427e-14 ***
RESIDUALS  16   18.5   1.156
CORRECTED TOTAL 31 2263.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE   y Mean
0.9918267 4.170824 1.075291 25.78125

```

```

$`Type I`
          Df Sum Sq Mean Sq  F value    Pr(>F)
block      1 1498.78 1498.78 1296.2432 < 2.2e-16 ***
A          1 132.03 132.03  114.1892 1.083e-08 ***
B          1   0.03   0.03   0.0270   0.87148
A:B        1   1.53   1.53   1.3243   0.26673
C          1 504.03 504.03  435.9189 4.926e-13 ***
A:C        1   0.78   0.78   0.6757   0.42316
B:C        1   3.78   3.78   3.2703   0.08938 .
A:B:C     1   2.53   2.53   2.1892   0.15840
block:A:B:C 7 101.47  14.50   12.5367 1.965e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
          Df Sum Sq Mean Sq  F value    Pr(>F)

```

```

block      1 1498.78 1498.78 1296.2432 < 2.2e-16 ***
A         1 132.03 132.03 114.1892 1.083e-08 ***
B         1 0.03 0.03 0.0270 0.87148
A:B       1 1.53 1.53 1.3243 0.26673
C         1 504.03 504.03 435.9189 4.926e-13 ***
A:C       1 0.78 0.78 0.6757 0.42316
B:C       1 3.78 3.78 3.2703 0.08938 .
A:B:C     1 2.53 2.53 2.1892 0.15840
block:A:B:C 7 101.47 14.50 12.5367 1.965e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block      1 1498.78 1498.78 1296.2432 < 2.2e-16 ***  

A         1 132.03 132.03 114.1892 1.083e-08 ***  

B         1 0.03 0.03 0.0270 0.87148  

A:B       1 1.53 1.53 1.3243 0.26673  

C         1 504.03 504.03 435.9189 4.926e-13 ***  

A:C       1 0.78 0.78 0.6757 0.42316  

B:C       1 3.78 3.78 3.2703 0.08938 .  

A:B:C     1 2.53 2.53 2.1892 0.15840  

block:A:B:C 7 101.47 14.50 12.5367 1.965e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.5 Chapter 8

9.5.1 p304

(137) MODEL

```

v2p304 = read.table("C:/G/Rt/Kemp/v2p304.txt", head=TRUE)
v2p304 = af(v2p304, c("rep", "block", "A", "B", "C"))
GLM(y ~ rep + block %in% rep + A*B*C - A:B:C, v2p304) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL    9 699.06 77.674 248.56 5.096e-07 ***
RESIDUALS 6 1.88 0.312
CORRECTED TOTAL 15 700.94
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness

```

```

R-square Coef Var Root MSE  y Mean
0.997325 2.423922 0.559017 23.0625

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep       1 390.06 390.06 1248.2 3.428e-08 ***  

rep:block 2   8.12   4.06   13.0 0.0065918 **  

A         1 18.06 18.06   57.8 0.0002696 ***  

B         1 175.56 175.56  561.8 3.702e-07 ***  

A:B       1   0.06   0.06    0.2 0.6704121  

C         1 68.06 68.06  217.8 6.083e-06 ***  

A:C       1   0.06   0.06    0.2 0.6704121  

B:C       1 39.06 39.06  125.0 3.056e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep       1 390.06 390.06 1248.2 3.428e-08 ***  

rep:block 2   8.12   4.06   13.0 0.0065918 **  

A         1 18.06 18.06   57.8 0.0002696 ***  

B         1 175.56 175.56  561.8 3.702e-07 ***  

A:B       1   0.06   0.06    0.2 0.6704121  

C         1 68.06 68.06  217.8 6.083e-06 ***  

A:C       1   0.06   0.06    0.2 0.6704121  

B:C       1 39.06 39.06  125.0 3.056e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep       1 390.06 390.06 1248.2 3.428e-08 ***  

rep:block 2   8.12   4.06   13.0 0.0065918 **  

A         1 18.06 18.06   57.8 0.0002696 ***  

B         1 175.56 175.56  561.8 3.702e-07 ***  

A:B       1   0.06   0.06    0.2 0.6704121  

C         1 68.06 68.06  217.8 6.083e-06 ***  

A:C       1   0.06   0.06    0.2 0.6704121  

B:C       1 39.06 39.06  125.0 3.056e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.5.2 p309

(138) MODEL

```
GLM(y ~ rep*A*B*C, v2p304) # OK
```

\$ANOVA

```
Response : y
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          15 700.94 46.729
RESIDUALS      0   0.00
CORRECTED TOTAL 15 700.94
```

\$Fitness

```
R-square Coef Var Root MSE y Mean
           1       NA       NA 23.0625
```

\$`Type I`

```
Df Sum Sq Mean Sq F value Pr(>F)
rep      1 390.06 390.06
A        1 18.06 18.06
rep:A    1 0.06 0.06
B        1 175.56 175.56
rep:B    1 1.56 1.56
A:B     1 0.06 0.06
rep:A:B  1 0.06 0.06
C        1 68.06 68.06
rep:C    1 0.06 0.06
A:C     1 0.06 0.06
rep:A:C  1 0.06 0.06
B:C     1 39.06 39.06
rep:B:C  1 0.06 0.06
A:B:C   1 7.56 7.56
rep:A:B:C 1 0.56 0.56
```

\$`Type II`

```
Df Sum Sq Mean Sq F value Pr(>F)
rep      1 390.06 390.06
A        1 18.06 18.06
rep:A    1 0.06 0.06
B        1 175.56 175.56
rep:B    1 1.56 1.56
A:B     1 0.06 0.06
rep:A:B  1 0.06 0.06
C        1 68.06 68.06
rep:C    1 0.06 0.06
A:C     1 0.06 0.06
rep:A:C  1 0.06 0.06
B:C     1 39.06 39.06
rep:B:C  1 0.06 0.06
A:B:C   1 7.56 7.56
```

```

rep:A:B:C 1 0.56 0.56

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

rep        1 390.06 390.06  

A          1 18.06 18.06  

rep:A      1 0.06 0.06  

B          1 175.56 175.56  

rep:B      1 1.56 1.56  

A:B        1 0.06 0.06  

rep:A:B    1 0.06 0.06  

C          1 68.06 68.06  

rep:C      1 0.06 0.06  

A:C        1 0.06 0.06  

rep:A:C    1 0.06 0.06  

B:C        1 39.06 39.06  

rep:B:C   1 0.06 0.06  

A:B:C     1 7.56 7.56  

rep:A:B:C 1 0.56 0.56

```

9.6 Chapter 9

9.6.1 p343

(139) MODEL

```

v2p343 = read.table("C:/G/Rt/Kemp/v2p343.txt", head=TRUE)
v2p343 = af(v2p343, c("rep", "block", "A", "B", "C"))
GLM(y ~ rep + block %in% rep + A*B*C - A:B:C, v2p343) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       17 1889.8 111.167 14.659 0.001608 ***
RESIDUALS    6   45.5   7.583
CORRECTED TOTAL 23 1935.3
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   y Mean
0.9764898 12.70978 2.753785 21.66667

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)
rep        2 1537.33 768.67 101.3626 2.375e-05 ***

```

```

rep:block 9 127.00 14.11 1.8608 0.23163
A 1 36.00 36.00 4.7473 0.07218 .
B 1 36.00 36.00 4.7473 0.07218 .
A:B 1 12.25 12.25 1.6154 0.25079
C 1 56.25 56.25 7.4176 0.03448 *
A:C 1 81.00 81.00 10.6813 0.01707 *
B:C 1 4.00 4.00 0.5275 0.49502
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`:
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 1537.33 768.67 101.3626 2.375e-05 ***
rep:block 9 119.83 13.31 1.7558 0.25388
A 1 36.00 36.00 4.7473 0.07218 .
B 1 36.00 36.00 4.7473 0.07218 .
A:B 1 12.25 12.25 1.6154 0.25079
C 1 56.25 56.25 7.4176 0.03448 *
A:C 1 81.00 81.00 10.6813 0.01707 *
B:C 1 4.00 4.00 0.5275 0.49502
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`:
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 1537.33 768.67 101.3626 2.375e-05 ***
rep:block 9 119.83 13.31 1.7558 0.25388
A 1 36.00 36.00 4.7473 0.07218 .
B 1 36.00 36.00 4.7473 0.07218 .
A:B 1 12.25 12.25 1.6154 0.25079
C 1 56.25 56.25 7.4176 0.03448 *
A:C 1 81.00 81.00 10.6813 0.01707 *
B:C 1 4.00 4.00 0.5275 0.49502
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.6.2 p348

(140) MODEL

```
GLM(y ~ rep + A*B*C + block %in% rep, v2p343) # OK
```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL 17 1889.8 111.167 14.659 0.001608 **
```

```

RESIDUALS      6   45.5   7.583
CORRECTED TOTAL 23 1935.3
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE   y Mean
  0.9764898 12.70978 2.753785 21.66667

$`Type I`
  Df  Sum Sq Mean Sq  F value    Pr(>F)
rep      2 1537.33  768.67 101.3626 2.375e-05 ***
A        1   88.17   88.17  11.6264  0.01432 *
B        1   37.50   37.50   4.9451  0.06785 .
A:B      1   2.67    2.67   0.3516  0.57484
C        1   66.67   66.67   8.7912  0.02512 *
A:C      1   37.50   37.50   4.9451  0.06785 .
B:C      1   0.17    0.17   0.0220  0.88700
A:B:C    1   24.00   24.00   3.1648  0.12555
rep:block 8   95.83   11.98   1.5797  0.29730
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq Mean Sq  F value    Pr(>F)
rep      2 1537.33  768.67 101.3626 2.375e-05 ***
A        1   36.00   36.00   4.7473  0.07218 .
B        1   36.00   36.00   4.7473  0.07218 .
A:B      1   12.25   12.25   1.6154  0.25079
C        1   56.25   56.25   7.4176  0.03448 *
A:C      1   81.00   81.00  10.6813  0.01707 *
B:C      1   4.00    4.00   0.5275  0.49502
A:B:C    0
rep:block 8   95.83   11.98   1.5797  0.29730
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
  Df  Sum Sq Mean Sq  F value    Pr(>F)
rep      2 1537.33  768.67 101.3626 2.375e-05 ***
A        1   36.00   36.00   4.7473  0.07218 .
B        1   36.00   36.00   4.7473  0.07218 .
A:B      1   12.25   12.25   1.6154  0.25079
C        1   56.25   56.25   7.4176  0.03448 *
A:C      1   81.00   81.00  10.6813  0.01707 *
B:C      1   4.00    4.00   0.5275  0.49502
A:B:C    0

```

```

rep:block 8 95.83 11.98 1.5797 0.29730
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.6.3 p353

(141) MODEL

```

v2p353 = read.table("C:/G/Rt/Kemp/v2p353.txt", head=TRUE)
v2p353 = af(v2p353, c("rep", "block", "A", "B", "C", "D"))
GLM(y ~ rep + rep:block + A*B*C*D - A:B:C:D, v2p353) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      21 7132.2 339.63 56.022 9.795e-08 ***
RESIDUALS   10   60.6    6.06
CORRECTED TOTAL 31 7192.9
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE  y Mean
0.9915715 6.621081 2.462214 37.1875

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 5940.5 5940.5 979.8763 2.600e-11 ***
rep:block 6 777.4   129.6 21.3711 3.675e-05 ***
A         1 171.1   171.1 28.2268 0.0003412 ***
B         1   18.0    18.0  2.9691 0.1155937
A:B       1     1.6    1.6  0.2577 0.6226914
C         1 120.1   120.1 19.8144 0.0012326 **
A:C       1     0.6    0.6  0.0928 0.7669127
B:C       1     2.0    2.0  0.3299 0.5784103
A:B:C     1     4.5    4.5  0.7423 0.4091189
D         1     6.1    6.1  1.0103 0.3385304
A:D       1     1.1    1.1  0.1856 0.6757693
B:D       1     5.1    5.1  0.8351 0.3823203
A:B:D     1     0.5    0.5  0.0825 0.7798349
C:D       1     1.6    1.6  0.2577 0.6226914
A:C:D     1    10.1   10.1  1.6701 0.2253083
B:C:D     1    72.0   72.0 11.8763 0.0062660 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep        1 5940.5  5940.5 979.8763  2.6e-11 ***  

rep:block  6  406.9    67.8  11.1856 0.0006129 ***  

A          1  171.1   171.1  28.2268 0.0003412 ***  

B          1   18.0    18.0   2.9691 0.1155937  

A:B        1    1.6    1.6   0.2577 0.6226914  

C          1  120.1   120.1  19.8144 0.0012326 **  

A:C        1    0.6    0.6   0.0928 0.7669127  

B:C        1    2.0    2.0   0.3299 0.5784103  

A:B:C     1    4.5    4.5   0.7423 0.4091189  

D          1    6.1    6.1   1.0103 0.3385304  

A:D        1    1.1    1.1   0.1856 0.6757693  

B:D        1    5.1    5.1   0.8351 0.3823203  

A:B:D     1    0.5    0.5   0.0825 0.7798349  

C:D        1    1.6    1.6   0.2577 0.6226914  

A:C:D     1   10.1   10.1   1.6701 0.2253083  

B:C:D     1   72.0   72.0  11.8763 0.0062660 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep        1 5940.5  5940.5 979.8763  2.6e-11 ***  

rep:block  6  406.9    67.8  11.1856 0.0006129 ***  

A          1  171.1   171.1  28.2268 0.0003412 ***  

B          1   18.0    18.0   2.9691 0.1155937  

A:B        1    1.6    1.6   0.2577 0.6226914  

C          1  120.1   120.1  19.8144 0.0012326 **  

A:C        1    0.6    0.6   0.0928 0.7669127  

B:C        1    2.0    2.0   0.3299 0.5784103  

A:B:C     1    4.5    4.5   0.7423 0.4091189  

D          1    6.1    6.1   1.0103 0.3385304  

A:D        1    1.1    1.1   0.1856 0.6757693  

B:D        1    5.1    5.1   0.8351 0.3823203  

A:B:D     1    0.5    0.5   0.0825 0.7798349  

C:D        1    1.6    1.6   0.2577 0.6226914  

A:C:D     1   10.1   10.1   1.6701 0.2253083  

B:C:D     1   72.0   72.0  11.8763 0.0062660 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.7 Chapter 10

9.7.1 p388

(142) MODEL

```
v2p388 = read.table("C:/G/Rt/Kemp/v2p388.txt", head=TRUE)
v2p388 = af(v2p388, c("rep", "block", "A", "B"))
GLM(y ~ rep + A*B + rep:block, v2p388) # OK
```

\$ANOVA

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	11	1136.8	103.343	124.01	3.698e-06 ***
RESIDUALS	6	5.0	0.833		
CORRECTED TOTAL	17	1141.8			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root MSE	y Mean
0.9956209	3.496101	0.9128709	26.11111	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	1	410.89	410.89	493.0667	5.455e-07 ***
A	2	228.11	114.06	136.8667	9.868e-06 ***
B	2	3.44	1.72	2.0667	0.207585
A:B	4	464.22	116.06	139.2667	4.801e-06 ***
rep:block	2	30.11	15.06	18.0667	0.002888 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	1	410.89	410.89	493.0667	5.455e-07 ***
A	2	228.11	114.06	136.8667	9.868e-06 ***
B	2	3.44	1.72	2.0667	0.207585
A:B	2	18.78	9.39	11.2667	0.009298 **
rep:block	2	30.11	15.06	18.0667	0.002888 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	1	410.89	410.89	493.0667	5.455e-07 ***
A	2	228.11	114.06	136.8667	9.868e-06 ***
B	2	3.44	1.72	2.0667	0.207585
A:B	2	18.78	9.39	11.2667	0.009298 **
rep:block	2	30.11	15.06	18.0667	0.002888 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

9.8 Chapter 14

9.8.1 p570

(143) MODEL

```
v2p570 = read.table("C:/G/Rt/Kemp/v2p570.txt", head=TRUE)
v2p570 = af(v2p570, c("A", "B", "C", "D"))
GLM(Y ~ A + B + C + D + A:B + A:C + A:D + B:C + B:D + C:D, v2p570) # OK
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      8 22.222 2.7778
RESIDUALS   0  0.000
CORRECTED TOTAL 8 22.222
```

```
$Fitness
R-square Coef Var Root MSE   Y Mean
      1       NA     NA 6.555556
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
A      2 2.8889 1.4444
B      2 2.8889 1.4444
C      2 1.5556 0.7778
D      2 14.8889 7.4444
A:B    0
A:C    0
A:D    0
B:C    0
B:D    0
C:D    0
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
A      0
B      0
C      0
D      0
A:B    0
A:C    0
A:D    0
B:C    0
B:D    0
C:D    0
```

```
$`Type III`  
CAUTION: Singularity Exists !  
Df Sum Sq Mean Sq F value Pr(>F)  
A 0  
B 0  
C 0  
D 0  
A:B 0  
A:C 0  
A:D 0  
B:C 0  
B:D 0  
C:D 0
```

9.8.2 p578

(144) MODEL

```
v2p578 = read.table("C:/G/Rt/Kemp/v2p578.txt", head=TRUE)  
v2p578 = af(v2p578, 1:11)  
GLM(Y ~ A + B + C + D + E + F + G + H + J + K + L, v2p578) # OK
```

```
$ANOVA  
Response : Y  
Df Sum Sq Mean Sq F value Pr(>F)  
MODEL 11 575 52.273  
RESIDUALS 0 0  
CORRECTED TOTAL 11 575
```

```
$Fitness  
R-square Coef Var Root MSE Y Mean  
1 NA NA 25.5
```

```
$`Type I`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 3.000 3.000  
B 1 27.000 27.000  
C 1 12.000 12.000  
D 1 16.333 16.333  
E 1 176.333 176.333  
F 1 133.333 133.333  
G 1 1.333 1.333  
H 1 21.333 21.333  
J 1 108.000 108.000  
K 1 1.333 1.333  
L 1 75.000 75.000
```

```
$`Type II`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A   1    3.000  3.000  

B   1   27.000 27.000  

C   1   12.000 12.000  

D   1   16.333 16.333  

E   1 176.333 176.333  

F   1 133.333 133.333  

G   1    1.333  1.333  

H   1   21.333 21.333  

J   1 108.000 108.000  

K   1    1.333  1.333  

L   1   75.000 75.000
```

```
$`Type III`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A   1    3.000  3.000  

B   1   27.000 27.000  

C   1   12.000 12.000  

D   1   16.333 16.333  

E   1 176.333 176.333  

F   1 133.333 133.333  

G   1    1.333  1.333  

H   1   21.333 21.333  

J   1 108.000 108.000  

K   1    1.333  1.333  

L   1   75.000 75.000
```

(145) MODEL

```
GLM(Y ~ E*F + E*J + F*J + E*L + F*L + J*L, v2p578) # OK
```

```
$ANOVA  

Response : Y  

          Df  Sum Sq Mean Sq F value Pr(>F)  

MODEL      10  574.5  57.45   114.9 0.07249 .  

RESIDUALS     1    0.5    0.50  

CORRECTED TOTAL 11  575.0  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE Y Mean  

0.9991304 2.772968 0.7071068   25.5
```

```
$`Type I`
```

```

      Df  Sum Sq Mean Sq F value Pr(>F)
E     1 176.333 176.333 352.6667 0.03387 *
F     1 133.333 133.333 266.6667 0.03894 *
E:F   1  65.333  65.333 130.6667 0.05555 .
J     1  66.667  66.667 133.3333 0.05500 .
E:J   1   2.667   2.667   5.3333 0.26015
F:J   1 112.667 112.667 225.3333 0.04235 *
L     1 10.800 10.800 21.6000 0.13492
E:L   1   5.486   5.486 10.9714 0.18666
F:L   1   0.176   0.176   0.3516 0.65925
J:L   1   1.038   1.038   2.0769 0.38618
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)
E     1 61.633 61.633 123.2667 0.05719 .
F     1 75.208 75.208 150.4167 0.05179 .
E:F   1  9.346  9.346 18.6923 0.14470
J     1 54.675 54.675 109.3500 0.06069 .
E:J   1   0.115   0.115   0.2308 0.71490
F:J   1 72.115 72.115 144.2308 0.05289 .
L     1 10.800 10.800 21.6000 0.13492
E:L   1   5.654   5.654 11.3077 0.18402
F:L   1   0.115   0.115   0.2308 0.71490
J:L   1   1.038   1.038   2.0769 0.38618
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value Pr(>F)
E     1 61.038 61.038 122.0769 0.05746 .
F     1 61.038 61.038 122.0769 0.05746 .
E:F   1  9.346  9.346 18.6923 0.14470
J     1 61.038 61.038 122.0769 0.05746 .
E:J   1   0.115   0.115   0.2308 0.71490
F:J   1 72.115 72.115 144.2308 0.05289 .
L     1   9.346   9.346 18.6923 0.14470
E:L   1   5.654   5.654 11.3077 0.18402
F:L   1   0.115   0.115   0.2308 0.71490
J:L   1   1.038   1.038   2.0769 0.38618
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.9 Chapter 16

9.9.1 p619

(146) MODEL

```
v2p619 = read.table("C:/G/Rt/Kemp/v2p619.txt", head=TRUE)
v2p619 = af(v2p619, c("A", "B", "C"))
GLM(y ~ A + B + C + A:B, v2p619) # OK
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      4 31.429  7.8571
RESIDUALS   2  0.000  0.0000
CORRECTED TOTAL 6 31.429
```

```
$Fitness
R-square Coef Var Root MSE   y Mean
      1       0       0 10.78571
```

```
$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
A      1 13.7619 13.7619     Inf < 2.2e-16 ***
B      1  1.6667  1.6667     Inf < 2.2e-16 ***
C      1 10.0000 10.0000     Inf < 2.2e-16 ***
A:B    1  6.0000  6.0000     Inf < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
A      1    19.6    19.6     Inf < 2.2e-16 ***
B      1     3.6     3.6     Inf < 2.2e-16 ***
C      1    13.5    13.5     Inf < 2.2e-16 ***
A:B    1     6.0     6.0     Inf < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
A      1    24.0    24.0     Inf < 2.2e-16 ***
B      1     6.0     6.0     Inf < 2.2e-16 ***
C      1    13.5    13.5     Inf < 2.2e-16 ***
A:B    1     6.0     6.0     Inf < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(147) MODEL

```
GLM(y ~ A + B + C + A:C, v2p619) # OK
```

```
$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       4 26.0952  6.5238  2.4464 0.3106
RESIDUALS    2  5.3333  2.6667
CORRECTED TOTAL 6 31.4286
```

```
$Fitness
R-square Coef Var Root MSE   y Mean
0.830303 15.14033 1.632993 10.78571
```

```
$`Type I` 
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 13.7619 13.7619  5.1607 0.1511
B      1  1.6667  1.6667  0.6250 0.5120
C      1 10.0000 10.0000  3.7500 0.1924
A:C    1  0.6667  0.6667  0.2500 0.6667
```

```
$`Type II` 
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 19.6000 19.6000    7.35 0.1134
B      1  2.6667  2.6667    1.00 0.4226
C      1 10.0000 10.0000    3.75 0.1924
A:C    1  0.6667  0.6667    0.25 0.6667
```

```
$`Type III` 
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 16.6667 16.6667  6.2500 0.1296
B      1  2.6667  2.6667  1.0000 0.4226
C      1  8.1667  8.1667  3.0625 0.2222
A:C    1  0.6667  0.6667  0.2500 0.6667
```

(148) MODEL

```
GLM(y ~ A + B + C + B:C, v2p619) # OK
```

```
$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       4 26.0952  6.5238  2.4464 0.3106
RESIDUALS    2  5.3333  2.6667
CORRECTED TOTAL 6 31.4286
```

```

$Fitness
R-square Coef Var Root MSE   y Mean
0.830303 15.14033 1.632993 10.78571

$`Type I`
  Df  Sum Sq Mean Sq F value Pr(>F)
A    1 13.7619 13.7619  5.1607 0.1511
B    1  1.6667  1.6667  0.6250 0.5120
C    1 10.0000 10.0000  3.7500 0.1924
B:C  1  0.6667  0.6667  0.2500 0.6667

$`Type II`
  Df  Sum Sq Mean Sq F value Pr(>F)
A    1 16.6667 16.6667   6.25 0.1296
B    1  3.6000  3.6000   1.35 0.3652
C    1 10.0000 10.0000   3.75 0.1924
B:C  1  0.6667  0.6667   0.25 0.6667

$`Type III`
  Df  Sum Sq Mean Sq F value Pr(>F)
A    1 16.6667 16.6667  6.2500 0.1296
B    1  2.6667  2.6667  1.0000 0.4226
C    1  8.1667  8.1667  3.0625 0.2222
B:C  1  0.6667  0.6667  0.2500 0.6667

```

9.9.2 p626

(149) MODEL

```

v2p626 = read.table("C:/G/Rt/Kemp/v2p626.txt", head=TRUE)
v2p626 = af(v2p626, c("A", "B", "C"))
GLM(y ~ A + B + C + A:B, v2p626) # OK

```

```

$ANOVA
Response : y
  Df  Sum Sq Mean Sq F value Pr(>F)
MODEL          4 42.092 10.5231  22.002 0.04395 *
RESIDUALS      2  0.957  0.4783
CORRECTED TOTAL 6 43.049
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE   y Mean
0.9777801 6.217804 0.6915708 11.12243

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

A     1 16.2088 16.2088  33.890 0.02826 *  

B     1  4.8150  4.8150  10.068 0.08662 .  

C     1 15.7339 15.7339  32.898 0.02908 *  

A:B   1  5.3346  5.3346  11.154 0.07916 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

A     1 25.4131 25.4131  53.136 0.01830 *  

B     1  8.6630  8.6630  18.113 0.05102 .  

C     1 19.5193 19.5193  40.812 0.02364 *  

A:B   1  5.3346  5.3346  11.154 0.07916 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

A     1 29.7950 29.7950  62.297 0.01568 *  

B     1 11.7460 11.7460  24.559 0.03839 *  

C     1 19.5193 19.5193  40.812 0.02364 *  

A:B   1  5.3346  5.3346  11.154 0.07916 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(150) MODEL

```
GLM(y ~ A + B + C + A:C, v2p626) # OK
```

```
$ANOVA  

Response : y  

      Df  Sum Sq Mean Sq F value Pr(>F)  

MODEL        4 39.229  9.8072  5.1346 0.1696  

RESIDUALS    2  3.820  1.9100  

CORRECTED TOTAL 6 43.049
```

```
$Fitness  

R-square Coef Var Root MSE   y Mean  

0.9112627 12.42564 1.382033 11.12243
```

```
$`Type I`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

A     1 16.2088 16.2088  8.4862 0.1004  

B     1  4.8150  4.8150  2.5209 0.2533
```

```

C      1 15.7339 15.7339  8.2376 0.1030
A:C   1  2.4711  2.4711  1.2937 0.3733

$`Type II` 
  Df  Sum Sq Mean Sq F value Pr(>F)
A     1 25.4131 25.4131 13.3052 0.06762 .
B     1  6.0361  6.0361  3.1602 0.21743
C     1 15.7339 15.7339  8.2376 0.10298
A:C   1  2.4711  2.4711  1.2937 0.37327
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df  Sum Sq Mean Sq F value Pr(>F)
A     1 20.1428 20.1428 10.5459 0.08317 .
B     1  6.0361  6.0361  3.1602 0.21743
C     1 11.8863 11.8863  6.2232 0.13007
A:C   1  2.4711  2.4711  1.2937 0.37327
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(151) MODEL

```
GLM(y ~ A + B + C + B:C, v2p626) # OK
```

```
$ANOVA
Response : y
  Df  Sum Sq Mean Sq F value Pr(>F)
MODEL        4 37.340  9.3349  3.2701 0.2477
RESIDUALS    2  5.709  2.8546
CORRECTED TOTAL 6 43.049
```

```
$Fitness
R-square Coef Var Root MSE   y Mean
0.8673781 15.19055 1.689558 11.12243
```

```
$`Type I` 
  Df  Sum Sq Mean Sq F value Pr(>F)
A     1 16.2088 16.2088  5.6781 0.1400
B     1  4.8150  4.8150  1.6867 0.3236
C     1 15.7339 15.7339  5.5118 0.1434
B:C   1  0.5819  0.5819  0.2038 0.6959
```

```
$`Type II` 
  Df  Sum Sq Mean Sq F value Pr(>F)
A     1 21.9995 21.9995  7.7067 0.1090
B     1  8.6630  8.6630  3.0347 0.2236
```

```
C      1 15.7339 15.7339  5.5118 0.1434
B:C   1  0.5819  0.5819  0.2038 0.6959
```

```
$`Type III`  
Df  Sum Sq Mean Sq F value Pr(>F)  
A   1 21.9995 21.9995  7.7067 0.1090  
B   1  7.0709  7.0709  2.4770 0.2562  
C   1 13.3221 13.3221  4.6669 0.1633  
B:C  1  0.5819  0.5819  0.2038 0.6959
```

9.10 Chapter 17

9.10.1 p642

(152) MODEL

```
v2p642 = read.table("C:/G/Rt/Kemp/v2p642.txt", head=TRUE)
v2p642 = af(v2p642, 2:11)
GLM(Y ~ A + B + C + D + E + F + G, v2p642) # OK
```

```
$ANOVA
Response : Y
Df  Sum Sq Mean Sq F value Pr(>F)
MODEL          7    11.0  1.57143  1.6688 0.1646
RESIDUALS       24    22.6  0.94167
CORRECTED TOTAL 31    33.6
```

```
$Fitness
R-square Coef Var Root MSE Y Mean
0.327381 43.12867 0.9703951    2.25
```

```
$`Type I`  
Df  Sum Sq Mean Sq F value Pr(>F)
A   1 5.7800  5.7800  6.1381 0.02066 *
B   1 0.1800  0.1800  0.1912 0.66587
C   1 0.1250  0.1250  0.1327 0.71879
D   1 2.5312  2.5312  2.6881 0.11415
E   1 0.6613  0.6613  0.7022 0.41031
F   1 0.0112  0.0112  0.0119 0.91387
G   1 1.7113  1.7113  1.8173 0.19023
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  
Df  Sum Sq Mean Sq F value Pr(>F)
A   1 5.7800  5.7800  6.1381 0.02066 *
```

```

B 1 0.1800 0.1800 0.1912 0.66587
C 1 0.1250 0.1250 0.1327 0.71879
D 1 2.5312 2.5312 2.6881 0.11415
E 1 0.6613 0.6613 0.7022 0.41031
F 1 0.0112 0.0112 0.0119 0.91387
G 1 1.7113 1.7113 1.8173 0.19023
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 5.7800 5.7800 6.1381 0.02066 *  

B 1 0.1800 0.1800 0.1912 0.66587  

C 1 0.1250 0.1250 0.1327 0.71879  

D 1 2.5312 2.5312 2.6881 0.11415  

E 1 0.6613 0.6613 0.7022 0.41031  

F 1 0.0112 0.0112 0.0119 0.91387  

G 1 1.7113 1.7113 1.8173 0.19023
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(153) MODEL

```
GLM(log(S) ~ A + B + C + D + E + F + G, v2p642) # OK
```

```
$ANOVA  

Response : log(S)  

  Df Sum Sq Mean Sq F value Pr(>F)  

MODEL 7 266.43 38.062  

RESIDUALS 24 0.00 0.000  

CORRECTED TOTAL 31 266.43
```

```
$Fitness  

R-square Coef Var Root MSE log(S) Mean  

      1       0       0    -2.233358
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 1.511 1.511 Inf < 2.2e-16 ***  

B 1 0.600 0.600 Inf < 2.2e-16 ***  

C 1 0.284 0.284 Inf < 2.2e-16 ***  

D 1 0.384 0.384 Inf < 2.2e-16 ***  

E 1 0.741 0.741 Inf < 2.2e-16 ***  

F 1 261.783 261.783 Inf < 2.2e-16 ***  

G 1 1.127 1.127 Inf < 2.2e-16 ***  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

A   1    1.511   1.511      Inf < 2.2e-16 ***  

B   1    0.600   0.600      Inf < 2.2e-16 ***  

C   1    0.284   0.284      Inf < 2.2e-16 ***  

D   1    0.384   0.384      Inf < 2.2e-16 ***  

E   1    0.741   0.741      Inf < 2.2e-16 ***  

F   1 261.783 261.783      Inf < 2.2e-16 ***  

G   1    1.127   1.127      Inf < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

A   1    1.511   1.511      Inf < 2.2e-16 ***  

B   1    0.600   0.600      Inf < 2.2e-16 ***  

C   1    0.284   0.284      Inf < 2.2e-16 ***  

D   1    0.384   0.384      Inf < 2.2e-16 ***  

E   1    0.741   0.741      Inf < 2.2e-16 ***  

F   1 261.783 261.783      Inf < 2.2e-16 ***  

G   1    1.127   1.127      Inf < 2.2e-16 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.11 Chapter 19

9.11.1 p700

(154) MODEL

```
v2p700 = read.table("C:/G/Rt/Kemp/v2p700.txt", head=TRUE)
v2p700 = af(v2p700, 2:5)
GLM(Y ~ P + S + T + C, v2p700) # OK
```

```
$ANOVA
Response : Y
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       12 378.80 31.5670  57.256 0.003319 **
RESIDUALS     3   1.65  0.5513
CORRECTED TOTAL 15 380.46
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var  Root MSE  Y Mean
```

```

0.9956526 3.781124 0.7425182 19.6375

$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

P 3 53.888 17.963 32.580 0.008646 **  

S 3 154.508 51.503 93.414 0.001845 **  

T 3 149.848 49.949 90.597 0.001930 **  

C 3 20.561  6.854 12.431 0.033708 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

P 2 2.220  1.110 2.0133 0.278974  

S 3 111.966 37.322 67.6941 0.002969 **  

T 3 161.828 53.943 97.8403 0.001722 **  

C 3 20.561  6.854 12.4311 0.033708 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

CAUTION: Singularity Exists !  

  Df Sum Sq Mean Sq F value    Pr(>F)  

P 2 2.220  1.110 2.0133 0.278974  

S 3 111.966 37.322 67.6941 0.002969 **  

T 3 161.828 53.943 97.8403 0.001722 **  

C 3 20.561  6.854 12.4311 0.033708 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.11.2 p703

(155) MODEL

```

v2p703 = read.table("C:/G/Rt/Kemp/v2p703.txt", head=TRUE)
v2p703$C = ifelse(v2p703$C == 0, 4, v2p703$C)
v2p703 = af(v2p703, 2:5)
GLM(Y ~ P + S + T + C, v2p703) # OK

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      13 385.18 29.6293 21.766 0.0005673 ***
RESIDUALS   6   8.17  1.3613
CORRECTED TOTAL 19 393.35
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE Y Mean
  0.9792359 5.99551 1.166726 19.46

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
P 4 56.408 14.102 10.3596 0.0073255 **
S 3 119.260 39.753 29.2036 0.0005620 ***
T 3 190.430 63.477 46.6312 0.0001498 ***
C 3 19.083 6.361 4.6728 0.0518237 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
P 4 52.288 13.072 9.6028 0.0088641 **
S 3 167.414 55.805 40.9952 0.0002163 ***
T 3 190.430 63.477 46.6312 0.0001498 ***
C 3 19.083 6.361 4.6728 0.0518237 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
P 4 52.287 13.072 9.6028 0.0088641 **
S 3 167.414 55.805 40.9952 0.0002163 ***
T 3 190.430 63.477 46.6312 0.0001498 ***
C 3 19.083 6.361 4.6728 0.0518237 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10 Lawson - DAE with SAS

Reference

- Lawson J. Design and Analysis of Experiments with SAS. Taylor and Francis Group. 2010.

```
require(daewr)
```

10.1 Chapter 2

10.1.1 p22

(156) MODEL

```
GLM(height ~ time, bread) # OK
```

```
$ANOVA
Response : height
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      2 21.573 10.7865 4.6022 0.042 *
RESIDUALS   9 21.094  2.3438
CORRECTED TOTAL 11 42.667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE height Mean
0.5056152 20.87633 1.530931    7.333333

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 21.573 10.787 4.6022 0.042 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 21.573 10.787 4.6022 0.042 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 21.573 10.787 4.6022 0.042 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.1.2 p32

(157) MODEL

```
GLM(height^(1 - 1.294869) ~ time, bread) # OK

$ANOVA
Response : height^(1 - 1.294869)
            Df   Sum Sq  Mean Sq F value Pr(>F)
MODEL          2 0.0130560 0.0065280 5.9356 0.02271 *
RESIDUALS      9 0.0098983 0.0010998
CORRECTED TOTAL 11 0.0229544
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE height^(1 - 1.294869) Mean
0.5687825 5.890685 0.03316344           0.5629811

$`Type I` 
Df   Sum Sq  Mean Sq F value Pr(>F)
time 2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
Df   Sum Sq  Mean Sq F value Pr(>F)
time 2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
Df   Sum Sq  Mean Sq F value Pr(>F)
time 2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.1.3 p42

(158) MODEL

```
GLM(yield ~ treat, sugarbeet) # OK

$ANOVA
Response : yield
            Df   Sum Sq  Mean Sq F value     Pr(>F)
```

```

MODEL           3 291.00  97.002    45.9 1.718e-07 ***
RESIDUALS      14 29.59   2.113
CORRECTED TOTAL 17 320.59
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
R-square Coef Var Root MSE yield Mean
0.9077128 3.182182 1.453727 45.68333

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291  97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291  97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291  97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2 Chapter 3

10.2.1 p63

(159) MODEL

```
GLM(CO ~ Eth + Ratio + Eth:Ratio, C0data) # OK
```

```

$ANOVA
Response : CO
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          8 1654.0 206.750  40.016 3.861e-06 ***
RESIDUALS      9   46.5   5.167
CORRECTED TOTAL 17 1700.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness

```

```

R-square Coef Var Root MSE CO Mean
0.9726551 3.120865 2.27303 72.83333

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Eth       2     324   162.0  31.355 8.790e-05 ***
Ratio     2     652   326.0  63.097 5.067e-06 ***
Eth:Ratio 4     678   169.5  32.806 2.240e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Eth       2     324   162.0  31.355 8.790e-05 ***
Ratio     2     652   326.0  63.097 5.067e-06 ***
Eth:Ratio 4     678   169.5  32.806 2.240e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Eth       2     324   162.0  31.355 8.790e-05 ***
Ratio     2     652   326.0  63.097 5.067e-06 ***
Eth:Ratio 4     678   169.5  32.806 2.240e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(160) MODEL

```
GLM(CO ~ Ratio + Eth + Ratio:Eth, COdata) # OK
```

```

$ANOVA
Response : CO
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      8 1654.0 206.750  40.016 3.861e-06 ***
RESIDUALS   9   46.5   5.167
CORRECTED TOTAL 17 1700.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
R-square Coef Var Root MSE CO Mean
0.9726551 3.120865 2.27303 72.83333

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Ratio      2     652   326.0  63.097 5.067e-06 ***

```

```

Eth          2     324    162.0   31.355 8.790e-05 ***
Ratio:Eth   4     678    169.5   32.806 2.240e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Ratio      2     652    326.0   63.097 5.067e-06 ***  

Eth        2     324    162.0   31.355 8.790e-05 ***  

Ratio:Eth  4     678    169.5   32.806 2.240e-05 ***  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Ratio      2     652    326.0   63.097 5.067e-06 ***  

Eth        2     324    162.0   31.355 8.790e-05 ***  

Ratio:Eth  4     678    169.5   32.806 2.240e-05 ***  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2.2 p74

(161) MODEL

```

GLM(CO ~ Eth + Ratio + Eth:Ratio, C0data[-18,]) # OK

$ANOVA
Response : CO
      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      8 1423.0 177.879   31.978 2.749e-05 ***  

RESIDUALS   8    44.5   5.563  

CORRECTED TOTAL 16 1467.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  CO Mean
0.9696769 3.199874 2.358495 73.70588

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth        2 472.66 236.33   42.486 5.482e-05 ***  

Ratio      2 395.33 197.66   35.535 0.0001048 ***  

Eth:Ratio  4 555.04 138.76   24.945 0.0001427 ***  

---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth       2 398.26 199.13 35.799 0.0001020 ***  

Ratio     2 395.33 197.66 35.535 0.0001048 ***  

Eth:Ratio 4 555.04 138.76 24.945 0.0001427 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth       2 319.45 159.73 28.715 0.0002235 ***  

Ratio     2 511.45 255.73 45.973 4.105e-05 ***  

Eth:Ratio 4 555.04 138.76 24.945 0.0001427 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2.3 p91

(162) MODEL

```

volt$XA = (as.numeric(as.character(volt$A)) - 27)/5
volt$XB = (as.numeric(as.character(volt$B)) - 2.75)/2.25
volt$XC = (as.numeric(as.character(volt$C)) - 2.75)/2.25
GLM(y ~ XA + XB + XC + XA:XB + XA:XC + XB:XC + XA:XB:XC, volt) # OK

```

```

$ANOVA  

Response : y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      7 8843.4 1263.35 3.8686 0.0385 *  

RESIDUALS   8 2612.5 326.56  

CORRECTED TOTAL 15 11455.9  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  R-square Coef Var Root MSE   y Mean  

0.7719523 2.702969 18.07104 668.5625

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

XA        1 4522.6 4522.6 13.8490 0.005859 **  

XB        1   14.1    14.1  0.0431 0.840793  

XC        1  473.1   473.1  1.4486 0.263154  

XA:XB     1  715.6   715.6  2.1912 0.177071

```

```

XA:XC      1 2525.1  2525.1  7.7322 0.023899 *
XB:XC      1   52.6    52.6  0.1610 0.698780
XA:XB:XC  1   540.6   540.6  1.6553 0.234218
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value Pr(>F)
XA      1 4522.6 4522.6 13.8490 0.005859 **
XB      1   14.1   14.1  0.0431 0.840793
XC      1   473.1   473.1  1.4486 0.263154
XA:XB    1   715.6   715.6  2.1912 0.177071
XA:XC    1 2525.1  2525.1  7.7322 0.023899 *
XB:XC    1   52.6    52.6  0.1610 0.698780
XA:XB:XC 1   540.6   540.6  1.6553 0.234218
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value Pr(>F)
XA      1 4522.6 4522.6 13.8490 0.005859 **
XB      1   14.1   14.1  0.0431 0.840793
XC      1   473.1   473.1  1.4486 0.263154
XA:XB    1   715.6   715.6  2.1912 0.177071
XA:XC    1 2525.1  2525.1  7.7322 0.023899 *
XB:XC    1   52.6    52.6  0.1610 0.698780
XA:XB:XC 1   540.6   540.6  1.6553 0.234218
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2.4 p97

(163) MODEL

```

chem2 = af(chem, c("A","B","C","D"))
GLM(y ~ A*B*C*D, chem2) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      15 6369.4  424.63
RESIDUALS   0    0.0
CORRECTED TOTAL 15 6369.4

$Fitness
R-square Coef Var Root MSE  y Mean

```

1 NA NA 62.3125

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	637.6	637.6		
B	1	5076.6	5076.6		
A:B	1	451.6	451.6		
C	1	0.6	0.6		
A:C	1	10.6	10.6		
B:C	1	1.6	1.6		
A:B:C	1	0.6	0.6		
D	1	7.6	7.6		
A:D	1	68.1	68.1		
B:D	1	0.1	0.1		
A:B:D	1	7.6	7.6		
C:D	1	7.6	7.6		
A:C:D	1	95.1	95.1		
B:C:D	1	3.1	3.1		
A:B:C:D	1	1.6	1.6		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	637.6	637.6		
B	1	5076.6	5076.6		
A:B	1	451.6	451.6		
C	1	0.6	0.6		
A:C	1	10.6	10.6		
B:C	1	1.6	1.6		
A:B:C	1	0.6	0.6		
D	1	7.6	7.6		
A:D	1	68.1	68.1		
B:D	1	0.1	0.1		
A:B:D	1	7.6	7.6		
C:D	1	7.6	7.6		
A:C:D	1	95.1	95.1		
B:C:D	1	3.1	3.1		
A:B:C:D	1	1.6	1.6		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	637.6	637.6		
B	1	5076.6	5076.6		
A:B	1	451.6	451.6		
C	1	0.6	0.6		
A:C	1	10.6	10.6		
B:C	1	1.6	1.6		
A:B:C	1	0.6	0.6		
D	1	7.6	7.6		

A:D	1	68.1	68.1
B:D	1	0.1	0.1
A:B:D	1	7.6	7.6
C:D	1	7.6	7.6
A:C:D	1	95.1	95.1
B:C:D	1	3.1	3.1
A:B:C:D	1	1.6	1.6

10.2.5 p104

(164) MODEL

```
GLM(y ~ A*B*C*D, BoxM) # OK
```

\$ANOVA

```
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      15 207.1 13.807
RESIDUALS   0    0.0
CORRECTED TOTAL 15 207.1
```

\$Fitness

```
R-square Coef Var Root MSE y Mean
      1       NA       NA 48.245
```

\$`Type I`

```
Df Sum Sq Mean Sq F value Pr(>F)
A      1 2.560  2.560
B      1 71.234 71.234
A:B    1 3.312  3.312
C      1 55.056 55.056
A:C    1 24.800 24.800
B:C    1 2.560  2.560
A:B:C  1 5.760  5.760
D      1 4.080  4.080
A:D    1 1.346  1.346
B:D    1 5.570  5.570
A:B:D  1 2.074  2.074
C:D    1 8.880  8.880
A:C:D  1 0.640  0.640
B:C:D  1 9.986  9.986
A:B:C:D 1 9.242  9.242
```

\$`Type II`

```
Df Sum Sq Mean Sq F value Pr(>F)
A      1 2.560  2.560
```

```

B           1 71.234 71.234
A:B         1 3.312  3.312
C           1 55.056 55.056
A:C         1 24.800 24.800
B:C         1 2.560  2.560
A:B:C      1 5.760  5.760
D           1 4.080  4.080
A:D         1 1.346  1.346
B:D         1 5.570  5.570
A:B:D      1 2.074  2.074
C:D         1 8.880  8.880
A:C:D      1 0.640  0.640
B:C:D      1 9.986  9.986
A:B:C:D    1 9.242  9.242

```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	2.560	2.560		
B	1	71.234	71.234		
A:B	1	3.312	3.312		
C	1	55.056	55.056		
A:C	1	24.800	24.800		
B:C	1	2.560	2.560		
A:B:C	1	5.760	5.760		
D	1	4.080	4.080		
A:D	1	1.346	1.346		
B:D	1	5.570	5.570		
A:B:D	1	2.074	2.074		
C:D	1	8.880	8.880		
A:C:D	1	0.640	0.640		
B:C:D	1	9.986	9.986		
A:B:C:D	1	9.242	9.242		

10.3 Chapter 4

10.3.1 p122

(165) MODEL

```
GLM(rate ~ rat + dose, drug) # OK
```

```

$ANOVA
Response : rate
              Df  Sum Sq  Mean Sq F value   Pr(>F)
MODEL          13 2.12867 0.163744 19.613 1.59e-12 ***
RESIDUALS     36 0.30055 0.008349

```

```

CORRECTED TOTAL 49 2.42922
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE rate Mean
0.8762762 9.994644 0.09137104    0.9142

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
rat   9 1.66846 0.18538 22.205 3.749e-12 ***
dose  4 0.46021 0.11505 13.781 6.535e-07 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
rat   9 1.66846 0.18538 22.205 3.749e-12 ***
dose  4 0.46021 0.11505 13.781 6.535e-07 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
rat   9 1.66846 0.18538 22.205 3.749e-12 ***
dose  4 0.46021 0.11505 13.781 6.535e-07 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.3.2 p127

(166) MODEL

```
GLM(y ~ block + treat + strain + treat:strain, bha) # OK
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      8 543.22 67.902 26.203 0.0001507 ***
RESIDUALS  7 18.14  2.591
CORRECTED TOTAL 15 561.36
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  y Mean

```

```

0.9676855 12.39493 1.609791 12.9875

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block       1  47.61   47.61  18.3721  0.003627 **  

treat       1 422.30  422.30 162.9613 4.194e-06 ***  

strain      3  32.96   10.99   4.2399  0.052741 .  

treat:strain 3  40.34   13.45   5.1892  0.033685 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block       1  47.61   47.61  18.3721  0.003627 **  

treat       1 422.30  422.30 162.9613 4.194e-06 ***  

strain      3  32.96   10.99   4.2399  0.052741 .  

treat:strain 3  40.34   13.45   5.1892  0.033685 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block       1  47.61   47.61  18.3721  0.003627 **  

treat       1 422.30  422.30 162.9613 4.194e-06 ***  

strain      3  32.96   10.99   4.2399  0.052741 .  

treat:strain 3  40.34   13.45   5.1892  0.033685 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.3.3 p129

(167) MODEL

```
GLM(cdistance ~ id + teehgt, rcb) # OK
```

```
$ANOVA  

Response : cdistance  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      10 126465 12646.5  161.72 < 2.2e-16 ***  

RESIDUALS  124  9697   78.2  

CORRECTED TOTAL 134 136162  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness  

R-square Coef Var Root MSE cdistance Mean
```

```

0.9287846 5.013727 8.8431 176.3778

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

id      8 124741   15593 199.394 < 2.2e-16 ***  

teehgt  2   1724     862 11.023 3.926e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

id      8 124741   15593 199.394 < 2.2e-16 ***  

teehgt  2   1724     862 11.023 3.926e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

id      8 124741   15593 199.394 < 2.2e-16 ***  

teehgt  2   1724     862 11.023 3.926e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.3.4 p136

(168) MODEL

```
GLM(AUC ~ Subject + Period + Treat, bioequiv) # OK
```

```
$ANOVA  

Response : AUC  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL       6 174461   29077  0.1315 0.9774  

RESIDUALS   2 442158   221079  

CORRECTED TOTAL 8 616618
```

```
$Fitness  

R-square Coef Var Root MSE AUC Mean  

0.2829314 41.18855 470.1902 1141.556
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Subject    2 114264   57132  0.2584 0.7946  

Period     2  45196   22598  0.1022 0.9073  

Treat      2 15000    7500  0.0339 0.9672
```

```
$`Type II`  
      Df Sum Sq Mean Sq F value Pr(>F)  
Subject  2 114264   57132  0.2584 0.7946  
Period   2  45196   22598  0.1022 0.9073  
Treat    2 15000    7500  0.0339 0.9672
```

```
$`Type III`  
      Df Sum Sq Mean Sq F value Pr(>F)  
Subject  2 114264   57132  0.2584 0.7946  
Period   2  45196   22598  0.1022 0.9073  
Treat    2 15000    7500  0.0339 0.9672
```

10.4 Chapter 5

10.4.1 p152

(169) MODEL

```
GLM(conc ~ lab, Apo) # OK
```

```
$ANOVA  
Response : conc  
      Df Sum Sq Mean Sq F value Pr(>F)  
MODEL       3 0.092233 0.0307444 42.107 4.009e-10 ***  
RESIDUALS   26 0.018984 0.0007302  
CORRECTED TOTAL 29 0.111217  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  
R-square Coef Var Root MSE conc Mean  
0.8293064 2.367047 0.02702142 1.141567
```

```
$`Type I`  
      Df Sum Sq Mean Sq F value Pr(>F)  
lab  3 0.092233 0.030744 42.107 4.009e-10 ***  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  
      Df Sum Sq Mean Sq F value Pr(>F)  
lab  3 0.092233 0.030744 42.107 4.009e-10 ***  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
```

```

      Df   Sum Sq  Mean Sq F value    Pr(>F)
lab  3 0.092233 0.030744  42.107 4.009e-10 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.4.2 p181

(170) MODEL

```
GLM(residue ~ form + tech + form:tech + plot:form:tech, pesticide) # OK
```

```

$ANOVA
Response : residue
      Df   Sum Sq  Mean Sq F value    Pr(>F)
MODEL      7 0.036857 0.0052653 11.804 0.001187 **
RESIDUALS  8 0.003569 0.0004461
CORRECTED TOTAL 15 0.040426
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var  Root MSE residue Mean
0.9117275 6.671729 0.02112019     0.3165625

```

```

$`Type I`
      Df   Sum Sq  Mean Sq F value    Pr(>F)
form       1 0.000018 0.000018 0.0405  0.84554
tech       1 0.032310 0.032310 72.4339 2.789e-05 ***
form:tech  1 0.002186 0.002186 4.8997  0.05776 .
form:tech:plot 4 0.002344 0.000586 1.3136  0.34317
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df   Sum Sq  Mean Sq F value    Pr(>F)
form       1 0.000018 0.000018 0.0405  0.84554
tech       1 0.032310 0.032310 72.4339 2.789e-05 ***
form:tech  1 0.002186 0.002186 4.8997  0.05776 .
form:tech:plot 4 0.002344 0.000586 1.3136  0.34317
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
      Df   Sum Sq  Mean Sq F value    Pr(>F)
form       1 0.000018 0.000018 0.0405  0.84554
tech       1 0.032310 0.032310 72.4339 2.789e-05 ***

```

```

form:tech      1 0.002186 0.002186  4.8997   0.05776 .
form:tech:plot 4 0.002344 0.000586  1.3136   0.34317
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.5 Chapter 7

10.5.1 p260

(171) MODEL

```
GLM(score ~ recipe + panelist, taste) # OK
```

```

$ANOVA
Response : score
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      14 28.458 2.03274  2.661 0.0719 .
RESIDUALS   9  6.875 0.76389
CORRECTED TOTAL 23 35.333
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  R-square Coef Var Root MSE score Mean
0.8054245 14.98298 0.8740074    5.833333

```

```

$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
recipe     3 21.0000  7.000  9.1636 0.004246 **
panelist  11  7.4583  0.678  0.8876 0.581099
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
recipe     3 9.1250 3.04167  3.9818 0.04649 *
panelist  11 7.4583 0.67803  0.8876 0.58110
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
recipe     3 9.1250 3.04167  3.9818 0.04649 *
panelist  11 7.4583 0.67803  0.8876 0.58110
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.5.2 p262

(172) MODEL

```
GLM(pressure ~ Block + Treatment, BPmonitor) # OK
```

```
$ANOVA
Response : pressure
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       8 321.00 40.125 4.4174 0.1245
RESIDUALS    3  27.25  9.083
CORRECTED TOTAL 11 348.25

$Fitness
  R-square Coef Var Root MSE pressure Mean
0.9217516 3.876343 3.013857          77.75

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Block       5 73.75 14.750 1.6239 0.36606
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Block       5 83.25 16.650 1.8330 0.32772
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Block       5 83.25 16.650 1.8330 0.32772
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.5.3 p276

(173) MODEL

```
GLM(weight ~ Blocks + A + B + C + D + E + F + G + H, Bff) # OK
```

```
$ANOVA
Response : weight
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	15	158.37	10.558		
RESIDUALS	0	0.00			
CORRECTED TOTAL	15	158.37			

\$Fitness

R-square	Coef	Var	Root	MSE	weight	Mean
1	NA	NA	5.925625			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Blocks	7	30.567	4.367		
A	1	21.879	21.879		
B	1	8.338	8.338		
C	1	6.213	6.213		
D	1	12.870	12.870		
E	1	0.098	0.098		
F	1	1.260	1.260		
G	1	71.868	71.868		
H	1	5.279	5.279		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Blocks	7	30.567	4.367		
A	1	21.879	21.879		
B	1	8.338	8.338		
C	1	6.213	6.213		
D	1	12.870	12.870		
E	1	0.098	0.098		
F	1	1.260	1.260		
G	1	71.868	71.868		
H	1	5.279	5.279		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Blocks	7	30.567	4.367		
A	1	21.879	21.879		
B	1	8.338	8.338		
C	1	6.213	6.213		
D	1	12.870	12.870		
E	1	0.098	0.098		
F	1	1.260	1.260		
G	1	71.868	71.868		
H	1	5.279	5.279		

10.6 Chapter 8

10.6.1 p315

(174) MODEL

```
GLM(ys ~ Block + A*B + Block:A:B + C*D + A:C + A:D + B:C + B:D + A:B:C + A:B:D +
    A:C:D + B:C:D + A:B:C:D, sausage) # OK
```

\$ANOVA

```
Response : ys
            Df   Sum Sq   Mean Sq F value Pr(>F)
MODEL          19 0.064059 0.0033715 14.134 1.74e-05 ***
RESIDUALS      12 0.002862 0.0002385
CORRECTED TOTAL 31 0.066922
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
R-square Coef Var Root MSE ys Mean
0.9572262 0.7632948 0.01544479 2.023438
```

\$`Type I`

```
Df   Sum Sq   Mean Sq F value Pr(>F)
Block 1 0.000903 0.000903 3.7860 0.075482 .
A     1 0.045753 0.045753 191.8035 9.647e-09 ***
B     1 0.002628 0.002628 11.0175 0.006119 **
A:B   1 0.001128 0.001128 4.7293 0.050371 .
Block:A:B 3 0.005484 0.001828 7.6638 0.004007 **
C     1 0.003828 0.003828 16.0480 0.001743 **
D     1 0.000528 0.000528 2.2140 0.162566
C:D   1 0.000253 0.000253 1.0611 0.323272
A:C   1 0.000153 0.000153 0.6419 0.438593
A:D   1 0.000903 0.000903 3.7860 0.075482 .
B:C   1 0.000078 0.000078 0.3275 0.577693
B:D   1 0.000253 0.000253 1.0611 0.323272
A:B:C 1 0.001378 0.001378 5.7773 0.033299 *
A:B:D 1 0.000703 0.000703 2.9476 0.111680
A:C:D 1 0.000028 0.000028 0.1179 0.737260
B:C:D 1 0.000028 0.000028 0.1179 0.737260
A:B:C:D 1 0.000028 0.000028 0.1179 0.737260
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
Df   Sum Sq   Mean Sq F value Pr(>F)
Block 1 0.000903 0.000903 3.7860 0.075482 .
```

```

A      1 0.045753 0.045753 191.8035 9.647e-09 ***
B      1 0.002628 0.002628  11.0175  0.006119 **
A:B    1 0.001128 0.001128   4.7293  0.050371 .
Block:A:B 3 0.005484 0.001828   7.6638  0.004007 **
C      1 0.003828 0.003828  16.0480  0.001743 **
D      1 0.000528 0.000528   2.2140  0.162566
C:D    1 0.000253 0.000253   1.0611  0.323272
A:C    1 0.000153 0.000153   0.6419  0.438593
A:D    1 0.000903 0.000903   3.7860  0.075482 .
B:C    1 0.000078 0.000078   0.3275  0.577693
B:D    1 0.000253 0.000253   1.0611  0.323272
A:B:C  1 0.001378 0.001378   5.7773  0.033299 *
A:B:D  1 0.000703 0.000703   2.9476  0.111680
A:C:D  1 0.000028 0.000028   0.1179  0.737260
B:C:D  1 0.000028 0.000028   0.1179  0.737260
A:B:C:D 1 0.000028 0.000028   0.1179  0.737260
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq  Mean Sq  F value    Pr(>F)
Block   1 0.000903 0.000903   3.7860  0.075482 .
A       1 0.045753 0.045753 191.8035 9.647e-09 ***
B       1 0.002628 0.002628  11.0175  0.006119 **
A:B     1 0.001128 0.001128   4.7293  0.050371 .
Block:A:B 3 0.005484 0.001828   7.6638  0.004007 **
C       1 0.003828 0.003828  16.0480  0.001743 **
D       1 0.000528 0.000528   2.2140  0.162566
C:D    1 0.000253 0.000253   1.0611  0.323272
A:C    1 0.000153 0.000153   0.6419  0.438593
A:D    1 0.000903 0.000903   3.7860  0.075482 .
B:C    1 0.000078 0.000078   0.3275  0.577693
B:D    1 0.000253 0.000253   1.0611  0.323272
A:B:C  1 0.001378 0.001378   5.7773  0.033299 *
A:B:D  1 0.000703 0.000703   2.9476  0.111680
A:C:D  1 0.000028 0.000028   0.1179  0.737260
B:C:D  1 0.000028 0.000028   0.1179  0.737260
A:B:C:D 1 0.000028 0.000028   0.1179  0.737260
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.6.2 p320

(175) MODEL

```
GLM(y ~ A*B*C*D*E, plasma) # OK
```

\$ANOVA

```
Response : y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          31 6672.9 215.26
RESIDUALS      0    0.0
CORRECTED TOTAL 31 6672.9
```

\$Fitness

```
R-square Coef Var Root MSE   y Mean
1           NA       NA 40.98125
```

\$`Type I`

```
Df Sum Sq Mean Sq F value Pr(>F)
A          1 1118.65 1118.65
B          1 142.81 142.81
A:B        1 141.96 141.96
C          1  91.80  91.80
A:C        1  70.81  70.81
B:C        1    5.78    5.78
A:B:C     1   65.55   65.55
D          1 1824.08 1824.08
A:D        1 2194.53 2194.53
B:D        1   87.78   87.78
A:B:D     1   87.12   87.12
C:D        1   22.45   22.45
A:C:D     1   42.78   42.78
B:C:D     1   12.25   12.25
A:B:C:D   1  375.38  375.38
E          1   78.75   78.75
A:E        1 278.48 278.48
B:E        1    0.72    0.72
A:B:E     1    0.10    0.10
C:E        1    0.15    0.15
A:C:E     1    0.24    0.24
B:C:E     1    6.48    6.48
A:B:C:E   1    1.53    1.53
D:E        1    8.40    8.40
A:D:E     1    5.28    5.28
B:D:E     1    0.28    0.28
A:B:D:E   1    0.60    0.60
C:D:E     1    0.85    0.85
A:C:D:E   1    0.55    0.55
B:C:D:E   1    6.30    6.30
A:B:C:D:E 1    0.50    0.50
```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	1118.65	1118.65		
B	1	142.81	142.81		
A:B	1	141.96	141.96		
C	1	91.80	91.80		
A:C	1	70.81	70.81		
B:C	1	5.78	5.78		
A:B:C	1	65.55	65.55		
D	1	1824.08	1824.08		
A:D	1	2194.53	2194.53		
B:D	1	87.78	87.78		
A:B:D	1	87.12	87.12		
C:D	1	22.45	22.45		
A:C:D	1	42.78	42.78		
B:C:D	1	12.25	12.25		
A:B:C:D	1	375.38	375.38		
E	1	78.75	78.75		
A:E	1	278.48	278.48		
B:E	1	0.72	0.72		
A:B:E	1	0.10	0.10		
C:E	1	0.15	0.15		
A:C:E	1	0.24	0.24		
B:C:E	1	6.48	6.48		
A:B:C:E	1	1.53	1.53		
D:E	1	8.40	8.40		
A:D:E	1	5.28	5.28		
B:D:E	1	0.28	0.28		
A:B:D:E	1	0.60	0.60		
C:D:E	1	0.85	0.85		
A:C:D:E	1	0.55	0.55		
B:C:D:E	1	6.30	6.30		
A:B:C:D:E	1	0.50	0.50		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	1118.64	1118.64		
B	1	142.80	142.80		
A:B	1	141.96	141.96		
C	1	91.80	91.80		
A:C	1	70.81	70.81		
B:C	1	5.78	5.78		
A:B:C	1	65.55	65.55		
D	1	1824.08	1824.08		
A:D	1	2194.53	2194.53		
B:D	1	87.78	87.78		
A:B:D	1	87.12	87.12		
C:D	1	22.45	22.45		

A:C:D	1	42.78	42.78
B:C:D	1	12.25	12.25
A:B:C:D	1	375.38	375.38
E	1	78.75	78.75
A:E	1	278.48	278.48
B:E	1	0.72	0.72
A:B:E	1	0.10	0.10
C:E	1	0.15	0.15
A:C:E	1	0.24	0.24
B:C:E	1	6.48	6.48
A:B:C:E	1	1.53	1.53
D:E	1	8.40	8.40
A:D:E	1	5.28	5.28
B:D:E	1	0.28	0.28
A:B:D:E	1	0.60	0.60
C:D:E	1	0.85	0.85
A:C:D:E	1	0.55	0.55
B:C:D:E	1	6.30	6.30
A:B:C:D:E	1	0.50	0.50

10.6.3 p335

(176) MODEL

```
gear$A = as.numeric(as.character(gear$A))
gear$B = as.numeric(as.character(gear$B))
gear$C = as.numeric(as.character(gear$C))
gear$P = as.numeric(as.character(gear$P))
gear$Q = as.numeric(as.character(gear$Q))
REG(y ~ A*B*C + P + Q + A:P + A:Q + B:P + B:Q + C:P + C:Q, gear) # OK
```

	Estimate	Std. Error	Df	t value	Pr(> t)
(Intercept)	15.4062		0		
A	-4.9062		0		
B	-0.1562		0		
A:B	0.5312		0		
C	3.9688		0		
A:C	2.9062		0		
B:C	0.4062		0		
A:B:C	0.5938		0		
P	-2.3438		0		
Q	-3.4062		0		
A:P	-0.9062		0		
A:Q	-0.3438		0		
B:P	1.0938		0		
B:Q	0.1562		0		

C:P	-0.2812	0
C:Q	0.7812	0

10.7 Chapter 9

10.7.1 p349

(177) MODEL

```
GLM(pl ~ Subject + Period + Treat, antifungal) # OK
```

```
$ANOVA
Response : pl
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      18 118.558  6.5866 1.4435 0.2388
RESIDUALS   15  68.444  4.5630
CORRECTED TOTAL 33 187.002
```

```
$Fitness
R-square Coef Var Root MSE  pl Mean
0.6339915 16.23328 2.136109 13.15882
```

```
$`Type I`
          Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1    0.922  0.9224  0.2021 0.6594
Treat     1    2.993  2.9932  0.6560 0.4306
```

```
$`Type II`
          Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1    0.734  0.7344  0.1609 0.6939
Treat     1    2.993  2.9932  0.6560 0.4306
```

```
$`Type III`
          Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1    0.734  0.7344  0.1609 0.6939
Treat     1    2.993  2.9932  0.6560 0.4306
```

10.7.2 p355

(178) MODEL

```
GLM(y ~ Group + Subject:Group + Period + Treat + Carry, bioequiv) # OK
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	39	417852	10714.1	20.367	< 2.2e-16 ***
RESIDUALS	68	35772	526.1		
CORRECTED TOTAL	107	453624			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	y	Mean
0.9211408	22.62287	22.93611	101.3846			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Group	1	43335	43335	82.3763	2.46e-13 ***
Group:Subject	34	370970	10911	20.7406	< 2.2e-16 ***
Period	2	287	143	0.2723	0.7624
Treat	1	2209	2209	4.1993	0.0443 *
Carry	1	1051	1051	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Group	1	32616	32616	61.9998	3.712e-11 ***
Group:Subject	34	370970	10911	20.7406	< 2.2e-16 ***
Period	1	38	38	0.0724	0.7888
Treat	1	2209	2209	4.1993	0.0443 *
Carry	1	1051	1051	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Group	1	32616	32616	61.9998	3.712e-11 ***
Group:Subject	34	370970	10911	20.7406	< 2.2e-16 ***
Period	1	38	38	0.0724	0.7888
Treat	1	2209	2209	4.1993	0.0443 *
Carry	1	1051	1051	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(179) MODEL

```
GLM(y ~ Subject + Period + Treat + Carry, bioequiv) # OK
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	39	417852	10714.1	20.367	< 2.2e-16 ***
RESIDUALS	68	35772	526.1		
CORRECTED TOTAL	107	453624			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root	MSE	y	Mean
0.9211408	22.62287	22.93611	101.3846			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	35	414306	11837.3	22.5016	<2e-16 ***
Period	2	287	143.3	0.2723	0.7624
Treat	1	2209	2209.1	4.1993	0.0443 *
Carry	1	1051	1050.6	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	35	403586	11531.0	21.9194	<2e-16 ***
Period	1	38	38.1	0.0724	0.7888
Treat	1	2209	2209.1	4.1993	0.0443 *
Carry	1	1051	1050.6	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	35	403586	11531.0	21.9194	<2e-16 ***
Period	1	38	38.1	0.0724	0.7888
Treat	1	2209	2209.1	4.1993	0.0443 *
Carry	1	1051	1050.6	1.9970	0.1622

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10.7.3 p361

(180) MODEL

```
GLM(Time ~ Subject + Period + Treat + Carry, chipman) # OK
```

\$ANOVA

Response : Time

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	17	28.0757	1.65151	64.421	1.139e-12 ***
RESIDUALS	18	0.4615	0.02564		
CORRECTED TOTAL	35	28.5372			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root MSE	Time Mean
0.9838299	2.561577	0.1601128	6.250556	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	11	24.2084	2.20076	85.8462	3.157e-13 ***
Period	2	3.2065	1.60325	62.5388	7.894e-09 ***
Treat	2	0.4276	0.21382	8.3406	0.002733 **
Carry	2	0.2332	0.11660	4.5484	0.025188 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	11	24.2547	2.20497	86.0105	3.104e-13 ***
Period	1	0.0018	0.00184	0.0717	0.7919554
Treat	2	0.6392	0.31958	12.4661	0.0004003 ***
Carry	2	0.2332	0.11660	4.5484	0.0251881 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Subject	11	24.2547	2.20497	86.0105	3.104e-13 ***
Period	1	0.0018	0.00184	0.0717	0.7919554
Treat	2	0.6392	0.31958	12.4661	0.0004003 ***
Carry	2	0.2332	0.11660	4.5484	0.0251881 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10.7.4 p372

(181) MODEL

```

residue$lc1 = log(residue$X1)
residue$lc2 = log(residue$X2)
residue$lc3 = log(residue$X3)
residue$lc4 = log(residue$X4)
residue$lc5 = log(residue$X5)
residue$sp = 7*residue$lc2+ 14*residue$lc3 + 30*residue$lc4 + 60*residue$lc5
residue$sm = residue$lc1 + residue$lc2+ residue$lc3 + residue$lc4 + residue$lc5
residue$num = 5*residue$sp - 111*residue$sm
residue$den = 5*4745 - 111^2
residue$k = residue$num/residue$den
residue$HL = -log(2)/residue$k
residue$logHL = log(residue$HL)
GLM(logHL ~ temp*moisture*soil, residue) # OK

```

\$ANOVA

Response : logHL

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	7	7.5133	1.07332	13.543	0.0007329 ***
RESIDUALS	8	0.6340	0.07925		
CORRECTED TOTAL	15	8.1473			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

R-square	Coef	Var	Root MSE	logHL	Mean
0.9221806	5.774532	0.2815174	4.875155		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
temp	1	6.0503	6.0503	76.3427	2.303e-05 ***
moisture	1	0.9521	0.9521	12.0134	0.008492 **
temp:moisture	1	0.0013	0.0013	0.0162	0.901779
soil	1	0.4098	0.4098	5.1712	0.052559 .
temp:soil	1	0.0086	0.0086	0.1081	0.750753
moisture:soil	1	0.0860	0.0860	1.0855	0.327921
temp:moisture:soil	1	0.0051	0.0051	0.0648	0.805427

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
temp	1	6.0503	6.0503	76.3427	2.303e-05 ***
moisture	1	0.9521	0.9521	12.0134	0.008492 **
temp:moisture	1	0.0013	0.0013	0.0162	0.901779
soil	1	0.4098	0.4098	5.1712	0.052559 .
temp:soil	1	0.0086	0.0086	0.1081	0.750753
moisture:soil	1	0.0860	0.0860	1.0855	0.327921

```

temp:moisture:soil 1 0.0051 0.0051 0.0648 0.805427
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

          Df Sum Sq Mean Sq F value    Pr(>F)  

temp           1 6.0503 6.0503 76.3427 2.303e-05 ***  

moisture       1 0.9521 0.9521 12.0134 0.008492 **  

temp:moisture  1 0.0013 0.0013 0.0162 0.901779  

soil           1 0.4098 0.4098 5.1712 0.052559 .  

temp:soil      1 0.0086 0.0086 0.1081 0.750753  

moisture:soil  1 0.0860 0.0860 1.0855 0.327921  

temp:moisture:soil 1 0.0051 0.0051 0.0648 0.805427
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.8 Chapter 11

10.8.1 p461

(182) MODEL

```

GLM(y ~ x1 + x2 + x1:x2 + x1:x3 + x2:x3, pest) # OK

$ANOVA
Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      5 275.642 55.128 160.38 4.631e-07 ***  

RESIDUALS   7  2.406  0.344  

CORRECTED TOTAL 12 278.048
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE  y Mean
0.9913463 1.113968 0.5862902 52.63077

$`Type I`  

          Df Sum Sq Mean Sq F value    Pr(>F)  

x1        1 83.402 83.402 242.6351 1.086e-06 ***  

x2        1 161.734 161.734 470.5191 1.116e-07 ***  

x1:x2    1  0.246  0.246  0.7169 0.4251627  

x1:x3    1 15.663 15.663 45.5660 0.0002649 ***  

x2:x3    1 14.596 14.596 42.4614 0.0003291 ***  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1  215.951 215.951 628.246 4.105e-08 ***  

x2     1  175.256 175.256 509.855 8.458e-08 ***  

x1:x2  1   0.025   0.025   0.072 0.7961658  

x1:x3  1  14.539  14.539  42.298 0.0003330 ***  

x2:x3  1  14.596  14.596  42.461 0.0003291 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1  178.372 178.372 518.922 7.958e-08 ***  

x2     1  145.518 145.518 423.341 1.608e-07 ***  

x1:x2  1   0.025   0.025   0.072 0.7961658  

x1:x3  1  14.539  14.539  42.298 0.0003330 ***  

x2:x3  1  14.596  14.596  42.461 0.0003291 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.8.2 p469

(183) MODEL

```
GLM(y ~ x1 + x2 + x1:x2 + x1:x3 + x2:x3 + x1:x2:x3, polvdat) # OK
```

```
$ANOVA  

Response : y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL       6  12.5313 2.08854  37.056 0.0005473 ***  

RESIDUALS    5   0.2818 0.05636  

CORRECTED TOTAL 11 12.8131  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  R-square Coef Var Root MSE  y Mean  

0.9780061   4.391 0.2374067 5.406667
```

```
$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1  5.4668  5.4668 96.9942 0.0001839 ***  

x2     1  0.3660  0.3660  6.4944 0.0513654 .  

x1:x2  1  4.6897  4.6897 83.2068 0.0002652 ***  

x1:x3  1  1.2450  1.2450 22.0887 0.0053378 **
```

```

x2:x3      1 0.4707  0.4707  8.3509  0.0341949 *
x1:x2:x3  1 0.2931  0.2931  5.2004  0.0714991 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

x1        1 0.0184  0.0184  0.3265  0.5924707  

x2        1 0.2419  0.2419  4.2911  0.0930613 .  

x1:x2     1 3.8824  3.8824  68.8834  0.0004147 ***  

x1:x3     1 1.4383  1.4383  25.5196  0.0039276 **  

x2:x3     1 0.4707  0.4707  8.3509  0.0341949 *  

x1:x2:x3 1 0.2931  0.2931  5.2004  0.0714991 .  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

x1        1 0.25744 0.25744  4.5677  0.08562 .  

x2        1 0.12956 0.12956  2.2987  0.18992  

x1:x2     1 0.65909 0.65909 11.6939  0.01885 *  

x1:x3     1 0.26323 0.26323  4.6704  0.08307 .  

x2:x3     1 0.12999 0.12999  2.3063  0.18931  

x1:x2:x3 1 0.29310 0.29310  5.2004  0.07150 .  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.8.3 p482

(184) MODEL

```

REG(y ~ x1 + x2 + x3 + x1:x2 + x1:x3 + x2:x3 + x1:z1 + x2:z1 + x3:z1 +
     x1:x2:z1 + x1:x3:z1 + x2:x3:z1 + x1:z2 + x2:z2 + x3:z2 +
     x1:x2:z2 + x1:x3:z2 + x2:x3:z2 + x1:z1:z2 + x2:z1:z2 + x3:z1:z2 +
     x1:x2:z1:z2 + x1:x3:z1:z2 + x2:x3:z1:z2 - 1, MPV) # OK

```

	Estimate	Std. Error	Df	t value	Pr(> t)
x1	346948	294197	11	1.1793	0.2631550
x2	8223	490	11	16.7869	3.467e-09 ***
x3	1656	459	11	3.6104	0.0040950 **
x1:x2	-414463	312262	11	-1.3273	0.2113017
x1:x3	-334747	311426	11	-1.0749	0.3054382
x2:x3	-6476	1199	11	-5.4032	0.0002156 ***
x1:z1	103044	328922	11	0.3133	0.7599297
x2:z1	-2241	548	11	-4.0924	0.0017824 **
x3:z1	823	513	11	1.6056	0.1366709

```

x1:x2:z1      -64013    349120 11 -0.1834 0.8578546
x1:x3:z1      -123730   348184 11 -0.3554 0.7290412
x2:x3:z1       4659     1340 11  3.4765 0.0051806 **
x1:z2         244320   328922 11  0.7428 0.4731733
x2:z2          886      548 11  1.6187 0.1338108
x3:z2          86       513 11  0.1670 0.8704301
x1:x2:z2      -266052   349120 11 -0.7621 0.4620497
x1:x3:z2      -253151   348184 11 -0.7271 0.4823761
x2:x3:z2      -1822    1340 11 -1.3593 0.2012686
x1:z1:z2      259038   328922 11  0.7875 0.4476062
x2:z1:z2      -137     548 11 -0.2500 0.8071853
x3:z1:z2       100     513 11  0.1955 0.8485983
x1:x2:z1:z2   -269527   349120 11 -0.7720 0.4563702
x1:x3:z1:z2   -269249   348184 11 -0.7733 0.4556454
x2:x3:z1:z2   -328     1340 11 -0.2448 0.8111141
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9 Chapter 12

10.9.1 p513

(185) MODEL

```
GLM(ybar ~ A + B + C + D + E + F + G, tile) # OK
```

```
$ANOVA
Response : ybar
            Df  Sum Sq  Mean Sq F value Pr(>F)
MODEL        7 0.68737 0.098196
RESIDUALS    0 0.00000
CORRECTED TOTAL 7 0.68737
```

```
$Fitness
R-square Coef Var Root MSE ybar Mean
      1      NA      NA 0.7424626
```

```
$`Type I`
            Df  Sum Sq  Mean Sq F value Pr(>F)
A 1 0.04984 0.04984
B 1 0.01992 0.01992
C 1 0.51534 0.51534
D 1 0.01532 0.01532
E 1 0.05965 0.05965
F 1 0.00879 0.00879
G 1 0.01851 0.01851
```

```
$`Type II`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 0.04984 0.04984  
B 1 0.01992 0.01992  
C 1 0.51534 0.51534  
D 1 0.01532 0.01532  
E 1 0.05965 0.05965  
F 1 0.00879 0.00879  
G 1 0.01851 0.01851
```

```
$`Type III`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 0.04984 0.04984  
B 1 0.01992 0.01992  
C 1 0.51534 0.51534  
D 1 0.01532 0.01532  
E 1 0.05965 0.05965  
F 1 0.00879 0.00879  
G 1 0.01851 0.01851
```

(186) MODEL

```
GLM(lns2 ~ A + B + C + D + E + F + G, tile) # OK
```

```
$ANOVA  
Response : lns2  
Df Sum Sq Mean Sq F value Pr(>F)  
MODEL 7 12.305 1.7578  
RESIDUALS 0 0.000  
CORRECTED TOTAL 7 12.305
```

```
$Fitness  
R-square Coef Var Root MSE lns2 Mean  
1 NA -2.623421
```

```
$`Type I`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 1.6436 1.6436  
B 1 0.3109 0.3109  
C 1 7.1858 7.1858  
D 1 2.3199 2.3199  
E 1 0.0248 0.0248  
F 1 0.7379 0.7379  
G 1 0.0820 0.0820
```

```
$`Type II`
```

```

Df Sum Sq Mean Sq F value Pr(>F)
A 1 1.6436 1.6436
B 1 0.3109 0.3109
C 1 7.1858 7.1858
D 1 2.3199 2.3199
E 1 0.0248 0.0248
F 1 0.7379 0.7379
G 1 0.0820 0.0820

```

```

$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
A 1 1.6436 1.6436
B 1 0.3109 0.3109
C 1 7.1858 7.1858
D 1 2.3199 2.3199
E 1 0.0248 0.0248
F 1 0.7379 0.7379
G 1 0.0820 0.0820

```

10.9.2 p521

(187) MODEL

```

strng = reshape(tile,
  direction = "long",
  varying = list(c("y1", "y2")),
  v.names = "y",
  idvar = c("A", "B", "C", "D", "E", "F", "G"),
  timevar = "H",
  times = c(-1, 1))
GLM(y ~ A/H + B/H + C/H + D/H + E/H + F/H + G/H, strng) # OK

```

```

$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      14 1.65427 0.11816 0.1433 0.9807
RESIDUALS   1 0.82473 0.82473
CORRECTED TOTAL 15 2.47901

```

```

$Fitness
R-square Coef Var Root MSE    y Mean
0.667313 122.3157 0.9081486 0.7424626

```

```

$`Type I`
Df  Sum Sq Mean Sq F value Pr(>F)
A   1 0.09968 0.09968 0.1209 0.7870

```

A:H	1	0.04015	0.04015	0.0487	0.8618
B	1	0.03984	0.03984	0.0483	0.8623
H:B	1	0.00043	0.00043	0.0005	0.9854
C	1	1.03069	1.03069	1.2497	0.4646
H:C	1	0.15307	0.15307	0.1856	0.7410
D	1	0.03064	0.03064	0.0372	0.8788
H:D	1	0.04690	0.04690	0.0569	0.8510
E	1	0.11929	0.11929	0.1446	0.7686
H:E	1	0.01883	0.01883	0.0228	0.9045
F	1	0.01758	0.01758	0.0213	0.9077
H:F	1	0.01384	0.01384	0.0168	0.9180
G	1	0.03702	0.03702	0.0449	0.8671
H:G	1	0.00632	0.00632	0.0077	0.9444

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	0.09968	0.09968	0.1209	0.7870
A:H	1	0.04015	0.04015	0.0487	0.8618
B	1	0.03984	0.03984	0.0483	0.8623
H:B	1	0.00043	0.00043	0.0005	0.9854
C	1	1.03069	1.03069	1.2497	0.4646
H:C	1	0.15307	0.15307	0.1856	0.7410
D	1	0.03064	0.03064	0.0372	0.8788
H:D	1	0.04690	0.04690	0.0569	0.8510
E	1	0.11929	0.11929	0.1446	0.7686
H:E	1	0.01883	0.01883	0.0228	0.9045
F	1	0.01758	0.01758	0.0213	0.9077
H:F	1	0.01384	0.01384	0.0168	0.9180
G	1	0.03702	0.03702	0.0449	0.8671
H:G	1	0.00632	0.00632	0.0077	0.9444

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	0.09968	0.09968	0.1209	0.7870
A:H	1	0.04015	0.04015	0.0487	0.8618
B	1	0.03984	0.03984	0.0483	0.8623
H:B	1	0.00043	0.00043	0.0005	0.9854
C	1	1.03069	1.03069	1.2497	0.4646
H:C	1	0.15307	0.15307	0.1856	0.7410
D	1	0.03064	0.03064	0.0372	0.8788
H:D	1	0.04690	0.04690	0.0569	0.8510
E	1	0.11929	0.11929	0.1446	0.7686
H:E	1	0.01883	0.01883	0.0228	0.9045
F	1	0.01758	0.01758	0.0213	0.9077
H:F	1	0.01384	0.01384	0.0168	0.9180
G	1	0.03702	0.03702	0.0449	0.8671
H:G	1	0.00632	0.00632	0.0077	0.9444

10.9.3 p525

(188) MODEL

```
prod2 = af(prodstd, 1:7)
GLM(Pof ~ A + B + C + D + E + F + G + A:G + A:E:F + B:E:G + C:E:G + C:E:G:F +
    D:E + D:F, prod2) # OK
```

```
$ANOVA
Response : Pof
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      47 769.49 16.3721  5.1667 2.737e-05 ***
RESIDUALS   24  76.05  3.1688
CORRECTED TOTAL 71 845.54
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
R-square Coef Var Root MSE Pof Mean
0.9100571 9.021403 1.780098 19.73194
```

```
$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
A           2 50.577 25.288  7.9806 0.0022023 **
B           2 13.384  6.692  2.1118 0.1429491
C           2 68.594 34.297 10.8234 0.0004463 ***
D           2 23.674 11.837  3.7355 0.0386914 *
E           1 275.733 275.733 87.0165 1.878e-09 ***
F           1 161.700 161.700 51.0296 2.204e-07 ***
G           1  1.051   1.051   0.3318 0.5699896
A:G         2 26.567 13.284  4.1921 0.0274494 *
A:E:F       7 28.404  4.058  1.2806 0.3013844
B:E:G       7 22.453  3.208  1.0123 0.4475160
C:E:G       6 35.546  5.924  1.8696 0.1277692
C:E:F:G    10 24.607  2.461  0.7766 0.6500534
D:E         2 21.745 10.873  3.4312 0.0489076 *
D:F         2 15.450  7.725  2.4379 0.1086730
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
A           2 50.577 25.288  7.9806 0.0022023 **
B           2 13.384  6.692  2.1118 0.1429491
C           2 68.594 34.297 10.8234 0.0004463 ***
D           2 23.674 11.837  3.7355 0.0386914 *
E           1 275.733 275.733 87.0165 1.878e-09 ***
```

```

F      1 161.700 161.700 51.0296 2.204e-07 ***
G      1   1.051   1.051   0.3318  0.5699896
A:G    2 26.567  13.284  4.1921  0.0274494 *
A:E:F  6 24.623   4.104   1.2951  0.2970196
B:E:G  6 19.770   3.295   1.0398  0.4246194
C:E:G  6 35.546   5.924   1.8696  0.1277692
C:E:F:G 10 24.607   2.461   0.7766  0.6500534
D:E     2 21.745  10.873  3.4312  0.0489076 *
D:F     2 15.450   7.725   2.4379  0.1086730
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type III`
CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	2	50.577	25.288	7.9806	0.0022023 **
B	2	13.384	6.692	2.1118	0.1429491
C	2	68.594	34.297	10.8234	0.0004463 ***
D	2	23.674	11.837	3.7355	0.0386914 *
E	1	275.733	275.733	87.0165	1.878e-09 ***
F	1	161.700	161.700	51.0296	2.204e-07 ***
G	1	1.051	1.051	0.3318	0.5699896
A:G	2	26.567	13.284	4.1921	0.0274494 *
A:E:F	6	24.623	4.104	1.2951	0.2970196
B:E:G	6	19.770	3.295	1.0398	0.4246194
C:E:G	6	35.546	5.924	1.8696	0.1277692
C:E:F:G	10	24.607	2.461	0.7766	0.6500534
D:E	2	21.745	10.873	3.4312	0.0489076 *
D:F	2	15.450	7.725	2.4379	0.1086730

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.9.4 p532

(189) MODEL

```
GLM(torque ~ A + B + C + D + E + A:B + A:C + A:D + A:E, Smotor) # OK
```

\$ANOVA
Response : torque

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	15	0.0112217	0.00074811	102.2	0.009731 **
RESIDUALS	2	0.0000146	0.00000732		
CORRECTED TOTAL	17	0.0112363			

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

$Fitness
  R-square Coef Var      Root MSE torque Mean
  0.9986971 1.051627 0.002705567   0.2572743

$`Type I`
  Df      Sum Sq   Mean Sq   F value   Pr(>F)
A     1 0.0039545 0.0039545 540.2187 0.001846 ***
B     2 0.0003817 0.0001909 26.0732 0.036937 *
C     2 0.0057241 0.0028620 390.9837 0.002551 ***
D     2 0.0000265 0.0000133   1.8104 0.355820
E     1 0.0000984 0.0000984 13.4406 0.067009 .
A:B   2 0.0010068 0.0005034 68.7668 0.014333 *
A:C   2 0.0000031 0.0000016   0.2134 0.824110
A:D   2 0.0000009 0.0000004   0.0599 0.943521
A:E   1 0.0000258 0.0000258   3.5198 0.201458
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df      Sum Sq   Mean Sq   F value   Pr(>F)
A     1 0.0039545 0.0039545 540.2187 0.001846 ***
B     2 0.0003817 0.0001909 26.0732 0.036937 *
C     2 0.0032014 0.0016007 218.6753 0.004552 ***
D     2 0.0000268 0.0000134   1.8319 0.353123
E     1 0.0000423 0.0000423   5.7744 0.138172
A:B   2 0.0010068 0.0005034 68.7668 0.014333 *
A:C   2 0.0000031 0.0000016   0.2134 0.824110
A:D   2 0.0000052 0.0000026   0.3536 0.738760
A:E   1 0.0000258 0.0000258   3.5198 0.201458
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df      Sum Sq   Mean Sq   F value   Pr(>F)
A     1 0.0034241 0.0034241 467.7636 0.002131 ***
B     2 0.0003817 0.0001909 26.0732 0.036937 *
C     2 0.0032014 0.0016007 218.6753 0.004552 ***
D     2 0.0000268 0.0000134   1.8319 0.353123
E     1 0.0000423 0.0000423   5.7744 0.138172
A:B   2 0.0010068 0.0005034 68.7668 0.014333 *
A:C   2 0.0000031 0.0000016   0.2134 0.824110
A:D   2 0.0000052 0.0000026   0.3536 0.738760
A:E   1 0.0000258 0.0000258   3.5198 0.201458
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9.5 p535

(190) MODEL

```
GLM(shrinkage ~ A + B + C + D + E + F + G + A:B + A:C + A:D + A:E + A:F + A:G +  
      B:D, inject) # OK
```

\$ANOVA

```
Response : shrinkage  
           Df Sum Sq Mean Sq F value    Pr(>F)  
MODEL          14 6659.4  475.67  129.08 1.97e-05 ***  
RESIDUALS       5   18.4     3.68  
CORRECTED TOTAL 19 6677.8  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
R-square Coef Var Root MSE shrinkage Mean  
0.9972409 7.083525 1.919635             27.1
```

\$`Type I`

```
           Df Sum Sq Mean Sq  F value    Pr(>F)  
A      1 770.1  770.1  208.9722 2.858e-05 ***  
B      1 5076.6 5076.6 1377.6289 2.674e-07 ***  
C      1   3.1    3.1   0.8311  0.403773  
D      1   7.6    7.6   2.0522  0.211416  
E      1   0.6    0.6   0.1526  0.712112  
F      1   0.6    0.6   0.1526  0.712112  
G      1  95.1   95.1   25.7972  0.003837 **  
A:B    1 564.1  564.1  153.0699 6.112e-05 ***  
A:C    1 10.6   10.6   2.8664  0.151230  
A:D    1 115.6  115.6   31.3602  0.002508 **  
A:E    1 14.1   14.1   3.8161  0.108185  
A:F    1   1.6    1.6   0.4240  0.543677  
A:G    1   0.1    0.1   0.0170  0.901459  
B:D    1   0.1    0.1   0.0170  0.901459  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
           Df Sum Sq Mean Sq  F value    Pr(>F)  
A      1 770.1  770.1  208.9722 2.858e-05 ***  
B      1 5076.6 5076.6 1377.6289 2.674e-07 ***  
C      1   3.1    3.1   0.8311  0.403773  
D      1   7.6    7.6   2.0522  0.211416  
E      1   0.6    0.6   0.1526  0.712112  
F      1   0.6    0.6   0.1526  0.712112
```

```

G     1    95.1    95.1   25.7972  0.003837 ** 
A:B   1    564.1   564.1   153.0699 6.112e-05 *** 
A:C   1    10.6    10.6    2.8664   0.151230 
A:D   1    115.6   115.6   31.3602  0.002508 ** 
A:E   1    14.1    14.1    3.8161   0.108185 
A:F   1     1.6    1.6    0.4240   0.543677 
A:G   1     0.1    0.1    0.0170   0.901459 
B:D   1     0.1    0.1    0.0170   0.901459 

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III` 
  Df Sum Sq Mean Sq F value Pr(>F) 
  A     1  770.1  770.1  208.9722 2.858e-05 *** 
  B     1 5076.6 5076.6 1377.6289 2.674e-07 *** 
  C     1     3.1     3.1    0.8311  0.403773 
  D     1     7.6     7.6    2.0522  0.211416 
  E     1     0.6     0.6    0.1526  0.712112 
  F     1     0.6     0.6    0.1526  0.712112 
  G     1    95.1    95.1   25.7972  0.003837 ** 
  A:B   1    564.1   564.1   153.0699 6.112e-05 *** 
  A:C   1    10.6    10.6    2.8664   0.151230 
  A:D   1    115.6   115.6   31.3602  0.002508 ** 
  A:E   1    14.1    14.1    3.8161   0.108185 
  A:F   1     1.6    1.6    0.4240   0.543677 
  A:G   1     0.1    0.1    0.0170   0.901459 
  B:D   1     0.1    0.1    0.0170   0.901459 

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9.6 p539

(191) MODEL

```

eptax = cbind(eptaxr[1:16,], y2=eptaxr[17:32,9], y3=eptaxr[33:48,9], 
               y5=eptaxr[49:64,9])
eptax$ybar = (eptax$y + eptax$y2 + eptax$y3 + eptax$y5)/4
GLM(ybar ~ A + B + C + D + E + F + G + H + A:B + A:C + A:D + A:E + A:F + A:G + 
     A:H, eptax) # OK

```

```

$ANOVA
Response : ybar
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      15 2.8452 0.18968
RESIDUALS   0 0.0000
CORRECTED TOTAL 15 2.8452

```

```
$Fitness
R-square Coef Var Root MSE ybar Mean
      1       NA       NA 14.36122
```

```
$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
A   1 0.02686 0.02686
B   1 0.00042 0.00042
C   1 0.06306 0.06306
D   1 2.49443 2.49443
E   1 0.00304 0.00304
F   1 0.03209 0.03209
G   1 0.02954 0.02954
H   1 0.12879 0.12879
A:B 1 0.00047 0.00047
A:C 1 0.03218 0.03218
A:D 1 0.01185 0.01185
A:E 1 0.00380 0.00380
A:F 1 0.01674 0.01674
A:G 1 0.00186 0.00186
A:H 1 0.00012 0.00012
```

```
$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
A   1 0.02686 0.02686
B   1 0.00042 0.00042
C   1 0.06306 0.06306
D   1 2.49443 2.49443
E   1 0.00304 0.00304
F   1 0.03209 0.03209
G   1 0.02954 0.02954
H   1 0.12879 0.12879
A:B 1 0.00047 0.00047
A:C 1 0.03218 0.03218
A:D 1 0.01185 0.01185
A:E 1 0.00380 0.00380
A:F 1 0.01674 0.01674
A:G 1 0.00186 0.00186
A:H 1 0.00012 0.00012
```

```
$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
A   1 0.02686 0.02686
B   1 0.00042 0.00042
C   1 0.06306 0.06306
D   1 2.49443 2.49443
E   1 0.00304 0.00304
```

F	1	0.03209	0.03209
G	1	0.02954	0.02954
H	1	0.12879	0.12879
A:B	1	0.00047	0.00047
A:C	1	0.03218	0.03218
A:D	1	0.01185	0.01185
A:E	1	0.00380	0.00380
A:F	1	0.01674	0.01674
A:G	1	0.00186	0.00186
A:H	1	0.00012	0.00012

11 Searle - Linear Models 2e

Reference

- Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

11.1 7.2 (p390, 59%)

(192) MODEL

```
weight = c(8,13,9,12,7,11,6,12,12,14,9,7,14,16,10,14,11,13)
treatment = c("ta","ta","ta","ta","ta","tb","tb","tb","tc","tc","tc",
             "tc","tc","tc")
variety = c("va","va","va","vc","vd","vd","va","va","vb","vb","vb","vb",
           "vc","vd","vd","vd")
d1 = data.frame(weight, treatment, variety)
GLM(weight ~ treatment*variety, d1)
```

\$ANOVA

```
Response : weight
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7    82   11.714  2.0918  0.14
RESIDUALS      10    56    5.600
CORRECTED TOTAL 17   138
```

\$Fitness

```
R-square Coef Var Root MSE weight Mean
0.5942029 21.51302 2.366432           11
```

\$`Type I`

```
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 10.500   5.250  0.9375 0.42348
variety        3 36.786  12.262  2.1896 0.15232
treatment:variety  2 34.714  17.357  3.0995 0.08965 .
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2  9.486  4.7429  0.8469 0.45731
variety        3 36.786 12.2619  2.1896 0.15232
treatment:variety  2 34.714 17.3571  3.0995 0.08965 .
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```

          Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 12.471  6.2353  1.1134 0.36595
variety        3 34.872 11.6240  2.0757 0.16719
treatment:variety 2 34.714 17.3571  3.0995 0.08965 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(weight ~ treatment*variety, d1), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: weight
          Sum Sq Df F values Pr(>F)
treatment      0.000  0
variety        0.000  0
treatment:variety 34.714  2   3.0995 0.08965 .
Residuals     56.000 10
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

11.2 7.2 (p393, 60%)

(193) MODEL

```

percent = c(31,33,44,36,38,26,37,59,42,42,34,42,28,39,36,32,38,42,36,22,42,46,
          26,37,43)
refinery = c(rep("g",9),rep("n",8),rep("s",8))
process = as.factor(c(1,1,1,1,1,1,2,2,2,1,1,1,2,2,2,2,1,1,1,2,2,2,2))
source0 = c("t","t","t","t","o","m","t","t","o","m","i","i","i","t","o","m","m",
           "t","o","i","o","o","m","i","i")
d2 = data.frame(percent, refinery, process, source=source0)
GLM(percent ~ refinery*source, d2)

```

```

$ANOVA
Response : percent
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL       10 442.56  44.256  0.6361 0.7616
RESIDUALS    14 974.00  69.571
CORRECTED TOTAL 24 1416.56

```

\$Fitness

```

R-square Coef Var Root MSE percent Mean
0.3124188 22.39782 8.340949      37.24

$`Type I` 
          Df Sum Sq Mean Sq F value Pr(>F)
refinery     2 20.963 10.481 0.1507 0.8615
source       3 266.124 88.708 1.2751 0.3212
refinery:source 5 155.474 31.095 0.4469 0.8086

$`Type II` 
          Df Sum Sq Mean Sq F value Pr(>F)
refinery     2 25.535 12.767 0.1835 0.8343
source       3 266.124 88.708 1.2751 0.3212
refinery:source 5 155.474 31.095 0.4469 0.8086

$`Type III` 
          Df Sum Sq Mean Sq F value Pr(>F)
refinery     2 10.766 5.383 0.0774 0.9259
source       3 282.633 94.211 1.3542 0.2972
refinery:source 5 155.474 31.095 0.4469 0.8086

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(percent ~ refinery*source, d2), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: percent
          Sum Sq Df F values Pr(>F)
refinery     2.52  1 0.0362 0.8518
source       268.19  2 1.9275 0.1822
refinery:source 155.47  5 0.4469 0.8086
Residuals    974.00 14

```

12 Web site examples

12.1 <https://github.com/djnavarro/psyr>

(194) MODEL

```
d21 = read.csv("http://r.acr.kr/psyr/coffee.csv")
GLM(babble ~ sugar*milk - 1, d21)

$ANOVA
Response : babble
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       6 472.54  78.756  298.84 2.39e-12 ***
RESIDUALS   12   3.16   0.264
UNCORRECTED TOTAL 18 475.70
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  R-square Coef Var Root MSE babble Mean
0.9933519 10.13217 0.5133631     5.066667

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      3 465.64 155.213 588.9486 2.756e-13 ***
milk       1   0.96   0.956   3.6279  0.081061 .
sugar:milk 2   5.94   2.972  11.2769  0.001754 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      2  3.0696 1.53482  5.8238 0.017075 *
milk       1  0.9561 0.95611  3.6279  0.081061 .
sugar:milk 2  5.9439 2.97193 11.2769  0.001754 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      2  2.1318 1.0659  4.0446 0.045426 *
milk       1  1.0041 1.0041  3.8102 0.074672 .
sugar:milk 2  5.9439 2.9719 11.2769  0.001754 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts=c("contr.sum", "contr.poly"))
r21 = lm(babble ~ sugar*milk - 1, d21)
anova(r21) # Type I SS OK
```

Analysis of Variance Table

```
Response: babble
          Df Sum Sq Mean Sq F value    Pr(>F)
sugar      3 465.64 155.213 588.9486 2.756e-13 ***
milk       1   0.96   0.956   3.6279  0.081061 .
sugar:milk 2   5.94   2.972  11.2769  0.001754 **
Residuals 12   3.16   0.264
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Anova(r21, type=2) # NOT OK
```

Anova Table (Type II tests)

```
Response: babble
          Sum Sq Df F value    Pr(>F)
sugar      453.76  3 573.9233 3.214e-13 ***
milk        0.96  1   3.6279  0.081061 .
sugar:milk  5.94  2  11.2769  0.001754 **
Residuals   3.16 12
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Anova(r21, type=3) # NOT OK
```

Anova Table (Type III tests)

```
Response: babble
          Sum Sq Df F value    Pr(>F)
sugar      454.77  3 575.1970 3.172e-13 ***
milk        1.00  1   3.8102  0.074672 .
sugar:milk  5.94  2  11.2769  0.001754 **
Residuals   3.16 12
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

13 Test Summary

Package	Version	Total Count	Identical to SAS	Different from SAS
sasLM	0.8.0	194	194 (100%)	0 (0%)
car	3.0.12	194	173 (89%)	21 (11%)

All of the results by sasLM 0.8.0 were practically identical to those of SAS.

Last digit difference by 1 is resulted from the round-to-even number way of R rounding function.

If you are uncertain about the equivalence of the ‘sasLM’ to ‘SAS,’ you can check these examples using ‘SAS onDemand’ for free.

If you have any question, please mail to the author, Kyun-Seop Bae k@acr.kr.

14 Session Information

```
R version 4.1.3 (2022-03-10)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 10 x64 (build 19044)
```

```
Matrix products: default
```

```
locale:
```

```
[1] LC_COLLATE=Korean_Korea.949  LC_CTYPE=Korean_Korea.949
[3] LC_MONETARY=Korean_Korea.949 LC_NUMERIC=C
[5] LC_TIME=Korean_Korea.949
```

```
attached base packages:
```

```
[1] stats      graphics   grDevices utils      datasets   methods    base
```

```
other attached packages:
```

```
[1] daewr_1.2-7     car_3.0-12      carData_3.0-5   sasLM_0.8.0    mvtnorm_1.1-3
[6] rmarkdown_2.13
```

```
loaded via a namespace (and not attached):
```

```
[1] gmp_0.6-5          compiler_4.1.3      mathjaxr_1.6-0
[4] numbers_0.8-2       tools_4.1.3        partitions_1.10-4
[7] digest_0.6.29       evaluate_0.15     lattice_0.20-45
[10] pkgconfig_2.0.3      rlang_1.0.2        igraph_1.2.11
[13] cli_3.2.0          yaml_2.3.5        polynom_1.4-0
[16] xfun_0.30          fastmap_1.1.0     stringr_1.4.0
[19] knitr_1.37          combinat_0.0-8    lmtest_0.9-39
[22] vcd_1.4-9           grid_4.1.3        scatterplot3d_0.3-41
[25] DoE.base_1.2         conf.design_2.0.0   FrF2_2.2-2
[28] magrittr_2.0.2       htmltools_0.5.2    MASS_7.3-55
[31] sfsmisc_1.1-12       abind_1.4-5       colorspace_2.0-3
[34] tinytex_0.37         stringi_1.7.6     zoo_1.8-9
```